IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the application of: R. Rogers Yocum et al.

Serial No.: Not Yet Assigned

Filed: Herewith

Methods and Microorganisms for

Production of Panto-Compounds

Attorney Docket No.: BBI-141CP

BOX PATENT APPLICATION **Assistant Commissioner for Patents** Washington, D.C. 20231

CERTIFICATION	UNDER 37	CFR 1.10
---------------	----------	----------

September 21, 2000 Date of Deposit:

Mailing Label Number: EL 095 525 893 US

I hereby certify that this 37 CFR 1.53(d) request and the documents referred to therein as enclosed are being deposited with the United States Postal Service on the date indicated above in an envelope as "Express Mail Post Office to Addressee" service under 37 CFR 1.10 and addressed to the Box Patent Application, Assistant Commissioner for Patents, Washington, D.C. 20231.

William J. McKinney

Name of Person Mailing Paper

Signature of Person Mailing Paper

PRELIMINARY AMENDMENT

Dear Sir:

Prior to examination of the above-identified application, please amend the application as follows:

In the Claims:

Please cancel claim 3-6, 8-11, 13, 45, 54-61, 63-66, 76-81, 84-87, 89, 91-96, 100-101, 103, 105 and 107, without prejudice.



Please amend claims 12, 20-25, 27-28, 33 and 46-50, as follows:

- 12. (Amended) The method of [any one of claims 7 to 11] <u>claim 7</u>, wherein the KPAR-O microorganism further overexpresses at least one pantothenate biosynthetic enzyme in addition to overexpressing ketopantoate reductase.
- 20. (Amended) The method of [any one of claims 14 to 19] <u>claim 14 or 19</u>, wherein the microorganism overexpresses acetohydroxyacid synthetase or is transformed with a vector comprising an *ilvBN* nucleic acid sequence or an *alsS* sequence.
- 21. (Amended) The method of [any one of claims 14 to 19] claim 14 or 19, wherein the microorganism overexpresses acetohydroxyacid isomeroreductase or is transformed with a vector comprising an *ilvC* nucleic acid sequence.
- 22. (Amended) The method of [any one of claims 14 to 19] <u>claim 14 or 19</u>, wherein the microorganism overexpresses dihydroxyacid dehydratase or is transformed with a vector comprising an *ilvD* nucleic acid sequence.
- 23. (Amended) The method of [any one of claims 19 to 22] <u>claim 19</u>, wherein the microorganism overexpresses aspartate- α -decarboxylase or is transformed with a vector comprising a *panD* nucleic acid sequence.
- 24. (Amended) The method of [any one of claims 14 to 19] <u>claim 14 or 19</u>, wherein the microorganism further has a deregulated pantothenate biosynthetic pathway.
- 25. (Amended) The method of [any one of claims 14 to 19] claim 14 or 19, wherein the microorganism further has at least one mutant gene selected from the group consisting of a mutant *avtA* gene, a mutant *ilvE* gene, a mutant *ansB* gene and a mutant *alsD* gene.
- 27. (Amended) The method of claim 24 [or 26], wherein the microorganism is transformed with a vector comprising a *panBCD* nucleic acid sequence or a vector comprising a *panE1* nucleic acid sequence.

- 28. (Amended) The method of [any one of claims 14 to 19] <u>claim 14 or 19</u>, wherein pantothenate is produced at a level selected from the group consisting of a level greater than 10g/L, a level greater than 20g/L and a level greater than 40g/L.
- 33. (Amended) The method of claim 24 [or 26], wherein the microorganism overexpresses any of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase and aspartate- α -decarboxylase derived from *Bacillus*.
- 46. (Amended) The method of [any one of claims 39 to 44] <u>claim 39 or 41</u>, wherein said panto-compound is pantothenate.
- 47. (Amended) The method of [any one of claims 39 to 44] <u>claim 39 or 41</u>, wherein said panto-compound is produced at a level selected from the group consisting of a level greater than 10g/L, a level greater than 20g/L and a level greater than 40g/L.
- 48. (Amended) The method of [any one of claims 39 to 44] <u>claim 39 or 41</u>, wherein said recombinant microorganism further has a deregulated pantothenate biosynthetic pathway or further has a deregulated isoleucine-valine (*ilv*) biosynthetic pathway.
- 49. (Amended) The method of [any one of claims 39 to 44] <u>claim 39 or 41</u>, wherein said recombinant microorganism further overexpresses *panD* and *panE*.
- 50. (Amended) The method of [any one of claims 39 to 44] claim 39 or 41, wherein said recombinant microorganism further has at least one mutant gene selected from the group consisting of a mutant avtA gene, a mutant ilvE gene, a mutant ansB gene and a mutant alsD gene.

THE RESERVE THE PARTY OF THE STATE OF THE ST

REMARKS

Applicants have attached herewith APPENDIX A setting forth the claims that will be pending after entry of the instant amendment. No new matter has been added to the application.

Date: September 21, 2000

LAHIVE & COCKFIELD, LLP

Attorneys at Law

Debra J. Milasincic, Esq.

Reg. No. 46,931 28 State Street Boston, MA 02109 (617) 227-7400

(617) 742-4214

APPENDIX A

- 1. A method of producing a panto-compound comprising culturing a microorganism which overexpresses at least one *Bacillus* pantothenate biosynthetic enzyme under conditions such that the panto-compound is produced.
- 2. The method of claim 1, wherein the microorganism overexpresses at least one *Bacillus subtilis* pantothenate biosynthetic enzyme.
- 7. A method of producing a panto-compound comprising culturing a ketopantoate reductase-overexpressing (KPAR-O) microorganism under conditions such that the panto-compound is produced.
- 12. The method of claim 7, wherein the KPAR-O microorganism further overexpresses at least one pantothenate biosynthetic enzyme in addition to overexpressing ketopantoate reductase.
- 14. A method of producing pantothenate in a manner independent of precursor feed comprising culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism having a deregulated isoleucine-valine (ilv) pathway under conditions such that pantothenate is produced.
- 15. A method of producing at least 2 g/L pantothenate in a manner independent of aspartate or β -alanine feed comprising culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism under conditions such that pantothenate is produced.
- 16. A method of producing at least 2 g/L pantothenate in a manner independent of valine or α -ketoisovalerate feed comprising culturing a microorganism having a deregulated isoleucine-valine (ilv) biosynthetic pathway under conditions such that pantothenate is produced.
- 17. A method of producing at least 30 g/L pantothenate in a manner independent of aspartate or β -alanine feed comprising culturing an aspartate- α -

decarboxylase-overexpressing (A α D-O) microorganism under conditions such that pantothenate is produced.

- 18. A method of producing at least 30 g/L pantothenate in a manner independent of valine or α -ketoisovalerate feed comprising culturing a microorganism having a deregulated isoleucine-valine (ilv) biosynthetic pathway under conditions such that pantothenate is produced.
- 19. A β -alanine independent high yield production method for producing pantothenate comprising culturing a manipulated microorganism under conditions such that pantothenate is produced at a significantly high yield.
- 20. The method of claim 14 or 19, wherein the microorganism overexpresses acetohydroxyacid synthetase or is transformed with a vector comprising an *ilvBN* nucleic acid sequence or an *alsS* sequence.
- 21. The method of claim 14 or 19, wherein the microorganism overexpresses acetohydroxyacid isomeroreductase or is transformed with a vector comprising an *ilvC* nucleic acid sequence.
- 22. The method of claim 14 or 19, wherein the microorganism overexpresses dihydroxyacid dehydratase or is transformed with a vector comprising an *ilvD* nucleic acid sequence.
- 23. The method of claim 19, wherein the microorganism overexpresses aspartate- α -decarboxylase or is transformed with a vector comprising a *panD* nucleic acid sequence.
- 24. The method of claim 14 or 19, wherein the microorganism further has a deregulated pantothenate biosynthetic pathway.
- 25. The method of claim 14 or 19, wherein the microorganism further has at least one mutant gene selected from the group consisting of a mutant *avtA* gene, a mutant *ilvE* gene, a mutant *ansB* gene and a mutant *alsD* gene.

- 26. The method of claim 24, wherein the microorganism overexpresses any of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase and aspartate-α-decarboxylase.
- 27. The method of claim 24, wherein the microorganism is transformed with a vector comprising a *panBCD* nucleic acid sequence or a vector comprising a *panE1* nucleic acid sequence.
- 28. The method of claim 14 or 19, wherein pantothenate is produced at a level selected from the group consisting of a level greater than 10g/L, a level greater than 20g/L and a level greater than 40g/L.
- 29. The method of claim 20, wherein the microorganism overexpresses acetohydroxyacid synthetase derived from *Bacillus* or is transformed with a vector comprising an *ilvBN* nucleic acid sequence or an *alsS* nucleic acid sequence derived from *Bacillus*.
- 30. The method of claim 21, wherein the microorganism overexpresses acetohydroxyacid isomeroreductase derived from *Bacillus* or is transformed with a vector comprising an *ilvC* nucleic acid sequence derived from *Bacillus*.
- 31. The method of claim 22, wherein the microorganism overexpresses dihydroxyacid dehydratase derived from *Bacillus* or is transformed with a vector comprising av *ilvD* nucleic acid sequence derived from *Bacillus*.
- 32. The method of claim 23, wherein the microorganism overexpresses aspartate-α-decarboxylase derived from *Bacillus* or is transformed with a vector comprising a *panD* nucleic acid sequence derived from *Bacillus*.
- 33. The method of claim 24 [or 26], wherein the microorganism overexpresses any of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase and aspartate-α-decarboxylase derived from *Bacillus*.

- 34. The method of claim 27, wherein the vector comprises a *panBCD* nucleic acid sequence or a *panE1* nucleic acid sequence derived from *Bacillus*.
- 35. A method of producing a panto-compound comprising contacting a composition comprising at least one pantothenate biosynthesis pathway precursor or isoleucine-valine biosynthesis pathway precursor with at least one isolated *Bacillus* enzyme selected from the group consisting of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase and aspartate- α -decarboxylase, under conditions such that the panto-compound is produced.
- 36. A method of producing β -alanine comprising culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism under conditions such that β -alanine is produced.
- 37. The method of claim 36, wherein the $A\alpha D$ -O microorganism has a mutation in a nucleic acid sequence encoding a pantothenate biosynthetic enzyme selected from the group consisting of ketopantoate hydroxymethyltransferase, ketopantoate reductase and pantothenate synthetase.
- 38. A method of producing β -alanine comprising contacting a composition comprising aspartate with an isolated *Bacillus* aspartate- α -decarboxylase enzyme under conditions such that β -alanine is produced.
- 39. A method for enhancing production of a panto-compound comprising culturing a mutant microorganism having a mutant *coaX* gene under conditions such that the panto-compound production is enhanced.
- 40. The method of claim 39, wherein said recombinant microorganism has a mutant *coaA* gene.
- 41. A method of producing a panto-compound comprising a pantothenate kinase mutant microorganism under conditions such that the panto-compound is produced at a significantly high yield.

- 42. The method of claim 41, wherein said mutant microorganism has a mutant *coaA* gene.
- 43. The method of claim 41, wherein said mutant microorganism has a mutant *coaX* gene.
- 44. The method of claim 41, where said mutant microorganism has a mutant *coaA* and *coaX* gene.
- 46. The method of claim 39 or 41, wherein said panto-compound is pantothenate.
- 47. The method of claim 39 or 41, wherein said panto-compound is produced at a level selected from the group consisting of a level greater than 10g/L, a level greater than 20g/L and a level greater than 40g/L.
- 48. The method of claim 39 or 41, wherein said recombinant microorganism further has a deregulated pantothenate biosynthetic pathway or further has a deregulated isoleucine-valine (*ilv*) biosynthetic pathway.
- 49. The method of claim 39 or 41, wherein said recombinant microorganism further overexpresses *panD* and *panE*.
- 50. The method of claim 39 or 41, wherein said recombinant microorganism further has at least one mutant gene selected from the group consisting of a mutant avtA gene, a mutant ilvE gene, a mutant ansB gene and a mutant alsD gene.
- 51. A method for enhancing production of a panto-compound comprising culturing a microorganism that has a deregulated pantothenate biosynthetic pathway and that also has a mutation that results in reduced pantothenate kinase activity under conditions such that the panto-compound production is enhanced.
- 52. A method for identifying compounds which modulate pantothenate kinase activity comprising contacting a recombinant cell expressing pantothenate kinase

encoded by the *coaX* gene with a test compound and determining the ability of the test compound to modulate pantothenate kinase activity in said cell.

- 53. The method of claim 52, wherein said cell further comprises a mutant *coaA* gene encoding a pantothenate kinase having reduced activity.
- 62. A recombinant microorganism which overexpresses at least one *Bacillus* pantothenate biosynthetic enzyme.
- 67. A recombinant microorganism having a mutant *coaX* gene, said mutant *coaX* gene encoding reduced pantothenate kinase activity in said microorganism.
- 68. The recombinant microorganism of claim 67 further having a mutant *coaA* gene, said mutant *coaA* gene encoding reduced pantothenate kinase activity in said microorganism.
- 69. A recombinant microorganism having a mutant *coaX* gene and optionally having a mutant *coaA* gene, said mutant microorganism having reduced pantothenate kinase activity as compared to a microorganism having wild-type *coaA* and *coaX* genes.
- 70. A recombinant microorganism comprising a vector comprising an isolated *coaX* gene.
- 71. A recombinant microorganism that overproduces a panto-compound, the microorganism having a deregulated pantothenate biosynthetic pathway and having at least one mutation that results in a decrease in the capacity of the microorganism to synthesize Coenzyme A (CoA).
- 72. The recombinant microorganism of claim 71, having at least one mutation that results in a reduced level of pantothenate kinase activity.
- 73. The recombinant microorganism of claim 72, having a mutation in a *coaA* gene, or homologue thereof, that results in a reduced level of CoaA enzyme activity.

- 74. The recombinant microorganism of claim 72, having a mutation in a *coaX* gene, or homologue thereof, that results in a reduced level of CoaX enzyme activity.
- 75. The recombinant microorganism of claim 72, having a mutation in a *coaA* gene, or homologue thereof, and having a mutation in a *coaX* gene, or homologue thereof, the mutations resulting in reduced levels of CoaA enzyme activity and reduced CoaX enzyme activity.
- 82. A recombinant microorganism selected from the group consisting of PA221, PA235, PA236, PA313, PA410, PA402, PA403, PA411, PA412, PA413, PA303, PA327, PA328, PA401, PA340, PA342, PA404, PA405, PA374, PA354, PA365, PA377, PA651 and PA824.
- 83. A recombinant vector for use in the production of panto-compounds comprising a nucleic acid sequence which encodes at least one *Bacillus* pantothenate biosynthetic enzyme operably linked to regulatory sequences.
- 88. A vector comprising a mutant *coaX* gene, said mutant encoding a pantothenate kinase enzyme having reduced activity.
 - 90. A vector comprising an isolated *Bacillus coaX* gene.
- 97. A vector selected from the group consisting of pAN004, pAN005, pAN006, pAN236, pAN423, pAN428, pAN429, pAN441, pAN442, pAN443, pAN251, pAN267, pAN256, pAN257, pAN263, pAN240, pAN294, pAN296, pAN336, pAN341 and pAN342.
- 99. An isolated nucleic acid molecule which encodes at least one *Bacillus* pantothenate biosynthetic gene.
 - 102. An isolated *Bacillus* pantothenate biosynthetic enzyme polypeptide.
 - 104. An isolated *Bacillus* ketopantoate reductase polypeptide.

- 106. An isolated *Bacillus* aspartate-α-decarboxylase polypeptide.
- 108. An isolated nucleic acid molecule comprising a mutant coaX gene.
- 109. An isolated nucleic acid molecule comprising a *coaX* gene.
- 110. An isolated pantothenate kinase protein encoded by a *coaX* gene.

10

15

20

25

30

35

.1

THE WEST OF

METHODS AND MICROORGANISMS FOR PRODUCTION OF PANTO-COMPOUNDS

Related Applications

The instant application is a continuation-in-part of U.S. Patent Application Serial No. 09/400,494, filed September 21, 1999 (pending). The instant application also claims the benefit of prior filed provisional U.S. Patent Application Serial No. 60/210,072, filed June 7, 2000, prior filed provisional U.S. Patent Application Serial No. 60/221,938, filed July 28, 2000 and prior filed provisional U.S. Patent Application Serial No. 60/227,860, filed August 24, 2000. The entire content of the above-referenced patent applications is incorporated herein by this reference.

Background of the Invention

Pantothenate, also known as pantothenic acid or vitamin B5, is a member of the B complex of vitamins and is a nutritional requirement for mammals, including livestock and humans (e.g., from food sources, as a water soluble vitamin supplement or as a feed additive). In cells, pantothenate is used primarily for the biosynthesis of coenzyme A (CoA) and acyl carrier protein (ACP). These coenzymes function in the metabolism of acyl moieties which form thioesters with the sulfhydryl group of the 4'phosphopantetheine portion of these molecules. These coenzymes are essential in all cells, participating in over 100 different intermediary reactions in cellular metabolism.

The conventional means of synthesizing pantothenate (in particular, the bioactive D isomer) is via chemical synthesis from bulk chemicals, a process which is hampered by excessive substrate cost as well as the requirement for optical resolution of racemic intermediates (e.g., resolution of DL-pantolactone to obtain D-pantolactone for chemical condensation with β -alanine). Accordingly, researchers have recently looked to bacterial or microbial systems that produce enzymes useful in pantothenate biosynthesis processes (as bacteria are themselves capable of synthesizing pantothenate). In particular, bioconversion processes have been evaluated as a means of favoring production of the D isomer of pantothenic acid, e.g., using microorganisms which selectively hydrolyze a DL-pantothenic acid ester to D-pantothenic acid; microorganisms which selectively decompose L-pantolactone resulting in Dpantolactone alone; and microorganisms which selectively hydrolyze DL-pantolactone to D-pantoic acid.

There is still, however, significant need for improved pantothenate production processes, in particular, for processes requiring reduced quantities of substrates and/or less expensive substrates. To this end, methods of direct microbial

20

synthesis have recently been examined as a means of improving D-pantothenate production. In microbes, pantothenate biosynthetis is a multistep pathway resulting in condensation of pantoate (derived from α -ketoisovalerate) and β -alanine to form D-pantothenate. The isoleucine-valine (ilv) pathway biosynthetic enzymes,

acetohydroxyacid synthetase (the *ilvBN* or *alsS* gene product), acetohydroxyacid isomeroreductase (the *ilvC* gene product) and dihydroxyacid dehydratase (the *ilvD* gene product) catalyze the conversion of pyruvate to α-ketoisovalerate. The reactions are further catalyzed by the pantothenate (*pan*) pathway biosynthetic enzymes ketopantoate hydroxymethyltransferase (the *panB* gene product), ketopantoate reductase (the *panE* gene product), aspartate-α-decarboxylase (the *panD* gene product) and pantothenate synthetase (the *panC* gene product).

The genes encoding the enzymes involved in the biosynthesis of pantothenic acid in *Salmonella typhimurium* and *Escherichia coli* have recently been identified and characterized (Frodyma and Downs (1998) *J. Biol. Chem.* 273:5572-5576 and Jackowski (1996) pp. 687-694, *In* Neidhardt *et al* (ed.) *Escherichia coli* and *Salmonella*: Cellular and Molecular Biology, 2^{nd} ed. *Am. Soc. Microbiol.* Wash, D.C). In *E. coli*, for example, the biosynthesis of pantothenic acid consists of four key steps. The first reaction is catalyzed by the *panB* gene product, ketopantoate hydroxymethyltransferase, and uses the L-valine intermediate α -ketoisovalerate to generate ketopantoate, which is subsequently reduced to pantoate by the *panE* gene product, ketopantoate reductase. The *panD* gene product, aspartate- α -decarboxylase, generates β -alanine from aspartate. The *panC* gene product, pantothenate synthetase, subsequently ligates β -alanine with pantoate to yield D-pantothenate.

The authors Dusch *et al.* described the identification of the *Corynebacterium glutamicum panD* gene and reported that expression of the *C. glutamicum panD* gene in *E. coli* yielded a strain producing pantothenate with a specific productivity of 140 ng of pantothenate per mg (dry weight) per hour. (Dusch *et al.* (1999) *Appl. Environ. Microbiol.* 65:1530-1539).

The authors Sahm and Eggeling have further identified the

Corynebacterium glutamicum panB and pan C genes and have described a genetically engineered strain of C. glutamicum which overexpresses the panBC genes (Sahm and Eggeling (1999) Appl. Environ. Microbiol. 65:1973-1979). The engineered strain produces pantothenate, however, it was necessary to overexpress the genes responsible for α-ketoisovalerate production in the host organism in order that pantothenic acid production could be detected. Moreover, without the addition of β-alanine, no substantial amounts of pantothenate accumulated with the strain constructed.

20

25

30

35

Likewise, a method of producing D-pantothenic acid has been described that takes advantage of a sodium salicylate resistant mutant strain of *E. coli* which produces D-pantothenic acid when cultured in the presence of β-alanine (U.S. Patent No. 5,518,906). Generation of *E. coli* strains resistant to α-ketoisovaleric acid and/or α-5 ketobutyric acid, and/or α-aminobutyric acid, and/or β-hydroxyaspartic acid and/or O-methyl-threonine, in addition to salicylic acid, further increased pantothenic acid production. Moreover, transformation of a plasmid DNA carrying the *panB*, *panC* and *panD* genes into the salicylic acid resistant mutant strain resulted in increased pantothenate production, however, up to 20 g/L β-alanine or more was fed in the examples given. The *panB-panC-panD* genes are clustered on the *E. coli* chromosome.

Finally, a method of producing D-pantothenic acid has been described which utilizes a salicylic acid-resistant, α -ketoisovalerate-resistant, α -ketobutyrate-resistant, β -hydroxyaspartate-resistant, o-methylthreonine-resistent *E. coli* strain transformed with pantothenate biosynthesis gene-containing DNA fragments and/or branched amino acid biosynthesis gene-containing DNA fragments and cultured in the presence of β -alanine (U.S. Patent No. 5,932,457).

Pantothenate production in bacteria results from the condensation of pantoate and β -alanine and involves the pantothenate biosynthetic enzymes ketopantoate hydroxymethyltransferase (the panB gene product), ketopantoate reductase (the panE gene product), aspartate- α -decarboxylase (the panD gene product) and pantothenate synthetase (the panC gene product). Although pantothenate is biologically active as a vitamin, it is further metabolized in all cells to Coenzyme A (CoA) which participates as an acyl group carrier in the tricarboxylic acid (TCA) cycle, fatty acid metabolism and numerous other reactions of intermediary metabolism. The initial (and possibly ratecontrolling) step in the conversion of pantothenate to Coenzyme A (CoA) is phosphorylation of pantothenate by pantothenate kinase. A pantothenate kinase activity was first identified in Salmonella typhimurium by screening for temperature-sensitive mutants which synthesized CoA at permissive temperatures but excreted pantothenate at non-permissive temperatures. The mutations were mapped in the Salmonella chromosome and the genetic locus was designated coaA. The gene encodes the enzyme that catalyzes the first step in the biosynthesis of coenzyme A from pantothenate (Dunn and Snell (1979) J. Bacteriol. 140:805-808). Escherichia coli temperature sensitive mutants have also been isolated and characterized (Vallari and Rock (1987) J. Bacteriol. 169:5795-5800). These mutants (named coaA15(Ts)) are defective in the conversion of pantothenate to CoA and further exhibit a temperature-sensitive growth phenotype, indicating that pantothenate kinase activity is essential for growth. Moreover, it was

The state of the s

::

The state of the s

noted that CoA inhibited pantothenate kinase activity to the same degree in the mutant as compared to the wild-type enzyme.

- 4 -

Feedback resistant *E. coli* mutants (named *coaA16(Fr)*) have also been isolated that posses a pantothenate kinase activity that is refractory to feedback inhibition by CoA (Vallari and Jackowski (1988) *J. Bacteriol.* 170:3961-3966). The mutation responsible for the reversion is, suprisingly, not genetically linked to the *coaA* gene by transduction. Additional data described therein support the view that the total cellular CoA content is controlled by both modulation of biosynthesis at the pantothenate kinase step and possibly by degradation of CoA to 4'-phosphopantetheine.

The wild-type E. coli coaA gene was cloned by functional 10 complementation of E. coli temperature-sensitive mutants. The sequence of the wildtype gene was determined (Song and Jackowski (1992) J. Bacteriol. 174:6411-6417 and Flamm et al. (1988) Gene (Amst.) 74:555-558). Strains containing multiple copies of the coaA gene possessed 76-fold higher specific activity of pantothenate kinase, however, there was only a 2.7-fold increase in the steady state level of CoA (Song and 15 Jackowski, supra). It has further been reported that the prokaryotic enzyme (encoded by coaA in E.coli and a variety of other microorganisms) is feedback inhibited by CoA both in vivo and in vitro with CoA being about five times more potent than acetyl-CoA in inhibiting the enzyme (Song and Jackowski, supra and Vallari et al., supra). Moreover, it has been reported that the panB gene product in E. coli is inhibited by CoA (Powers 20 and Snell (1976) J. Biol. Chem. 251:3786-3793). These data further support the view that feedback inhibition of pantothenate kinase activity is a critical factor controlling intracellular CoA concentration.

Using standard search and alignment tools, coaA homologues have been identified in Hemophilus influenzae, Mycobacterium tuberculosis, Vibrio cholerae, 25 Streptococcus pyogenes and Bacillus subtilis. By contrast, proteins with significant similarity could not be identified in eukaryotic cells including Saccharomyces cerevisiae or in mammalian expressed sequence tag (EST) databases. Using a genetic selection strategy, a cDNA encoding pantothenate kinase activity has recently been identified from Aspergillus nidulans (Calder et al. (1999) J. Biol. Chem. 274:2014-2020). The 30 eukaryotic pantothenate kinase gene (panK) has distinct primary structure and unique regulatory properties that clearly distinguish it from its prokaryotic counterpart. A mammalian pantothenate kinase gene $(mpanK1\alpha)$ has also been isolated which encodes a protein having homology to the A. nidulans PanK protein and to the predicted gene product of GenBankTM Accession Number 927798 identified in the S. cerevisiae 35 genome (Rock et al. (2000) J. Biol. Chem. 275:1377-1383).

Summary of the Invention

5

10

15

20

25

30

35

The present invention is based, at least in part, on the discovery of key enzyme-encoding genes of the pantothenate biosynthetic pathway in *Bacillus subtilis*. In particular, the present inventors have identified the *panE* gene of *B. subtilis*. Overexpression or deregulation of the *panE* gene in *B. subtilis* results in enhanced production of the *panE* gene product, ketopantoate reductase, further resulting in increased production of pantothenate. Likewise, mutations in this gene reduce pantothenate production in *B. subtilis* >90%. The present inventors have further identified the presumptive *panBCD* operon in *B. subtilis*, overexpression or deregulation of which results in increased pantothenate production. The present inventors have further demonstrated that overexpression or deregulation of the *panD* gene in *B. subtilis* (resulting in enhanced production of the *panD* gene product, aspartate-α-decarboxylase) further results in increased production of pantothenate, in particular, in combination with deregulation of genes encoding key enzymes of the isoleucine-valine (*ilv*) biosynthetic pathway.

Accordingly, the present invention features methods of producing pantothenate, as well as other compounds of the pantothenate biosynthetic pathway (e.g., ketopantoate, pantoate and β-alanine), termed "panto-compounds" herein, using microorganisms in which the pantothenate biosynthetic pathway and/or isoleucinevaline biosynthetic pathway has been manipulated such that pantothenate or other desired panto-compounds are produced. In one embodiment, the invention features a method of producing a panto-compound (e.g., pantothenate or pantoate) that involves culturing a microorganism which overexpresses the panE gene product, ketopantoate reductase, also referred to herein as a ketopantoate reductase-overexpressing or "KPAR-O" microorganism, under conditions such that the panto-compound (e.g., pantothenate or pantoate) is produced. In another embodiment, the present invention features a method of producing panto-compounds (e.g., pantothenate or pantoate) which includes culturing a microorganism which overexpresses at least one pantothenate biosynthetic enzyme (e.g., at least one of the panB, panC or panD gene products), preferably in a KPAR-O microorganism, under conditions such that the panto-compound (e.g., pantothenate or pantoate) is produced.

Yet another aspect of the invention features methods of producing panto-compounds which are independent of the need to feed precursors (e.g., β -alanine or aspartate and/or α -ketoisovalerate or valine). In one embodiment, the invention features a method of producing pantothenate in a manner independent of precursor feed that includes culturing an aspartate- α -decarboxylase-overexpressing (A α D-O)

10

15

microorganism having a deregulated isoleucine-valine (ilv) pathway under conditions such that pantothenate is produced. In another embodiment, the invention features a method of producing pantothenate in a manner independent of precursor feed that includes culturing an AaD-O microorganism having a deregulated pantothenate (pan) pathway and a deregulated isoleucine-valine (ilv) pathway, under conditions such that pantothenate is produced. In another embodiment, the invention features a method of producing pantothenate in a manner independent of aspartate or β-alanine feed that includes culturing an AaD-O microorganism under conditions such that pantothenate is produced. In another embodiment, the invention features a method of producing pantothenate in a manner independent of valine or α -ketoisovalerate feed that includes culturing a microorganism having a deregulated isoleucine-valine (ilv) biosynthetic pathway under conditions such that pantothenate is produced. In yet another embodiment, the invention features a high yield production method for producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a significantly high yield (e.g., at a level greater than 10 g/L, 20 g/L, 30 g/L or 40g/L).

The methods of the present invention further feature microorganisms that overexpresses acetohydroxyacid synthetase or acetohydroxyacid isomeroreductase (e.g., microorganisms transformed with a vector that includes an ilvBNC nucleic acid sequence), microorganisms that overexpresses dihydroxyacid dehydratase (e.g., 20 microorganisms transformed with a vector that includes an ilvD nucleic acid sequence), microorganisms that overexpresses aspartate-α-decarboxylase (e.g., microorganisms transformed with a vector that includes a panD nucleic acid sequence), microorganisms having a deregulated isoleucine-valine (ilv) biosynthetic pathway and microorganisms having a deregulated pantothenate biosynthetic pathway (e.g., microorganisms that 25 overexpress any of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase and aspartate- α -decarboxylase, for example, microorganisms transformed with a vector comprising a panBCD nucleic acid sequence or a vector comprising a panE1 nucleic acid sequence). In one embodiment, the recombinant microorganism is Gram positive (e.g., microorganisms belonging to the genus Bacillus, 30 Cornyebacterium, Lactobacillus, Lactococci or Streptomyces). In another embodiment, the recombinant microorganism is Gram negative. Particularly preferred is a Bacillus recombinant microorganism (e.g., Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus subtilis, Bacillus pumilus, Bacillus halodurans, and the like). Recombinant vectors that contain the genes encoding Bacillus pantothenate and/or isoleucine-valine 35 biosynthetic enzymes (e.g., B. subtilis pantothenate and/or isoleucine-valine biosynthetic enzymes) are also described.

10

15

20

25

30

35

Also featured are methods of producing β -alanine that include culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism under conditions such that β -alanine is produced and methods of producing β -alanine that involve contacting a composition comprising aspartate with an isolated *Bacillus* aspartate- α -decarboxylase enzyme under conditions such that β -alanine is produced.

The production methods of the present invention further can include recovering the panto-compound (e.g., pantothenate or pantoate).

The present invention further features recombinant microorganisms (e.g., AαD-O microorganisms, microorganisms having a deregulated isoleucine-valine (ilv) pathway, microorganisms overexpressing at least one of ketopantoate hydroxymethyltransferase (the panB gene product), pantothenate synthetase (the panC gene product), aspartate-α-decarboxylase (the panD gene product), ketopantoate reductase (the panE1 gene product) and microorganisms having a deregulated panBCD operon. Also featured are panB, panC, panD, panE, ilvB, ilvN, alsS, ilvC, and/or ilvD nucleic acid molecules, as well as vectors including such nucleic acid molecules and gene products encoded by such nucleic acid molecules.

The methodology of the present invention further includes, for example in addition to overexpressing at least one pantothenate biosynthetic enzyme, deleting or mutating a second pantothenate biosynthetic enzyme, said second pantothenate biosynthetic enzyme preferably being downstream of the desired product in the pantothenate biosynthetic pathway. For example, mutating panC, in addition to overexpressing the panE gene product, results in even further enhanced or increased production of pantoate. Accordingly, in one embodiment, the invention features a method of producing pantoate which includes culturing a microorganism which overexpresses the panE gene product and which has a deletion in the panC gene. In another embodiment, the invention features a method of producing pantoate which includes culturing a microorganism which overexpresses the panE gene product and/or panB gene product and which has a deletion in the panC gene. Other exemplary embodiments include a method of producing ketopantoate which includes culturing a microorganism which overexpresses the panB gene product and which has a deletion in the panE gene and a method of producing β-alanine which includes culturing a microorganism which overexpresses the panD gene product and which has a deletion in the panC gene. Also included are methods of producing panto-compounds which include overexpressing at least one valine biosynthetic enzyme in a microorganism which has at least one pantothenate biosynthetic enzyme deleted.

The present invention is also based at least in part, on the identification and characterization of a previously unidentified microbial pantothenate kinase gene,

10

15

20

25

30

35

coaX. CoaX was first identified in Bacillus subtilis and corresponds to an open reading frame in a portion of the chromosomal DNA that includes the 5' end of the ftsH gene, and all of the yacB, yacC, yacD, cysK and pabB genes. The present inventors have demonstrated that the yacB open reading frame encodes a novel pantothenate kinase activity, the gene being unrelated by homology to any previously known pantothenate kinase gene. The gene has been renamed coaX, as it encodes the enzyme which catalyzes the first step in the pathway from pantothenate to CoaA.

Accordingly, the present invention features new and improved methods of producing pantothenate and other key compounds of the pantothenate biosynthetic pathway (e.g., panto-compounds) utilizing microorganisms having modified pantothenate kinase activity. In particular, the present invention features recombinant microorganisms that contain the coaX gene or that contain a mutant coaX gene, having reduced pantothenate kinase activity. In one embodiment, the invention features such recombinant microorganisms further having a deregulated pantothenate biosynthetic pathway. In another embodiment, the invention features such recombinant microorganisms further having a deregulated isoleucine-valine (ilv) pathway. In a preferred embodiment, the microorganisms belong to the genus Bacillus (e.g., B. subtilis).

The present invention also features recombinant microorganisms (e.g., microorganisms belonging to the genus Bacillus, for example, B. subtilis) that contain the coaA gene or that contain a mutant coaA gene, optionally including a coaX gene or mutant thereof, having reduced pantothenate kinase activity. In one embodiment, the invention features such recombinant microorganisms further having a deregulated pantothenate biosynthetic pathway or having a deregulated isoleucine-valine (ilv) pathway.

Also featured are vectors that contain isolated coaX or coaA genes as well as mutant coaX and/or coaA genes. Isolated nucleic acid molecules that contain isolated coaX genes or mutant coaX genes are featured in addition to isolated CoaX proteins and mutant CoaX proteins.

The nucleic acids, vectors and recombinant microorganisms described above are particularly useful in the methodologies of the present invention. In particular, the invention features methods of enhancing panto-compound production (e.g., ketopantoate, pantoate and or pantothenate production) that include culturing a recombinant microorganism having a mutant coaX gene under conditions such that panto-compound production is enhanced. In one embodiment, the recombinant microorganism further includes a mutant coaA gene. In another embodiment, the recombinant microorganism further includes a mutant avtA and/or mutant ilvE gene

15

and/or mutant *ansB* gene and/or mutant *alsD* gene. Also featured are methods for identifying pantothenate modulators utilizing the recombinant microorganisms and purified CoaX proteins of the present invention.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

Brief Description of the Drawings

Figure 1 is a schematic representation of the pantothenate biosynthetic pathway.

Figure 2 is a schematic representation of the plasmid pAN240, containing sequences ligated upstream of the $P_{26}panBCD$ cassette, equivalent to the integrated version in strain PA221.

Figure 3A is a schematic representation of the plasmid pAN004, containing the panBCD operon expressed from P_{26} and RBS1.

Figure 3B is a schematic representation of the plasmid pAN006, containing the panBCD operon expressed from P_{26} and RBS2.

Figure 4 is a schematic representation of the plasmid pAN236, containing an integratable and amplifiable P_{26} -RBS2-panE1 expression cassette.

20 Figure 5 is a schematic representation of the construction of plasmid pAN423.

Figure 6 is a schematic representation of the construction of plasmids pAN426 and pAN427.

Figure 7 is a schematic representation of the construction of plasmids pAN428 and pAN429.

Figure 8 is a schematic representation of the construction of plasmid pAN431.

Figure 9 is a schematic representation of the construction of plasmid pAN441.

Figure 10 is a schematic representation of the construction of plasmid pAN440.

Figure 11 is a schematic representation of the plasmid pAN251 designed to integrate a single copy of a P_{26} -panE1 cassette at the panE1 locus by double crossover.

35 Figure 12 is a schematic representation of the plasmid pAN267 designed to integrate a single copy of a P_{26} -ilvBNC cassette at the amyE locus.

10

15

20

25

30

35

Figure 13 is a schematic representation of the plasmid pAN257, a clone of Bacillus subtilis ilvD in a low copy vector.

Figure 14 is a schematic representation of the plasmid pAN263, designed to integrate a single copy of a P_{26} -ilvD cassette at the ilvD locus.

Figure 15 is a schematic representation of the plasmid pAN261, designed to disrupt the Bacillus subtilis ilvD gene with the cat gene.

Figure 16 is a schematic representation of the Coenzyme A biosynthetic pathway in E. coli.

Figure 17 is a schematic representation of the structure of pAN296, a plasmid designed to delete most of the *B. subtilis coaA* gene and substitute a chloramphenicol resistance gene.

Figure 18 is a schematic representation of the structure of the Bacillus subtilis genome in the region of the coaA gene. The scale is in base pairs and the significant open reading frames are shown by open arrows.

Figure 19 is a schematic representation of the plasmid pAN281, a plasmid for expressing Bacillus subtilis coaA after integration at the bpr locus.

Figure 20A-B depicts a multiple sequence alignment (MSA) of the amino acid sequences encoded by six known or predicted microbial coaA genes. SEQ ID NOs:4-6 and 1-3 correspond to the amino acid sequences of Mycobacterium leprae (SwissProtTM Accession No. Q9X795), Mycobacterium tuberculosis (SwissProtTM Accession No. O53440), Streptomyces coelicolor (SwissProtTM Accession No. O86799), Haemophilus influenzae (SwissProtTM Accession No. P44793), Escherichia coli SwissProtTM Accession No. P15044) and Bacillus subtilis (SwissProtTM Accession No. P54556), respectively. The alignment was generated using ClustalW MSA software at the GenomeNet CLUSTALW Server at the Institute for Chemical Research, Kyoto University. The following parameters were used: Pairwise Alignment, K-tuple (word) size = 1, Window size = 5, Gap Penalty = 3, Number of Top Diagonals = 5, Scoring Method = Percent; Multiple Alignment, Gap Open Penalty = 10, Gap Extension Penalty = 0.0, Weight Transition = No, Hydrophilic residues = Gly, Pro, Ser, Asn, Asp, Gln, Glu, Arg and Lys, Hydrophobic Gaps = Yes; and Scoring Matrix = BLOSUM.

Figure 21 is a schematic representation of the structure of the Bacillus subtilis genome in the region of the coaX (yacB) gene. The scale is in base pairs, the significant open reading frames are shown by open arrows and certain predicted restriction fragments are indicated by thick bars.

Figure 22 is a schematic representation of the structure of pAN341 and pAN342, two independent PCR-derived clones of B. subtilis yacB (remaned herein as coaX).

Figure 23A-D depicts a multiple sequence alignment (MSA) of the amino acid sequences encoded by fourteen known or predicted microbial coaX genes. SEQ ID NOs:9, 74, 7-8, 75, 11, 10 and 12-18 correspond to the amino acid sequences of Bacillus subtilis (SwissProtTM Accession No. P37564), Clostridium acetobulyticum (WITTM

Accession No. RCA03301, Argonne National Laboratories), Streptomyces coelicolor (PIRTM Accession No. T36391), Mycobacterium tuberculosis (SwissProtTM Accession No. O06282), Rhodobacter capsulatus (WITTM Accession No. RRC02473), Desulfovibrio vulgaris (DBJTM Accession No. BAA21476.1), Deinococcus radiodurans (SwissProtTM Accession No. Q9RX54), Thermotoga maritima (GenBankTM Accession

No. AAD35964.1), Treponema pallidum (SwissProtTM Accession No. O83446),

Borrelia burgdorferi (SwissProtTM Accession No.O51477), Aquifex aeolicus

(SwissProtTM Accession No. O67753), Synechocystis sp. (SwissProtTM Accession No. P74045), Helicobacter pylori (SwissProtTM Accession No. O25533), and Bordetella pertussis (SwissProtTM Accession No. Q45338), respectively. The alignment was

generated using ClustalW MSA software at the GenomeNet CLUSTALW Server at the Institute for Chemical Research, Kyoto University. The following parameters were used: Pairwise Alignment, K-tuple (word) size = 1, Window size = 5, Gap Penalty = 3, Number of Top Diagonals = 5, Scoring Method = Percent; Multiple Alignment, Gap Open Penalty = 10, Gap Extension Penalty = 0.0, Weight Transition = No, Hydrophilic residues = Gly, Pro, Ser, Asn, Asp, Gln, Glu, Arg and Lys, Hydrophobic Gaps = Yes;

residues = Gly, Pro, Ser, Asn, Asp, Gln, Glu, Arg and Lys, Hydrophobic Gaps = Yes; and Scoring Matrix = BLOSUM.

Figure 24 depicts a multiple sequence alignment of a portion of the protein sequences of the coaA gene products from the following microorganisms: Bacillus subtilis, Escherichia coli, Haemophilus influenzae, Mycobacterium leprae,

25 Mycobacterium tuberculosis, and Streptomyces coelicolor. The residues that are mutated in E. coli coaA15(Ts) and B. subtilis coaA282A are indicated below and above the alignment, respectively. The portions correspond to amino acid residues 168-187 of SEQ ID NO:3, 167-186 of SEQ ID NO:2, 165-184 of SEQ ID NO:1, 169-188 of SEQ ID NO:4, 169-188 of SEQ ID NO:5 and 179-198 of SEQ ID NO:6, respectively.

Figure 25 is a schematic representation of the structure of pAN294, a plasmid for integrating mutagenized B. subtilis coaA at its native locus.

Figure 26 is a schematic representation of the structure of pAN336, a plasmid designed to delete *B. subtilis coaX* from its chromosomal locus and replace it with a kanamycin resistence gene.

30

10

15

20

- 12 -

Detailed Description of the Invention

The present invention features new and improved methods of producing pantothenate and other key compounds of the pantothenate biosynthetic pathway (referred to herein as "panto-compounds", for example, pantothenate, ketopantoate, pantoate and β -alanine) using microorganisms in which the pantothenate biosynthetic pathway has been manipulated such that pantothenate or other desired panto-compounds are produced.

The new and improved methodologies of the present invention include methods of producing panto-compounds (e.g., pantothenate) in microorganisms having at least one enzyme of the pantothenate biosynthetic pathway manipulated such that pantothenate or other desired panto-compounds are produced (e.g., produced at an increased level). For example, the invention features methods of producing pantocompounds (e.g., pantothenate) in microorganisms having at least one of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase or aspartateα-decarboxylase manipulated such that pantothenate or other desired panto-compounds are produced. The methodologies of the present invention also include methods of producing panto-compounds (e.g., pantothenate) in microorganisms having at least one valine-isoleucine biosynthetic enzyme, described herein, manipulated such that pantothenate or other desired panto-compounds are produced. For example, the invention features methods of producing panto-compounds (e.g., pantothenate) in microorganisms having at least one of acetohydroxyacid synthetase, acetohydroxyacid isomeroreductase or dihydroxyacid dehydratase manipulated such that pantothenate or other desired panto-compounds are produced.

The invention also features methods of producing panto-compounds that involve culturing a ketopantoate reductase-overexpressing (KPAR-O) microorganism 25 under conditions such that the panto-compound is produced. The invention also features methods of producing pantothenate in a manner independent of precursor feed that involve culturing an aspartate-α-decarboxylase-overexpressing (AαD-O) microorganism under conditions such that pantothenate is produced. Also featured are β -alanine independent high yield pantothenate production methods as well as methods of 30 producing \(\textit{B-alanine}\). The present invention also features methods for enhancing production of panto-compounds that involve culturing pantothenate kinase mutants. In particular, the present invention features new and improved methods of producing pantothenate and other key compounds of the pantothenate biosynthetic pathway (e.g., panto-compounds) utilizing microorganisms having modified pantothenate kinase 35 activity, for example, microorganisms that include the coaX gene or that include a mutant *coaX* gene, having reduced pantothenate kinase activity.

15

20

35

In order that the present invention may be more readily understood, certain terms are first defined herein.

The term "pantothenate biosynthetic pathway" includes the biosynthetic pathway involving pantothenate biosynthetic enzymes (e.g., polypeptides encoded by biosynthetic enzyme-encoding genes), compounds (e.g., precursors, substrates, intermediates or products), cofactors and the like utilized in the formation or synthesis of pantothenate. The term "pantothenate biosynthetic pathway" includes the biosynthetic pathway leading to the synthesis of pantothenate in a microorganisms (e.g., in vivo) as well as the biosynthetic pathway leading to the synthesis of pantothenate in vitro. Figure 1 includes a schematic representation of the pantothenate biosynthetic pathway. Pantothenate biosynthetic enzymes are depicted in bold and their corresponding genes indicated in italics.

The term "pantothenate biosynthetic enzyme" includes any enzyme utilized in the formation of a compound (e.g., intermediate or product) of the pantothenate biosynthetic pathway. According to Figure 1, synthesis of pantoate from α -ketoisovalerate (α -KIV) proceeds via the intermediate, ketopantoate. Formation of ketopantoate is catalyzed by the pantothenate biosynthetic enzyme ketopantoate hydroxymethyltransferase (the panB gene product). Formation of pantoate is catalyzed by the pantothenate biosynthetic enzyme ketopantoate reductase (the panE gene product). Synthesis of β -alanine from aspartate is catalyzed by the pantothenate biosynthetic enzyme aspartate- α -decarboxylase (the panD gene product). Formation of pantothenate from pantoate and β -alanine (e.g., condensation) is catalyzed by the pantothenate biosynthetic enzyme pantothenate synthetase (the panC gene product).

The term "isoleucine-valine biosynthetic pathway" includes the

25 biosynthetic pathway involving isoleucine-valine biosynthetic enzymes (e.g.,
polypeptides encoded by biosynthetic enzyme-encoding genes), compounds (e.g.,
precursors, substrates, intermediates or products), cofactors and the like utilized in the
formation or synthesis of conversion of pyruvate to valine or isoleucine. The term
"isoleucine-valine biosynthetic pathway" includes the biosynthetic pathway leading to

30 the synthesis of valine or isoleucine in a microorganisms (e.g., in vivo) as well as the
biosynthetic pathway leading to the synthesis of valine or isoleucine in vitro. Figure 1
includes a schematic representation of the isoleucine-valine biosynthetic pathway.
Isoleucine-valine biosynthetic enzymes are depicted in bold italics and their
corresponding genes indicated in italics

The term "isoleucine-valine biosynthetic enzyme" includes any enzyme utilized in the formation of a compound (e.g., intermediate or product) of the isoleucine-valine biosynthetic pathway. According to Figure 1, synthesis of valine from pyruvate

15

20

proceeds via the intermediates, acetolactate, α,β -dihydroxyisovalerate (α,β -DHIV) and α -ketoisovalerate (α -KIV). Formation of acetolactate from pyruvate is catalyzed by the isoleucine-valine biosynthetic enzyme acetohydroxyacid synthetase (the ilvBN gene product, or alternatively, the alsS gene product). Formation of α,β -DHIV from acetolactate is catalyzed by the isoleucine-valine biosynthetic enzyme acetohydroxyacidisomero reductase (the ilvC gene product). Synthesis of α -KIV from α,β -DHIV is catalyzed by the isoleucine-valine biosynthetic enzyme dihydroxyacid dehydratase (the ilvD gene product). Moreover, valine and isoleucine can be interconverted by branched chain amino acid transaminases.

As used herein, each of ketopantoate, pantoate, β -alanine and pantothenate are "panto-compounds". The term "panto-compound" includes a compound (*e.g.*, a substrate, intermediate or product) in the pantothenate biosynthetic pathway which is downstream from a particular pantothenate biosynthetic enzyme. In one example, a panto-compound is downstream of the pantothenate biosynthetic enzyme ketopantoate hydroxymethyltransferase (the *panB* gene product) and can include ketopantoate, pantoate and/or pantothenate. In another example, a panto-compound is downstream of the pantothenate biosynthetic enzyme ketopantoate reductase (the *panE* gene product) and can include pantoate and/or pantothenate. In yet another example, a panto-compound is downstream of the pantothenate biosynthetic enzyme pantothenate synthetase (the *panC* gene product) and can include pantothenate. In yet another example, a panto-compound is downstream of the pantothenate biosynthetic enzyme aspartate- α -decarboxylase (the *panD* gene product) and can include β -alanine and/or pantothenate.

Preferred panto-compounds include pantothenate and pantoate. The term "pantothenate" includes the free acid form of pantothenate, also referred to as 25 "pantothenic acid" as well as any salt thereof (e.g., derived by replacing the acidic hydrogen of pantothenate or pantothenic acid with a cation, for example, calcium, sodium, potassium, ammonium), also referred to as a "pantothenate salt". The term "panto-compound" also includes alcohol derivatives of pantothenate. Preferred pantothenate salts are calcium pantothenate or sodium pantothenate. A preferred 30 alcohol derivative is pantothenol. Pantothenate salts and/or alcohols of the present invention include salts and/or alcohols prepared via conventional methods from the free acids described herein. In another embodiment, calcium pantothenate is synthesized directly by a microorganism of the present invention. A pantothenate salt of the present invention can likewise be converted to a free acid form of pantothenate or pantothenic 35 acid by conventional methodology.

The term "pantoate" includes the free acid form of pantoate, also referred to as "pantoic acid" as well as any salt thereof (e.g., derived by replacing the acidic hydrogen of pantoate or pantoic acid with a cation, for example, calcium, sodium, potassium, ammonium), also referred to as a "pantoate salt". Preferred pantoate salts are calcium pantoate or sodium pantoate. Pantoate salts of the present invention include salts prepared via conventional methods from the free acids described herein. A pantoate salt of the present invention can likewise be converted to a free acid form of pantoate or pantoic acid by conventional methodology. Moreover, a free acid form of pantoate or pantoic acid can be converted to pantolactone by conventional methodology.

The term "CoA biosynthetic pathway" includes the biosynthetic pathway involving CoA biosynthetic enzymes (e.g., polypeptides encoded by biosynthetic enzyme-encoding genes), compounds (e.g., precursors, substrates, intermediates or products), cofactors and the like utilized in the formation or synthesis of CoA from pantothenate. A schematic representation of the CoA biosynthetic pathway in E. coli is set forth as Figure 16. (The pathway depicted is also presumed to be that utilized by other microorganisms.) The term "CoA biosynthetic pathway" includes the biosynthetic pathway leading to the synthesis of CoA in microorganisms (e.g., in vivo) as well as the biosynthetic pathway leading to the synthesis of CoA in vitro. The term "Coenzyme A or CoA biosynthetic enzyme" includes any enzyme utilized in the formation of a compound (e.g., intermediate or product) of the CoA biosynthetic pathway, for example, the coaA, panK or coaX gene product which catalyzes the phosphorylation of pantothenate to form 4'-phosphopantothenate, or the coaD gene product which catalyzes the conversion of 4'-phosphopantetheine to dephosphocoenzyme A.

25

30

35

20

10

15

I. Recombinant Microorganisms and Methods for Culturing Microorganisms Such That Panto-Compounds are Produced

The methodologies of the present invention feature microorganisms, *e.g.*, recombinant microorganisms, preferably including vectors or genes (*e.g.*, wild-type and/or mutated genes) as described herein and/or cultured in a manner which results in the production of a desired product (*e.g.* a panto-compound or panto-compounds). The term "recombinant" microorganism includes a microorganism (*e.g.*, bacteria, yeast cell, fungal cell, etc.) which has been genetically altered, modified or engineered (*e.g.*, genetically engineered) such that it exhibits an altered, modified or different genotype and/or phenotype (*e.g.*, when the genetic modification affects coding nucleic acid sequences of the microorganism) as compared to the naturally-occurring microorganism from which it was derived. Preferably, a "recombinant" microorganism of the present

15

20

25

30

35

invention has been genetically engineered such that it overexpresses at least one bacterial gene or gene product (e.g., a pantothenate or isoleucine-valine biosynthetic enzyme encoding-gene) as described herein, preferably a biosynthetic enzyme encoding-gene included within a recombinant vector as described herein and/or a biosynthetic enzyme expressed from a recombinant vector. The ordinary skilled will appreciate that a microorganism expressing or overexpressing a gene product produces or overproduces the gene product as a result of expression or overexpression of nucleic acid sequences and/or genes encoding the gene product.

The term "manipulated microorganism" includes a microorganism that has been engineered (e.g., genetically engineered) or modified such that the microorganism has at least one enzyme of the pantothenate biosynthetic pathway and/or at least one enzyme of the isoleucine-valine biosynthetic pathway modified such that pantothenate or other desired panto-compounds are produced. Modification or engineering of such microorganisms can be according to any methodology described herein including, but not limited to, deregulation of a biosynthetic pathway and/or overexpression of at least one biosynthetic enzyme. A "manipulated" enzyme (e.g., a "manipulated" biosynthetic enzyme) includes an enzyme, the expression or production of which has been altered or modified such that at least one upstream or downstream precursor, substrate or product of the enzyme is altered or modified, for example, as compared to a corresponding wild-type or naturally occurring enzyme.

The term "overexpressed" or "overexpression" includes expression of a gene product (e.g., a pantothenate biosynthetic enzyme or isoleucine-valine biosynthetic enzyme) at a level greater than that expressed prior to manipulation of the microorganism or in a comparable microorganism which has not been manipulated. In one embodiment, the microorganism can be genetically manipulated (e.g., genetically engineered) to overexpress a level of gene product greater than that expressed prior to manipulation of the microorganism or in a comparable microorganism which has not been manipulated. Genetic manipulation can include, but is not limited to, altering or modifying regulatory sequences or sites associated with expression of a particular gene (e.g., by adding strong promoters, inducible promoters or multiple promoters or by removing regulatory sequences such that expression is constitutive), modifying the chromosomal location of a particular gene, altering nucleic acid sequences adjacent to a particular gene such as a ribosome binding site or transcription terminator, increasing the copy number of a particular gene, modifying proteins (e.g., regulatory proteins, suppressors, enhancers, transcriptional activators and the like) involved in transcription of a particular gene and/or translation of a particular gene product, or any other conventional means of deregulating expression of a particular gene routine in the art

20

25

30

35

(including but not limited to use of antisense nucleic acid molecules, for example, to block expression of repressor proteins).

In another embodiment, the microorganism can be physically or environmentally manipulated to overexpress a level of gene product greater than that expressed prior to manipulation of the microorganism or in a comparable microorganism which has not been manipulated. For example, a microorganism can be treated with or cultured in the presence of an agent known or suspected to increase transcription of a particular gene and/or translation of a particular gene product such that transcription and/or translation are enhanced or increased. Alternatively, a microorganism can be cultured at a temperature selected to increase transcription of a particular gene and/or translation of a particular gene product such that transcription and/or translation are enhanced or increased.

The term "deregulated" or "deregulation" includes the alteration or modification of at least one gene in a microorganism that encodes an enzyme in a biosynthetic pathway, such that the level or activity of the biosynthetic enzyme in the microorganism is altered or modified. Preferably, at least one gene that encodes an enzyme in a biosynthetic pathway is altered or modified such that the gene product is enhanced or increased. The phrase "deregulated pathway" can also include a biosynthetic pathway in which more than one gene that encodes an enzyme in a biosynthetic pathway is altered or modified such that the level or activity of more than one biosynthetic enzyme is altered or modified. The ability to "deregulate" a pathway (e.g., to simultaneously deregulate more than one gene in a given biosynthetic pathway) in a microorganism arises from the particular phenomenon of microorganisms in which more than one enzyme (e.g., two or three biosynthetic enzymes) are encoded by genes occurring adjacent to one another on a contiguous piece of genetic material termed an "operon".

The term "operon" includes a coordinated unit of gene expression that contains a promoter and possibly a regulatory element associated with one or more, preferably at least two, structural genes (e.g., genes encoding enzymes, for example, biosynthetic enzymes). Expression of the structural genes can be coordinately regulated, for example, by regulatory proteins binding to the regulatory element or by antitermination of transcription. The structural genes can be transcribed to give a single mRNA that encodes all of the structural proteins. Due to the coordinated regulation of genes included in an operon, alteration or modification of the single promoter and/or regulatory element can result in alteration or modification of each gene product encoded by the operon. Alteration or modification of the regulatory element can include, but is not limited to removing the endogenous promoter and/or regulatory element(s), adding

15

20

strong promoters, inducible promoters or multiple promoters or removing regulatory sequences such that expression of the gene products is modified, modifying the chromosomal location of the operon, altering nucleic acid sequences adjacent to the operon or within the operon such as a ribosome binding site, increasing the copy number of the operon, modifying proteins (*e.g.*, regulatory proteins, suppressors, enhancers, transcriptional activators and the like) involved in transcription of the operon and/or translation of the gene products of the operon, or any other conventional means of deregulating expression of genes routine in the art (including but not limited to use of antisense nucleic acid molecules, for example, to block expression of repressor proteins). Deregulation can also involve altering the coding region of one or more genes to yield, for example, an enzyme that is feedback resistant or has a higher or lower specific activity.

A particularly preferred "recombinant" microorganism of the present invention has been genetically engineered to overexpress a bacterially-derived gene or gene product. The term "bacterially-derived" or "derived-from", for example bacteria, includes a gene which is naturally found in bacteria or a gene product (*e.g.*, ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase, aspartate-α-decarboxylate, acetohydroxyacid synthetase, acetohydroxyacid isomeroreductase or dihydroxyacid dehydratase) which is encoded by a bacterial gene (*e.g.*, encoded by *panB*, *panC*, *panD*, *ilvB*, *ilvN*, *alsS*, *ilvC*, or *ilvD*).

The methodologies of the present invention feature recombinant microorganisms which overexpress at least one of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase or aspartate- α -decarboxylase. A particularly preferred recombinant microorganism of the present invention has been genetically engineered to overexpress a Bacillus (e.g., Bacillus 25 licheniformis, Bacillus amyloliquefaciens, Bacillus halodurans, Bacillus subtilis, and Bacillus pumilus, etc.) biosynthetic enzyme (e.g., has been engineered to overexpress at least one of B. subtilis ketopantoate reductase (the panE gene product) (e.g., ketopantoate reductase having the amino acid sequence of SEQ ID NO:30 or encoded by the nucleic acid sequence of SEQ ID NO:29), B. subtilis ketopantoate 30 hydroxymethyltransferase (the panB gene product) (e.g., ketopantoate hydroxymethyltransferase having the amino acid sequence of SEQ ID NO:24 or encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:23), B. subtilis pantothenate synthetase (the panC gene product) (e.g., pantothenate synthetase having the amino acid sequence of SEQ ID NO:26 or encoded by a nucleic 35 acid molecule having the nucleotide sequence of SEQ ID NO:25) and/or B. subtilis

aspartate- α -decarboxylase (the panD gene product) (e.g., aspartate- α -decarboxylase

15

20

25

30

35

having the amino acid sequence of SEQ ID NO:28 or encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:27).

In an exemplary embodiment, the invention features a microorganism (e.g., a KPAR-O microorganism) that has been transformed with a vector comprising a panE nucleic acid sequence (e.g., a panE nucleic acid sequence as set forth in SEQ ID NO:29). In another embodiment, the invention features a microorganism that has been transformed with a vector comprising a panB nucleic acid sequence (e.g., a panB nucleic acid sequence as set forth in SEQ ID NO:23), a vector comprising a panC nucleic acid sequence (e.g., a panC nucleic acid sequence as set forth in SEQ ID NO:25) or a vector comprising a panD nucleic acid sequence (e.g., a panD nucleic acid sequence as set forth in SEQ ID NO:27). In yet another embodiment, the invention features a microorganism having a deregulated panBCD operon (e.g., SEQ ID NO:59).

Other preferred "recombinant" microorganisms of the present invention have a deregulated isoleucine-valine (ilv) pathway. The phrase "microorganism having a deregulated isoleucine-valine (ilv) pathway" includes a microorganism having an alteration or modification in at least one gene encoding an enzyme of the isoleucinevaline (ilv) pathway or having an alteration or modification in an operon including more than one gene encoding an enzyme of the isoleucine-valine (ilv) pathway. A preferred "microorganism having a deregulated isoleucine-valine (ilv) pathway" has been genetically engineered to overexpress a Bacillus (e.g., B. subtilis) ilv biosynthetic enzyme (e.g., has been engineered to overexpress at least one of acetohydroxyacid synthetase (the ilvBN gene products or the alsS gene product) (e.g., acetohydroxyacid synthetase having subunits having the amino acid sequences of SEQ ID NO:32 and SEQ ID NO:34 or encoded by nucleic acid molecules having the nucleotide sequence of SEQ ID NO:31 and SEQ ID NO:33 or the nucleotide sequence of SEQ ID NO:58 from nucleotides 1-2246 or acetohydroxyacid synthetase encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:86), acetohydroxyacid isomeroreductase (the ilvC gene product) (e.g., acetohydroxyacid isomeroreductase having the amino acid sequence of SEQ ID NO:36 or encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:35), dihydroxyacid dehydratase (the ilvD gene product) (e.g., dihydroxyacid dehydratase having the amino acid sequence of SEQ ID NO:38 or encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:37), and/or has been transformed with a vector comprising an ilvBNC nucleic acid sequence (SEQ ID NO:58, coding regions from nucleotides 1-1725, 1722-2246 and 2263-3291) and/or an ilvD nucleic acid sequence (SEQ ID NO:37).

In another preferred embodiment, a recombinant microorganism is designed or engineered such that a mutant CoaA and/or CoaX biosynthetic enzyme is

į. **4**

13

expressed and at least one pantothenate biosynthetic enzyme and/or at least one isoleucine-valine biosynthetic enzyme is overexpressed or deregulated.

In another preferred embodiment, a microorganism of the present invention overexpresses or is mutated for a gene or biosynthetic enzyme (*e.g.*, a CoA biosynthetic enzyme, pantothenate biosynthetic enzyme or isoleucine-valine biosynthetic enzyme) which is bacterially-derived. The term "bacterially-derived" or "derived-from", for example bacteria, includes a gene product (*e.g.*, ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase, aspartate-α-decarboxylate, acetohydroxyacid synthetase, acetohydroxyacid isomeroreductase, dihydroxyacid dehydratase or pantothenate kinase) which is encoded by a bacterial gene (*e.g.*, *panB*, *panE*, *panC*, *panD*, *ilvBN* (or *alsS*), *ilvC*, *ilvD*, or encoded by *coaA* or *coaX*).

Still other preferred recombinant microorganisms of the present invention are mutant microorganisms. As used herein, the term "mutant microorganism" includes a recombinant microorganism that has been genetically engineered to express a mutated 15 gene or protein that is normally or naturally expressed by the microorganism. Preferably, a mutant microorganism expresses a mutated gene or protein such that the microorganism exhibits an altered, modified or different phenotype (e.g., has been engineered to express a mutated CoaA biosynthetic enzyme, for example, pantothenate kinase). In one embodiment, a mutant microorganism is designed or engineered such 20 that it includes a mutant coaX gene, as defined herein. In another embodiment, a recombinant microorganism is designed or engineered such that it includes a mutant coaA gene, as defined herein. In another embodiment, a mutant microorganism is designed or engineered such that a coaX gene has been deleted (i.e., the protein encoded by the coaX gene is not produced). In another embodiment, a mutant microorganism is 25 designed or engineered such that a coaA gene has been deleted (i.e., the protein encoded by the coaA gene is not produced). Preferably, a mutant microorganism has a mutant coaX gene or a mutant coaA gene, or has been engineered to have a coaX gene and/or coaA deleted, such that that the mutant microorganism encodes a "reduced pantothenate kinase activity". In the context of a whole microorganism, a "reduced pantothenate 30 kinase activity" can be determined by measuring or assaying for a decrease in an intermediate or product of the CoA biosynthetic pathway, for example, measuring or assaying for 4'-phosphopantothenate, 4'-phosphopantothenylcysteine, 4'phosphopantetheine, dephosphocoenzyme A, Coenzyme A, apo-acyl carrier protein (apo-ACP) or holo-acyl carrier protein (ACP) in the microorganism (e.g., in a lysate 35 isolated or derived from the microorganism) or in the medium in which the microorganism is cultured (see e.g., Figure 16). Alternatively, a "reduced pantothenate

30

35

kinase activity" can be determined by measuring or assaying for decreased growth of the microorganism. Alternatively, a "reduced pantothenate kinase activity" can be determined by measuring or assaying for an increase in a panto-compound (e.g., pantothenate) in the microorganism or surrounding media, as panto-compounds lie upstream of the CoA biosynthetic pathway, the first step of which is catalyzed by pantothenate kinase. The invention also features recombinant microorganisms that, in addition to having reduced pantothenate kinase activity (e.g., expressing mutant coaA and/or mutant coaX genes) have a deregulated pantothenate biosynthesis pathway and/or a deregulated isoleucine-valine (ilv) biosynthetic pathway.

In one embodiment, a recombinant microorganism of the present 10 invention is a Gram positive organism (e.g., a microorganism which retains basic dye, for example, crystal violet, due to the presence of a Gram-positive wall surrounding the microorganism). In a preferred embodiment, the recombinant microorganism is a microorganism belonging to a genus selected from the group consisting of Bacillus, Cornyebacterium, Lactobacillus, Lactococci and Streptomyces. In a more preferred 15 embodiment, the recombinant microorganism is of the genus Bacillus. In another preferred embodiment, the recombinant microorganism is selected from the group consisting of Bacillus subtilis, Bacillus lentimorbus, Bacillus lentus, Bacillus firmus, Bacillus pantothenticus, Bacillus amyloliquefaciens, Bacillus cereus, Bacillus circulans, Bacillus coagulans, Bacillus licheniformis, Bacillus megaterium, Bacillus pumilus, 20 Bacillus thuringiensis, and other Group 1 Bacillus species, for example, as characterized by 16S rRNA type (Priest (1993) in Bacillus subtilis and Other Gram-Positive Bacteria eds. Sonenshein et al., ASM, Washington, D.C., p. 6). In another preferred embodiment, the recombinant microorganism is Bacillus brevis or Bacillus stearothermophilus. In another preferred embodiment, the recombinant microorganism 25 is selected from the group consisting of Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus halodurans, Bacillus subtilis, and Bacillus pumilus.

In another embodiment, the recombinant microorganism is a Gram negative (excludes basic dye) organism. In a preferred embodiment, the recombinant microorganism is a microorganism belonging to a genus selected from the group consisting of *Salmonella*, *Escherichia*, *Klebsiella*, *Serratia*, and *Proteus*. In a more preferred embodiment, the recombinant microorganism is of the genus *Escherichia*. In an even more preferred embodiment, the recombinant microorganism is *Escherichia coli*. In another embodiment, the recombinant microorganism is *Saccharomyces* (e.g., *S. cerevisiae*).

An important aspect of the present invention involves culturing the recombinant microorganisms described herein, such that a desired compound (e.g., a

25

30

35

desired panto-compound) is produced. The term "culturing" includes maintaining and/or growing a living microorganism of the present invention (e.g., maintaining and/or growing a culture or strain). In one embodiment, a microorganism of the invention is cultured in liquid media. In another embodiment, a microorganism of the invention is cultured in solid media or semi-solid media. In a preferred embodiment, a microorganism of the invention is cultured in media (e.g., a sterile, liquid media) comprising nutrients essential or beneficial to the maintenance and/or growth of the microorganism (e.g., carbon sources or carbon substrate, for example complex carbohydrates such as bean or grain meal, starches, sugars, sugar alcohols, hydrocarbons, oils, fats, fatty acids, organic acids and alcohols; nitrogen sources, for 10 example, vegetable proteins, peptones, peptides and amino acids derived from grains, beans and tubers, proteins, peptides and amino acids derived form animal sources such as meat, milk and animal byproducts such as peptones, meat extracts and casein hydrolysates; inorganic nitrogen sources such as urea, ammonium sulfate, ammonium chloride, ammonium nitrate and ammonium phosphate; phosphorus sources, for 15 example, phosphoric acid, sodium and potassium salts thereof; trace elements, for example, magnesium, iron, manganese, calcium, copper, zinc, boron, molybdenum, and/or cobalt salts; as well as growth factors such as amino acids, vitamins, growth promoters and the like).

Preferably, microorganisms of the present invention are cultured under controlled pH. The term "controlled pH" includes any pH which results in production of the desired product (*e.g.*, a panto-compound). In one embodiment, microorganisms are cultured at a pH of about 7. In another embodiment, microorganisms are cultured at a pH of between 6.0 and 8.5. The desired pH may be maintained by any number of methods known to those skilled in the art.

Also preferably, microorganisms of the present invention are cultured under controlled aeration. The term "controlled aeration" includes sufficient aeration (e.g., oxygen) to result in production of the desired product (e.g., panto-compound). In one embodiment, aeration is controlled by regulating oxygen levels in the culture, for example, by regulating the amount of oxygen dissolved in culture media. Preferably, aeration of the culture is controlled by agitating the culture. Agitation may be provided by a propeller or similar mechanical agitation equipment, by revolving or shaking the growth vessel (e.g., fermentor) or by various pumping equipment. Aeration may be further controlled by the passage of sterile air or oxygen through the medium (e.g., through the fermentation mixture). Also preferably, microorganisms of the present invention are cultured without excess foaming (e.g., via addition of antifoaming agents).

15

20

25

Moreover, microorganisms of the present invention can be cultured under controlled temperatures. The term "controlled temperature" includes any temperature which results in production of the desired product (*e.g.*, a panto-compound). In one embodiment, controlled temperatures include temperatures between 15°C and 95°C. In another embodiment, controlled temperatures include temperatures between 15°C and 70°C. Preferred temperatures are between 20°C and 55°C, more preferably between 30°C and 45°C or between 30°C and 50°C.

Microorganisms can be cultured (e.g., maintained and/or grown) in liquid media and preferably are cultured, either continuously or intermittently, by conventional culturing methods such as standing culture, test tube culture, shaking culture (e.g., rotary shaking culture, shake flask culture, etc.), aeration spinner culture, or fermentation. In a preferred embodiment, the microorganisms are cultured in shake flasks. In a more preferred embodiment, the microorganisms are cultured in a fermentor (e.g., a fermentation process). Fermentation processes of the present invention include, but are not limited to, batch, fed-batch and continuous methods of fermentation. The phrase "batch process" or "batch fermentation" refers to a closed system in which the composition of media, nutrients, supplemental additives and the like is set at the beginning of the fermentation and not subject to alteration during the fermentation, however, attempts may be made to control such factors as pH and oxygen concentration to prevent excess media acidification and/or microorganism death. The phrase "fedbatch process" or "fed-batch" fermentation refers to a batch fermentation with the exception that one or more substrates or supplements are added (e.g., added in increments or continuously) as the fermentation progresses. The phrase "continuous process" or "continuous fermentation" refers to a system in which a defined fermentation media is added continuously to a fermentor and an equal amount of used or "conditioned" media is simultaneously removed, preferably for recovery of the desired product (e.g., panto-compound). A variety of such processes have been developed and are well-known in the art.

The phrase "culturing under conditions such that a desired compound (e.g., a panto-compound, for example, pantothenate) is produced" includes maintaining and/or growing microorganisms under conditions (e.g., temperature, pressure, pH, duration, etc.) appropriate or sufficient to obtain production of the desired compound or to obtain desired yields of the particular compound being produced. For example, culturing is continued for a time sufficient to produce the desired amount of a panto-compound (e.g., pantothenate, pantoate or β-alanine). Preferably, culturing is continued for a time sufficient to substantially reach maximal production of the panto-compound. In one embodiment, culturing is continued for about 12 to 24 hours. In another

20

25

30

35

embodiment, culturing is continued for about 24 to 36 hours, 36 to 48 hours, 48 to 72 hours, 72 to 96 hours, 96 to 120 hours, 120 to 144 hours, or greater than 144 hours. In another embodiment, culturing is continued for a time sufficient to reach production yields of panto-compound, for example, cells are cultured such that at least about 15 to 20 g/L of panto-compound are produced, at least about 20 to 25 g/L panto-compound are produced, at least about 30 to 35 g/L panto-compound are produced, at least about 35 to 40 g/L panto-compound are produced (e.g., at least about 37 g/L panto-compound) or at least about 40 to 50 g/L panto compound are produced. In yet another embodiment, microorganisms are cultured under conditions such that a preferred yield of panto-compound, for example, a yield within a range set forth above, is produced in about 24 hours, in about 36 hours, in about 48 hours, in about 72 hours, or in about 96 hours.

The methodology of the present invention can further include a step of recovering a desired compound (e.g., a panto-compound). The term "recovering" a desired compound (e.g., a panto-compound) includes extracting, harvesting, isolating or purifying the compound from culture media. Recovering the compound can be performed according to any conventional isolation or purification methodology known in the art including, but not limited to, treatment with a conventional resin (e.g., anion or cation exchange resin, non-ionic adsorption resin, etc.), treatment with a conventional adsorbent (e.g., activated charcoal, silicic acid, silica gel, cellulose, alumina, etc.), alteration of pH, solvent extraction (e.g., with a conventional solvent such as an alcohol, ethyl acetate, hexane and the like), dialysis, filtration, concentration, crystallization, recrystallization, pH adjustment, lyophilization and the like. For example, a compound (e.g., a panto-compound) can be recovered from culture media by first removing the microorganisms from the culture. Media is then passed through or over a cation exchange resin to remove unwanted cations and then through or over an anion exchange resin to remove unwanted inorganic anions and organic acids having stronger acidities than the panto-compound of interest (e.g., pantothenate). The resulting panto-compound (e.g., pantothenate) can subsequently be converted to a pantothenate salt (e.g., calcium pantothenate) as described herein.

Preferably, a desired compound of the present invention is "extracted", "isolated" or "purified" such that the resulting preparation is substantially free of other components (e.g., free of media components and/or fermentation byproducts). The language "substantially free of other components" includes preparations of desired compound in which the compound is separated (e.g., purified or partially purified) from media components or fermentation byproducts of the culture from which it is produced. In one embodiment, the preparation has greater than about 80% (by dry weight) of the

25

30

35

desired compound (e.g., less than about 20% of other media components or fermentation byproducts), more preferably greater than about 90% of the desired compound (e.g., less than about 10% of other media components or fermentation byproducts), still more preferably greater than about 95% of the desired compound (e.g., less than about 5% of other media components or fermentation byproducts), and most preferably greater than about 98-99% desired compound (e.g., less than about 1-2% other media components or fermentation byproducts). When the desired compound is a panto-compound that has been derivatized to a salt (e.g. a pantothenate salt or pantoate salt), the panto-compound is preferably further free (e.g., substantially free) of chemical contaminants associated with the formation of the salt. When the desired compound is a panto-compound that has been derivatized to an alcohol, the panto-compound is preferably further free (e.g., substantially free) of chemical contaminants associated with the formation of the alcohol.

In an alternative embodiment, the desired panto-compound is not purified from the microorganism, for example, when the microorganism is biologically non-hazardous (e.g., safe). For example, the entire culture (or culture supernatant) can be used as a source of product (e.g., crude product). In one embodiment, the culture (or culture supernatant) supernatant is used without modification. In another embodiment, the culture (or culture supernatant) is concentrated. In yet another embodiment, the culture (or culture supernatant) is dried or lyophilized.

II. Panto-Compound Production Methodologies Featuring Ketopantoate Reductase-Overexpressing Microorganisms

One aspect of the invention features methods of producing a panto-compounds that involve culturing a ketopantoate reductase-overexpressing (KPAR-O) microorganism under conditions such that the panto-compound is produced. The term "ketopantoate reductase-overexpressing (KPAR-O) microorganism" includes a microorganism which has been manipulated such that ketopantoate reductase is overexpressed (e.g., a B. subtilis ketopantoate reductase protein having the amino acid sequence of SEQ ID NO:30) and/or has been transformed with a vector comprising a panE1 nucleic acid sequence (e.g., a B. subtilis panE1 nucleic acid sequence as set forth in SEQ ID NO:29). In one embodiment, the panto-compound is pantothenate. In another embodiment, the panto-compound is pantoate. In another embodiment, the ketopantoate reductase is bacterial-derived. In another embodiemnt, the ketopantoate reductase is derived from Bacillus (e.g., is derived from Bacillus subtilis). In yet another embodiment, the KPAR-O microorganism is Gram positive. In yet another

15

20

25

embodiment, the KPAR-O microorganism is a microorganism belonging to a genus selected from the group consisting of Bacillus, Cornyebacterium, Lactobacillus, Lactococci and Streptomyces. In a preferred embodiemnt, the KPAR-O microorganism is of the genus Bacillus. In a more preferred embodiment, the KPAR-O microorganism is selected from the group consisting of Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus halodurans, Bacillus subtilis and Bacillus pumilus. In a particularly preferred embodiemnt, the KPAR-O microorganism is Bacillus subtilis.

In still other embodiments, the KPAR-O microorganism further overexpresses at least one pantothenate biosynthetic enzyme in addition to ketopantoate reductase. In an exemplary embodiment, the KPAR-O microorganism further overexpresses at least one of ketopantoate hydroxymethyltransferase, pantothenate synthetase and aspartate- α -decarboxylase. Also featured are methods of producing panto-compounds, for example, methods that involve culturing a KPAR-O microorganism, which further include the step of recovering the panto-compound.

Methods of Producing Panto-Compounds Independent of III.Precursor Feed Requirements

Depending on the biosynthetic enzyme or combination of biosynthetic enzymes manipulated, it may be desirable or necessary to provide (e.g., feed) microorganisms of the present invention at least one pantothenate biosynthetic precursor such that pantothenate or other desired panto-compounds are produced. The term "pantothenate biosynthetic precursor" or "precursor" includes an agent or compound which, when provided to, brought into contact with, or included in the culture medium of a microorganism, serves to enhance or increase pantothenate biosynthesis. In one embodiment, the pantothenate biosynthetic precursor or precursor is aspartate. In another embodiment, the pantothenate biosynthetic precursor or precursor is β -alanine. The amount of aspartate or β -alanine added is preferably an amount that results in a concentration in the culture medium sufficient to enhance productivity of the microorganism (e.g., a concentration sufficient to enhance production of a pantocompound, for example, β -alanine, ketopantoate, pantoate or pantothenate). 30 Pantothenate biosynthetic precursors of the present invention can be added in the form of a concentrated solution or suspension (e.g., in a suitable solvent such as water or buffer) or in the form of a solid (e.g., in the form of a powder). Moreover, pantothenate biosynthetic precursors of the present invention can be added as a single aliquot, continuously or intermittently over a given period of time. 35

In yet another embodiment, the pantothenate biosynthetic precursor is valine, see e.g., Example III. In yet another embodiment, the pantothenate biosynthetic

15

20

25

30

35

- 27 -

precursor is α -ketoisovalerate. Preferably, valine or α -ketoisovalerate is added in an amount that results in a concentration in the medium sufficient for production of the desired product (e.g., panto-compound) to occur. Pantothenate biosynthetic precursors are also referred to herein as "supplemental pantothenate biosynthetic substrates".

Providing pantothenate biosynthetic precursors in the pantothenate biosynthetic methodologies of the present invention, can be associated with high costs, for example, when the methodologies are used to produce high yields of pantocompounds. Accordingly, preferred methodologies of the present invention feature microorganisms having at least one biosynthetic enzyme or combination of biosynthetic enzymes (e.g., at least one pantothenate biosynthetic enzyme and/or valine-isoleucine biosynthetic enzyme) manipulated such that pantothenate or other desired pantocompounds are produced in a manner independent of precursor feed. The phrase "a manner independent of precursor feed", for example, when referring to a method for producing a desired compound (e.g., a panto-compound), includes an approach to or a mode of producing the desired compound that does not depend or rely on precursors being provided (e.g., fed) to the microorganism being utilized to produce the desired compound. For example, microorganisms featured in the methodologies of the present invention can be used to produce panto-compounds in a manner requiring no feeding of the precursors aspartate, β -alanine, valine and/or α -KIV.

Alternative preferred methodologies of the present invention feature microorganisms having at least one biosynthetic enzyme or combination of biosynthetic enzymes manipulated such that pantothenate or other desired panto-compounds are produced in a manner substantially independent of precursor feed. The phrase "a manner substantially independent of precursor feed" includes an approach to or a method of producing the desired compound that depends or relies to a lesser extent on precursors being provided (e.g., fed) to the microorganism being utilized. For example, microorganisms featured in the methodologies of the present invention can be used to produce panto-compounds in a manner requiring feeding of substantially reduced amounts of the precursors aspartate, β -alanine, valine and/or α -KIV. In one embodiment, the invention features methods of producing panto-compounds (e.g., pantothenate) in a manner that requires feeding of less than 5%-10% of the amount of precursor required by a control microorganism (e.g., a microorganism that is dependent, for example is wholly dependent, on precursor feed to efficiently produce the desired compound). In another embodiment, the invention features methods of producing panto-compounds in a manner that requires feeding of less than 15-20% of the amount of precursor required by a control microorganism. In another embodiment, the invention features methods of producing panto-compounds in a manner that requires

1.3

13

10

15

20

feeding of less than 25-30%, 35-40%, 45-50% or 55-60% of the amount of precursor required by a control microorganism. As described in Examples I-III herein, particular microorganisms featured in the methodologies of the present invention require, for example, 5 g/L of aspartate, β-alanine, valine or α-KIV (*e.g.*, in test tube or in shake flask cultures). Accordingly, in a preferred embodiment, the present invention features methods of producing panto-compounds (*e.g.*, pantothenate) in a manner requiring feeding of less than 0.25 g/L, 0.5 g/L, 0.75 g/L, 1 g/L, 1.25 g/L, 1.5 g/L, 1.75 g/L, 2 g/L, 2.25 g/L, 2.5 g/L, 2.75 g/L or 3 g/L.

Preferred methods of producing desired compounds (*e.g.*, pantocompounds) in a manner independent of precursor feed or alternatively, in a manner substantially independent of precursor feed, involve culturing microorganisms which have been manipulated (*e.g.*, designed or engineered, for example, genetically engineered) such that expression of at least one pantothenate biosynthetic enzyme, and/or at least one isoleucine-valine biosynthetic enzyme, is modified. For example, in one embodiment, a microorganism is manipulated (*e.g.*, designed or engineered) such that the production of at least one pantothenate biosynthetic enzyme, and/or at least one isoleucine/valine biosynthetic enzyme is deregulated. In a preferred embodiment, a microorganism is manipulated (*e.g.*, designed or engineered) such that it has a deregulated biosynthetic pathway, for example, a deregulated pantothenate biosynthesis pathway and/or a deregulated isoleucine-valine biosynthetic pathway, as defined herein. In another preferred embodiment, a microorganism is manipulated (*e.g.*, designed or engineered) such that at least one pantothenate biosynthetic enzyme, and/or at least one isoleucine-valine biosynthetic enzyme is overexpressed.

Preferred methods of producing desired compounds (e.g., pantocompounds) in a manner independent of precursor feed or alternatively, in a manner 25 substantially independent of precursor feed, are as follows. In one embodiment, the invention features a method of producing pantothenate in a manner independent of precursor feed comprising culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism having a deregulated isoleucine-valine (ilv) pathway under conditions such that pantothenate is produced. In another embodiment, the invention features a 30 method of producing pantothenate in a manner independent of precursor feed comprising culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism having a deregulated pantothenate (pan) pathway and a deregulated isoleucine-valine (ilv) pathway, under conditions such that pantothenate is produced. In another embodiment, the invention features a method of producing pantothenate in a 35 manner independent of aspartate or β -alanine feed comprising culturing an aspartate- α decarboxylase-overexpressing (AaD-O) microorganism under conditions such that

15

20

25

30

35

pantothenate is produced. In yet another embodiment, the invention features a method of producing pantothenate in a manner independent of valine or α -ketoisovalerate feed comprising culturing a microorganism having a deregulated isoleucine-valine (ilv) biosynthetic pathway under conditions such that pantothenate is produced.

The term "aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism" includes a microorganism which has been manipulated such that aspartate- α -decarboxylase is overexpressed. A preferred "aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism" has been transformed with a vector comprising a *B. subtilis panD* nucleic acid sequence (*e.g.*, a *panD* nucleic acid sequence that encodes an aspartate- α -decarboxylase protein having the amino acid sequence of SEQ ID NO:28, for example, a *panD* nucleic acid sequence as set forth in SEQ ID NO:27).

The phrase "microorganism having a deregulated isoleucine-valine (*ilv*) pathway" includes a microorganism having an alteration or modification in at least one gene encoding an enzyme of the isoleucine-valine (*ilv*) pathway or having an alteration or modification in an operon including more than one gene encoding an enzyme of the isoleucine-valine (*ilv*) pathway. A preferred "microorganism having a deregulated isoleucine-valine (*ilv*) pathway" overexpresses acetohydroxyacid synthetase (*e.g.*, acetohydroxyacid synthetase having subunits having the amino acid sequences of SEQ ID NO:34 or acetohydroxyacid synthetase having the amino acid sequence of SEQ ID NO:87), acetohydroxyacid isomeroreductase (having the amino acid sequence of SEQ ID NO:36), or dihydroxyacid dehydratase (having the amino acid sequence of SEQ ID NO:38) and/or has been transformed with a vector comprising *ilvB*, *ilvN*, *ilvC*, *ilvBN*, *ilvBNC* or *alsS* nucleic acid sequences (SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, nucleotides 1-2246 of SEQ ID NO:58, SEQ ID NO:58 having coding regions from nucleotides 1-1725, 1722-2246 and 2263-3291, or SEQ ID NO:86, respectively) and/or an *ilvD* nucleic acid sequence (SEQ ID NO:37).

IV. High Yield Production Methodologies

A particularly preferred embodiment of the present invention is a high yield production method for producing pantothenate comprising culturing a manipulated microorganism under conditions such that pantothenate is produced at a significantly high yield. The phrase "high yield production method", for example, a high yield production method for producing a desired compound (*e.g.*, for producing a pantocompound) includes a method that results in production of the desired compound at a level which is elevated or above what is usual for comparable production methods. Preferably, a high yield production method results in production of the desired

15

20

25

30

35

compound at a significantly high yield. The phrase "significantly high yield" includes a level of production or yield which is sufficiently elevated or above what is usual for comparable production methods, for example, which is elevated to a level sufficient for commercial production of the desired product (e.g., production of the product at a commercially feasible cost). In one embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 2 g/L. In another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 10 g/L. In another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 20 g/L. In yet another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 30 g/L. In yet another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 40 g/L.

The invention further features a high yield production method for producing a desired compound (e.g., for producing a panto-compound) that involves culturing a manipulated microorganism under conditions such that a sufficiently elevated level of compound is produced within a commercially desireable period of time. In an exemplary embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 15-20 g/L in 36 hours. In another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 25-30 g/L in 48 hours. In another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 35-40 g/L in 72 hours, for example, greater that 37 g/L in 72 hours. In another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 30-40 g/L in 60 hours, for example, greater that 30, 35 or 40 g/L in 60 hours. Values and ranges included and/or intermediate within the ranges

set forth herein are also intended to be within the scope of the present invention. For example, pantothenate production at levels of at least 31, 32, 33, 34, 35, 36, 37, 38 and 39 g/L in 60 hours are intended to be included within the range of 30-40 g/L in 60 hours. In another example, ranges of 30-35 g/L or 35-40 g/L are intended to be included within the range of 30-40 g/L in 60 hours. Moreover, the skilled artisan will appreciate that culturing a manipulated microorganism to achieve a production level of, for example, "30-40 g/L in 60 hours" includes culturing the microorganism for additional time periods (*e.g.*, time periods longer than 60 hours), optionally resulting in even higher yields of pantothenate being produced.

10

15

20

25

30

35

V. Panto-Compound Production Methodologies Featuring Pantothenate Kinase Mutant Microorganisms

The present invention relates to methods of producing pantothenate using microorganisms engineered to produce high yields of pantothenate as well as other panto-compounds. Cells overproducing pantothenate result in high intracellular pantothenate levels that could overcome the feedback inhibition of pantothenate kinase by CoA, leading to overproduction of CoA. Besides consuming pantothenate, increased synthesis of CoA may cause increased feedback inhibition of the PanB, PanD, PanE or PanC reaction, thereby limiting pantothenate production. Accordingly, a reduction in pantothenate kinase activity may lead to a decrease in CoA levels with resulting increases in PanB, PanD, PanE or PanC activity and pantothenate production.

Thus, certain methodologies of the present invention are based, at least in part, on the identification and characterization of the *B. subtilis coaA* gene and the demonstration that the gene is neither essential for *B. subtilis* growth (*i.e.*, deletion of the *coaA* gene from the chromosome of *B. subtilis* is not a lethal event) nor for pantothenate kinase activity in *B. subtilis*. A second pantothenate kinase-encoding gene has been identified and characterized in *B. subtilis*, and is termed "*coaX*". This gene complements an *E. coli* mutant that contains a temperature sensitive pantothenate kinase and is not related by homology to any previously known pantothenate kinase gene.

In one aspect, the methodologies of the invention feature recombinant microorganisms that include the *coaX* gene or that include a mutant *coaX* gene, having reduced pantothenate kinase activity. In one embodiment, the methodologies feature such recombinant microorganisms further having a deregulated pantothenate biosynthetic pathway. In another embodiment, the methodologies feature such recombinant microorganisms further having a deregulated isoleucine-valine (*ilv*) pathway. In a preferred embodiment, the microorganisms belong to the genus *Bacillus* (*e.g.*, *B. subtilis*).

10

15

20

25

30

35

The methodologies of the invention also feature recombinant microorganisms (e.g., microorganisms belong to the genus Bacillus, for example, Bacillus) that include the coaA gene or that include a mutant coaA gene, optionally including a coaX gene or mutant thereof, having reduced pantothenate kinase activity.

In one embodiment, the methodologies feature such recombinant microorganisms further having a deregulated pantothenate biosynthetic pathway or having a deregulated isoleucine-valine (*ilv*) pathway. Also featured are vectors that include isolated *coaX* or *coaA* genes as well as mutant *coaX* and/or *coaA* genes. Isolated nucleic acid molecules that include isolated *coaX* genes or mutant *coaX* genes are features in addition to isolated CoaX proteins and mutant CoaX proteins.

The above-described nucleic acid molecules (e.g., genes), proteins, vectors, and recombinant microorganisms (e.g., mutant microorganisms), are particularly suited for use in methods of producing panto-compounds and/or methods of enhancing panto-compound production. In one embodiment, the invention features a method for producing a panto-compound (e.g., pantothenate) that includes culturing a pantothenate kinase mutant (e.g., a recombinant microorganism that misexpresses, e.g., is mutated for, pantothenate kinase, as defined herein) under conditions such that pantocompound is produced. In another embodiment, the invention features a method for enhancing production of a panto-compound (e.g., pantothenate) that includes culturing a pantothenate kinase mutant (e.g., a recombinant microorganism that misexpresses, e.g., is mutated for, pantothenate kinase, as defined herein) under conditions such that production of the panto-compound is produced. As used herein, the term "enhancing" (for example, in the context of the phrase "enhancing production") includes increasing the level or rate of production of panto-compound (e.g., pantothenate) as compared to the level or rate of production in a non-mutant microorganism (e.g., a microorganism having a normal pantothenate kinase gene(s) and/or having normal pantothenate production rates and/or levels.

Preferably, the level of panto-compound produced in methodologies featuring the pantothenate kinase mutants of the present invention is increased by at least 5% as compared to the level produced by a non-mutant (e.g., a recombinant microorganism expressing non-mutated pantothenate kinase). Even more preferably, the level of panto-compound is increased 10% as compared to methodologies featuring non-mutants. Even more preferably, panto-compound levels (e.g., pantothenate levels) are increased 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, are increased 2-fold, 5-fold, 10-fold, 50-fold, 100-fold or more as compared to methodologies featuring non-mutants.

35

- 33 -

VI. Additional Mutations Resulting in Enhanced Panto-Compound

Production

The methodologies of the present invention further can include, for example in addition to overexpressing or deregulating a pantothenate biosynthetic 5 enzyme and/or an isoleucine-valine biosynthetic enzyme, or in addition to mutating a pantothenate-kinase encoding gene, deleting or mutating an enzyme that catalyzes the conversion of key pantothenate biosynthesis substrates or precursors to unwanted or undesirable products. For example, mutating the ilvE gene (Kuramitsu et al. (1985) J. Biochem. 97:993-999) or a homologue thereof (SEQ ID NO:62 or SEQ ID NO:64), 10 thereby limiting the conversion of α -ketoisovalerate to valine, in addition to mutating a pantothenate kinase encoding enzyme, is predicted to result in even further enhanced or increased production of panto-compound. Alternatively, mutating the ansB gene (Sun and Seflow (1991) J. Bacteriol. 173:3831-3845) or a homologue thereof (SEQ ID NO:66), thereby limiting the degradation of aspartate, in addition to mutating a 15 pantothenate kinase encoding enzyme, is predicted to result in even further enhanced or increased production of panto-compound. Alternatively, mutating the alsD gene (Renna et al. (1993) J. Bacteriol. 175:3863-3875) or a homologue thereof (SEQ ID NO:68), thereby limiting the conversion of acetolactate to acetoin, in addition to mutating a pantothenate kinase encoding enzyme, is predicted to result in even further enhanced or 20 increased production of panto-compound. Alternatively, mutating the avtA gene encoding alanine-valine transaminase or a homologue thereof, thereby limiting the conversion of α -ketoisovalerate to valine, in addition to mutating a pantothenate kinase encoding enzyme, is predicted to result in even further enhanced or increased production of panto-compound. Mutating the avtA gene can include mutating, for example, an avtA 25 gene having the nucleotide sequence of SEQ ID NO:70 (e.g., the E. coli avtA gene), or a structural homolog thereof (e.g., a homologue encoding a protein having 30-40%, 40-50%, 50-60%, 60-70%, 70-80%, 80-90%, 90-95% or more identity with the amino acid sequence of SEQ ID NO:71) or a functional homologue (e.g., a gene encoding a structurally unrelated protein having alanine-valine transmainase activity. Such 30 mutations can be accomplished using the methodologies as exemplified in the Examples

Accordingly, in one embodiment, the invention features a method of producing a panto-compound which includes culturing a microorganism having a mutant pantothenate kinase-encoding gene and which further has a deletion or mutation in an *avtA*, *ilvE*, *ansB*, and/or *alsD* gene, or homologue thereof. In another embodiment, the invention features a method of producing a panto-compound which includes

(e.g., Examples XIII, XV, XVI and XVII).

15

20

25

30

35

culturing a microorganism having a mutant pantothenate-kinase encoding gene and a deregulated pantothenate biosynthetic pathway enzyme and which further has a deletion or mutation in an avtA, ilvE, ansB, and/or alsD gene, or homologue thereof. In another embodiment, the invention features a method of producing a panto-compound which includes culturing a microorganism having a mutant pantothenate-kinase encoding gene and a deregulated isoleucine-valine biosynthetic pathway enzyme and which further has a deletion or mutation in an avtA, ilvE, ansB, and/or alsD gene, or homologue thereof.

- 34 -

Mutating the alsD gene can be particularly useful when accomplished in conjunction with overexpression or deregulation of the alsS gene, for example, to prevent carbon (e.g., acetolactate) from being drawn away from the precursor pool utilized for α -KIV production. Accordingly, to maximize the contribution of the als locus to panto-compound production, it is desirable to disrupt the alsD gene in addition to overexpressing the alsS gene. To disrupt the alsD gene, appropriate fragments of the als operon, flanking the alsD gene, are amplified by PCR and cloned to provide homology for creating the disruptions. A drug resistance gene, such as the cat gene, is cloned between the flanking DNA fragments in place of the alsD gene, and the linearized DNA is transformed into a pantothenate production strain such as PA824, selecting for drug-resistance. To overexpress alsS, the alsS coding sequence (e.g., an alsS coding sequence that has been engineered by PCR for expression) is cloned into an expression vector. Vectors which express alsS (or alternatively, vectors which express alsS plus ilvC) are the introduced into panto-compound production strains (e.g., the pantothenate producing strain PA824).

The methodologies of the present invention further can include, for example in addition to overexpressing or deregulating a pantothenate biosynthetic enzyme and/or an isoleucine-valine biosynthetic enzyme, or in addition to mutating a pantothenate-kinase encoding gene, deleting or mutating an enzyme that catalyzes the conversion of desired panto-compounds to unwanted or undesireable downstream products.

Isolated Nucleic Acid Molecules and Genes VII.

Another aspect of the present invention features isolated nucleic acid molecules that encode Bacillus proteins (e.g., B. subtilis proteins), for example, Bacillus pantothenate biosynthetic enzymes (e.g., B. subtilis pantothenate biosynthetic enzymes) or Bacillus valine-isoleucine biosynthetic enzymes (e.g., B. subtilis valine-isoleucine biosynthetic enzymes). Also featured are isolated coaX and/or coaA nucleic acid molecules (e.g., isolated coaX and/or coaA genes) as well as isolated nucleic acid molecules that include such coaX and/or coaA nucleic acid molecules or genes.

i. 4

13

10

15

The term "nucleic acid molecule" includes DNA molecules (*e.g.*, linear, circular, cDNA or chromosomal DNA) and RNA molecules (*e.g.*, tRNA, rRNA, mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA. The term "isolated" nucleic acid molecule includes a nucleic acid molecule which is free of sequences which naturally flank the nucleic acid molecule (*i.e.*, sequences located at the 5' and 3' ends of the nucleic acid molecule) in the chromosomal DNA of the organism from which the nucleic acid is derived. In various embodiments, an isolated nucleic acid molecule can contain less than about 10 kb, 5 kb, 4kb, 3kb, 2kb, 1 kb, 0.5 kb, 0.1 kb, 50 bp, 25 bp or 10 bp of nucleotide sequences which naturally flank the nucleic acid molecule in chromosomal DNA of the microorganism from which the nucleic acid molecule is derived. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular materials when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized.

The term "gene", as used herein, includes a nucleic acid molecule (e.g., a DNA molecule or segment thereof), for example, a protein or RNA-encoding nucleic acid molecule, that in an organism, is separated from another gene or other genes, by intergenic DNA (i.e., intervening or spacer DNA which naturally flanks the gene and/or separates genes in the chromosomal DNA of the organism). A gene may direct 20 synthesis of an enzyme or other protein molecule (e.g., may comprise coding sequences, for example, a contiguous open reading frame (ORF) which encodes a protein) or may itself be functional in the organism. A gene in an organism, may be clustered in an operon, as defined herein, said operon being separated from other genes and/or operons by the intergenic DNA. Individual genes contained within an operon may overlap 25 without intergenic DNA between said individual genes. An "isolated gene", as used herein, includes a gene which is essentially free of sequences which naturally flank the gene in the chromosomal DNA of the organism from which the gene is derived (i.e., is free of adjacent coding sequences which encode a second or distinct protein or RNA molecule, adjacent structural sequences or the like) and optionally includes 5' and 3' 30 regulatory sequences, for example promoter sequences and/or terminator sequences. In one embodiment, an isolated gene includes predominantly coding sequences for a protein (e.g., sequences which encode Bacillus proteins). In another embodiment, an isolated gene includes coding sequences for a protein (e.g., for a Bacillus protein) and adjacent 5' and/or 3' regulatory sequences from the chromosomal DNA of the organism 35 from which the gene is derived (e.g., adjacent 5' and/or 3' Bacillus regulatory sequences). Preferably, an isolated gene contains less than about 10 kb, 5 kb, 2 kb, 1 kb,

35

0.5 kb, 0.2 kb, 0.1 kb, 50 bp, 25 bp or 10 bp of nucleotide sequences which naturally flank the gene in the chromosomal DNA of the organism from which the gene is derived.

In one aspect, the present invention features isolated *panB* nucleic acid sequences or genes, isolated *panC* nucleic acid sequences or genes, isolated *panD* nucleic acid sequences or genes, isolated *panE* nucleic acid sequences or genes, isolated *ilvB*, *ilvN*, *ilvBN* nucleic acid sequences or genes, isolated *alsS* nucleic acid sequences or genes, isolated *ilvC* nucleic acid sequences or genes and/or isolated *ilvD* nucleic acid sequences or genes.

In a preferred embodiment, the nucleic acid or gene is derived from 10 Bacillus (e.g., is Bacillus-derived). The term "derived from Bacillus" or "Bacillusderived" includes a nucleic acid or gene which is naturally found in microorganisms of the genus Bacillus. Preferably, the nucleic acid or gene is derived from a microorganism selected from the group consisting of Bacillus subtilis, Bacillus lentimorbus, Bacillus lentus, Bacillus firmus, Bacillus pantothenticus, Bacillus 15 amyloliquefaciens, Bacillus cereus, Bacillus circulans, Bacillus coagulans, Bacillus licheniformis, Bacillus megaterium, Bacillus pumilus, Bacillus thuringiensis, and other Group 1 Bacillus species, for example, as characterized by 16S rRNA type (Priest, supra). In another preferred embodiment, the nucleic acid or gene is derived from Bacillus brevis or Bacillus stearothermophilus. In another preferred embodiment, the 20 nucleic acid molecules and/or genes of the present invention are derived from a microorganism selected from the group consisting of Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus halodurans, Bacillus subtilis, and Bacillus pumilus. In a particularly preferred embodiment, the nucleic acid or gene is derived from Bacillus subtilis (e.g., is Bacillus subtilis-derived). The term "derived from Bacillus subtilis" or 25 "Bacillus subtilis-derived" includes a nucleic acid or gene which is naturally found in Bacillus subtilis. In yet another preferred embodiment, the nucleic acid or gene is a Bacillus gene homologue (e.g., is derived from a species distinct from Bacillus but having significant homology to a Bacillus gene of the present invention, for example, a Bacillus pan gene or Bacillus ilv gene). 30

Included within the scope of the present invention are bacterial-derived nucleic acid molecules or genes and/or *Bacillus*-derived nucleic acid molecules or genes (e.g., B. subtilis-derived nucleic acid molecules or genes), for example, the genes identified by the present inventors, for example, *Bacillus* or *B. subtilis coaX* genes, coaA genes, pan genes and/or ilv genes. Further included within the scope of the present invention are bacterial-derived nucleic acid molecules or genes and/or *Bacillus*-derived nucleic acid molecules or genes (e.g., B. subtilis-derived nucleic acid molecules or

genes) (e.g., B. subtilis nucleic acid molecules or genes) which differ from naturallyoccurring bacterial and/or Bacillus nucleic acid molecules or genes (e.g., B. subtilis nucleic acid molecules or genes), for example, nucleic acid molecules or genes which have nucleic acids that are substituted, inserted or deleted, but which encode proteins substantially similar to the naturally-occurring gene products of the present invention. In one embodiment, an isolated nucleic acid molecule comprises at least one of the nucleotide sequences set forth as SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:86, SEQ ID NO:35 or SEQ ID NO:37. In another preferred embodiment, an isolated nucleic acid molecule comprises at least two, three or four of the nucleotide sequences set forth as SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35 or SEQ ID NO:37. For example, a preferred isolated nucleic acid molecule of the present invention can include the nucleotide sequences of SEQ ID NO:23, SEQ ID NO:25 and SEQ ID NO:27, preferably linked such that the proteins encoded by the nucleotide sequences of SEQ ID NO:23, SEQ ID NO:25 and SEQ ID 15 NO:27 are each produced when the isolated nucleic acid molecule is expressed in a microorganism (e.g., SEQ ID NO:59). In another example, a preferred isolated nucleic acid molecule of the present invention can include the nucleotide sequences of SEQ ID NO:31 and SEQ ID NO:33, preferably linked such that the proteins encoded by the nucleotide sequences of SEQ ID NO:31 and SEQ ID NO:33 are each produced when the 20 isolated nucleic acid molecule is expressed in a microorganism (e.g., nucleotides 1-2246 of SEQ ID NO:58). In another example, a preferred isolated nucleic acid molecule of the present invention can include the nucleotide sequence of SEQ ID NO:86. In another example, a preferred isolated nucleic acid molecule of the present invention can include the nucleotide sequences of SEQ ID NO:31, SEQ ID NO:33 and SEQ ID NO:35, 25 preferably linked such that the proteins encoded by the nucleotide sequences of SEQ ID NO:31, SEQ ID NO:33 and SEQ ID NO:35 are each produced when the isolated nucleic acid molecule is expressed in a microorganism (e.g., SEQ ID NO:58).

In another embodiment, an isolated nucleic acid molecule of the present invention comprises a nucleotide sequence which is at least about 60-65%, preferably at least about 70-75%, more preferable at least about 80-85%, and even more preferably at least about 90-95% or more identical to a nucleotide sequence set forth as SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35 or SEQ ID NO:37. In another embodiment, an isolated nucleic acid molecule hybridizes under stringent conditions to a nucleic acid molecule having a nucleotide sequence set forth as SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:88, SEQ

ļ. i

15

20

25

30

35

ID NO:35 or SEQ ID NO:37. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. A preferred, non-limiting example of stringent (*e.g.* high stringency) hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35 or SEQ ID NO:37 corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature.

A nucleic acid molecule of the present invention (e.g., a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35 or SEQ ID NO:37 can be isolated using standard molecular biology techniques and the sequence information provided herein. For example, nucleic acid molecules can be isolated using standard hybridization and cloning techniques (e.g., as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) or can be isolated by the polymerase chain reaction using synthetic oligonucleotide primers designed based upon the sequence of SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35 or SEQ ID NO:37. A nucleic acid of the invention can be amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule which is a complement of the nucleotide sequence shown in SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:33, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35.

Additional panC nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:25, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:26 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:26 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:25 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the

15

20

25

30

35

amino acid sequence of SEQ ID NO:26, or are complementary to a *panC* nucleotide sequence as set forth herein.

Aditional *panD* nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:27, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:28 (*e.g.*, encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:28 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:27 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:28, or are complementary to a *panD* nucleotide sequence as set forth herein.

Additional *panE* nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:29, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:30 (*e.g.*, encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:30 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:29 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:30, or are complementary to a *panE* nucleotide sequence as set forth herein.

Additional *ilvB* nucleic acid sequences are those that comprise the nucleotide sequence of SEQ ID NO:31, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:32 (*e.g.*, encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:32 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:31 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:32, or are complementary to an *ilvB* nucleotide sequence as set forth herein.

Additional *ilvN* nucleic acid sequences are those that comprise the nucleotide sequence of SEQ ID NO:33, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:34 (*e.g.*, encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:34 and a substantially

20

25

30

35

identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:33 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:34, or are complementary to an *ilvN* nucleotide sequence as set forth herein.

Additional *ilvC* nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:35, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:36 (*e.g.*, encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:36 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:35 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:36, or are complementary to an *ilvC* nucleotide sequence as set forth herein.

Additional *ilvD* nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:37, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:38 (*e.g.*, encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:38 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:37 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:38, or are complementary to an *ilvD* nucleotide sequence as set forth herein.

Additional *alsS* nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:86, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:87 (*e.g.*, encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:87 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:86 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:87, or are complementary to an *alsS* nucleotide sequence as set forth herein.

In another embodiment, an isolated nucleic acid molecule is or includes a coaX gene, or portion or fragment thereof. In one embodiment, an isolated coaX nucleic

acid molecule or gene comprises the nucleotide sequence as set forth in SEQ ID NO:19 (e.g., comprises the B. subtilis coaX nucleotide sequence). In another embodiment, an isolated coaX nucleic acid molecule or gene comprises a nucleotide sequence that encodes the amino acid sequence as set forth in SEQ ID NO:9 (e.g., encodes the B. subtilis CoaX amino acid sequence). In yet another embodiment, an isolated coaX nucleic acid molecule or gene encodes a homologue of the CoaX protein having the amino acid sequence of SEQ ID NO:9. As used herein, the term "homologue" includes a protein or polypeptide sharing at least about 30-35%, preferably at least about 35-40%, more preferably at least about 40-50%, and even more preferably at least about 60%, 70%, 80%, 90% or more identity with the amino acid sequence of a wild-type protein or polypeptide described herein and having a substantially equivalent functional or biological activity as said wild-type protein or polypeptide. For example, a CoaX homologue shares at least about 30-35%, preferably at least about 35-40%, more preferably at least about 40-50%, and even more preferably at least about 60%, 70%, 80%, 90% or more identity with the protein having the amino acid sequence set forth as 15 SEQ ID NO:9 and has a substantially equivalent functional or biological activity (i.e., is a functional equivalent) of the protein having the amino acid sequence set forth as SEQ ID NO:9 (e.g., has a substantially equivalent pantothenate kinase activity). In a preferred embodiment, an isolated coaX nucleic acid molecule or gene comprises a nucleotide sequence that encodes a polypeptide as set forth in any one of SEQ ID NO:7, 20 SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:74 or SEQ ID NO:75. In another embodiment, an isolated coaX nucleic acid molecule hybridizes to all or a portion of a nucleic acid molecule having the nucleotide sequence set forth in SEQ ID NO:19 or hybridizes to all or a portion of a nucleic acid 25 molecule having a nucleotide sequence that encodes a polypeptide having the amino acid sequence of any of SEQ ID NOs:7-18, 74 or 75. Such hybridization conditions are known to those skilled in the art and can be found in Current Protocols in Molecular Biology, Ausubel et al., eds., John Wiley & Sons, Inc. (1995), sections 2, 4 and 6. Additional stringent conditions can be found in Molecular Cloning: A Laboratory 30 Manual, Sambrook et al., Cold Spring Harbor Press, Cold Spring Harbor, NY (1989), chapters 7, 9 and 11. A preferred, non-limiting example of stringent hybridization conditions includes hybridization in 4X sodium chloride/sodium citrate (SSC), at about 65-70°C (or hybridization in 4X SSC plus 50% formamide at about 42-50°C) followed by one or more washes in 1X SSC, at about 65-70°C. A preferred, non-limiting 35 example of highly stringent hybridization conditions includes hybridization in 1X SSC, at about 65-70°C (or hybridization in 1X SSC plus 50% formamide at about 42-50°C)

35

followed by one or more washes in 0.3X SSC, at about 65-70°C. A preferred, nonlimiting example of reduced stringency hybridization conditions includes hybridization in 4X SSC, at about 50-60°C (or alternatively hybridization in 6X SSC plus 50% formamide at about 40-45°C) followed by one or more washes in 2X SSC, at about 50-60°C. Ranges intermediate to the above-recited values, e.g., at 65-70°C or at 42-50°C are also intended to be encompassed by the present invention. SSPE (1X SSPE is 0.15 M NaCl, 10mM NaH₂PO₄, and 1.25 mM EDTA, pH 7.4) can be substituted for SSC (1X SSC is 0.15 M NaCl and 15 mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes each after hybridization is complete. The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length 10 should be 5-10°C less than the melting temperature (T_m) of the hybrid, where T_m is determined according to the following equations. For hybrids less than 18 base pairs in length, $T_m(^{\circ}C) = 2(\# \text{ of } A + T \text{ bases}) + 4(\# \text{ of } G + C \text{ bases})$. For hybrids between 18 and 49 base pairs in length, $T_m(^{\circ}C) = 81.5 + 16.6(log_{10}[Na^+]) + 0.41(\%G+C) - (600/N)$, where N is the number of bases in the hybrid, and [Na+] is the concentration of sodium 15 ions in the hybridization buffer ($[Na^+]$ for 1X SSC = 0.165 M). It will also be recognized by the skilled practitioner that additional reagents may be added to hybridization and/or wash buffers to decrease non-specific hybridization of nucleic acid molecules to membranes, for example, nitrocellulose or nylon membranes, including but not limited to blocking agents (e.g., BSA or salmon or herring sperm carrier DNA), 20 detergents (e.g., SDS), chelating agents (e.g., EDTA), Ficoll, PVP and the like. When using nylon membranes, in particular, an additional preferred, non-limiting example of stringent hybridization conditions is hybridization in 0.25-0.5M NaH₂PO₄, 7% SDS at about 65°C, followed by one or more washes at 0.02M NaH₂PO₄, 1% SDS at 65°C, see e.g., Church and Gilbert (1984) Proc. Natl. Acad. Sci. USA 81:1991-1995, (or, 25 alternatively, 0.2X SSC, 1% SDS). In another preferred embodiment, an isolated nucleic acid molecule comprises a nucleotide sequence that is complementary to a coaX nucleotide sequence as set forth herein (e.g., is the full complement of the nucleotide sequence set forth as SEQ ID NO:19).

In another preferred embodiment, an isolated nucleic acid molecule is or includes a *coaA* gene, for example, a *Bacillus* (*e.g.*, *B. subtilis*) *coaA* gene, or portion or fragment thereof. Exemplary isolated *coaA* nucleic acid molecules and/or genes include (1) an isolated *coaA* nucleic acid molecule or gene comprising the nucleotide sequence as set forth in any one of SEQ ID NOs:20-22; (2) an isolated *coaA* nucleic acid molecule or gene comprising a nucleotide sequence that encodes the amino acid sequence as set forth in SEQ ID NO:3; (3) an isolated *coaA* nucleic acid molecule or gene comprising a nucleotide sequence which encodes a CoaA homologue (*e.g.*, a polypeptide having an

15

20

25

30

35

amino acid sequence at least about 30-35%, preferably at least about 35-40%, more preferably at least about 40-50%, and even more preferably at least about 60%, 70%, 80%, 90% or more identical to the amino acid sequence set forth as SEQ ID NO:3 and having a substantially equivalent enzymatic activity; (4) an isolated *coaA* nucleic acid molecule or gene comprising a nucleotide sequence that encodes a polypeptide as set forth in any one of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6; (5) an isolated nucleic acid molecule that hybridizes under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence set forth in SEQ ID NO:20, SEQ ID NO:21 or SEQ ID NO:22 or hybridizes to all or a portion of a nucleic acid molecule having a nucleotide sequence that encodes a polypeptide having the amino acid sequence of SEQ ID NO:3; and (6) an isolated nucleic acid molecule comprising a nucleotide sequence that is complementary to a *coaA* nucleotide sequence as set forth herein (*e.g.*, is the full complement of the nucleotide sequence set forth in SEQ ID NO:20, SEQ ID NO:21 or SEQ ID NO:22).

- 43 -

A nucleic acid molecule of the present invention (*e.g.*, a *coaX* nucleic acid molecule or gene or a *coaA* nucleic acid molecule or gene), can be isolated using standard molecular biology techniques and the sequence information provided herein. For example, nucleic acid molecules can be isolated using standard hybridization and cloning techniques (*e.g.*, as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) or can be isolated by the polymerase chain reaction using synthetic oligonucleotide primers designed based upon the *coaX* or *coaA* nucleotide sequences set forth herein, or flanking sequences thereof. A nucleic acid of the invention (*e.g.*, a *coaX* nucleic acid molecule or gene or a *coaA* nucleic acid molecule or gene), can be amplified using cDNA, mRNA or alternatively, chromosomal DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques.

Yet another embodiment of the present invention features mutant coaX and coaA nucleic acid molecules or genes. The phrase "mutant nucleic acid molecule" or "mutant gene" as used herein, includes a nucleic acid molecule or gene having a nucleotide sequence which includes at least one alteration (e.g., substitution, insertion, deletion) such that the polypeptide or protein that may be encoded by said mutant exhibits an activity that differs from the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene. Preferably, a mutant nucleic acid molecule or mutant gene (e.g., a mutant coaA or coaX gene) encodes a polypeptide or protein having a reduced activity (e.g., having a reduced pantothenate kinase activity) as compared to the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene, for

15

20

25

30

35

example, when assayed under similar conditions (e.g., assayed in microorganisms cultured at the same temperature). A mutant gene also can encode no polypeptide or have a reduced level of production of the wild-type polypeptide.

As used herein, a "reduced activity" or "reduced enzymatic activity" is one that is at least 5% less than that of the polypeptide or protein encoded by the wildtype nucleic acid molecule or gene, preferably at least 5-10% less, more preferably at least 10-25% less and even more preferably at least 25-50%, 50-75% or 75-100% less than that of the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene. Ranges intermediate to the above-recited values, e.g., 75-85%, 85-90%, 90-95%, are also intended to be encompassed by the present invention. As used herein, a "reduced activity" or "reduced enzymatic activity" also includes an activity that has been deleted or "knocked out" (e.g., approximately 100% less activity than that of the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene). Activity can be determined according to any well accepted assay for measuring activity of a particular protein of interest. Activity can be measured or assayed directly, for example, measuring an activity of a protein isolated or purified from a cell. Alternatively, an activity can be measured or assayed within a cell or in an extracellular medium. For example, assaying for a mutant coaA gene or a mutant coaX gene (i.e., said mutant encoding a reduced pantothenate kinase activity) can be accomplished by expressing the mutated gene in a microorganism, for example, a mutant microorganism which expresses pantothenate kinase in a temperature-sensitive manner, assaying the mutant gene for the ability to complement a temperature sensitive (Ts) mutant for pantothenate kinase activity. A coaX mutant gene or coaA mutant gene that encodes a "reduced pantothenate kinase activity" is one that complements the Ts mutant less effectively than, for example, a corresponding wild-type coaX gene or coaA gene.

It will be appreciated by the skilled artisan that even a single substitution in a nucleic acid or gene sequence (e.g., a base substitution that encodes an amino acid change in the corresponding amino acid sequence) can dramatically affect the activity of an encoded polypeptide or protein as compared to the corresponding wild-type polypeptide or protein. A mutant nucleic acid or mutant gene (e.g., encoding a mutant polypeptide or protein), as defined herein, is readily distinguishable from a nucleic acid or gene encoding a protein homologue, as described above, in that a mutant nucleic acid or mutant gene encodes a protein or polypeptide having an altered activity, optionally observable as a different or distinct phenotype in a microorganism expressing said mutant gene or nucleic acid or producing said mutant protein or polypeptide (i.e., a mutant microorganism) as compared to a corresponding microorganism expressing the wild-type gene or nucleic acid or producing said mutant protein or polypeptide. By

15

20

25

30

35

contrast, a protein homologue has an identical or substantially similar activity, optionally phenotypically indiscernable when produced in a microorganism, as compared to a corresponding microorganism expressing the wild-type gene or nucleic acid. Accordingly it is not, for example, the degree of sequence identity between nucleic acid molecules, genes, protein or polypeptides that serves to distinguish between homologues and mutants, rather it is the activity of the encoded protein or polypeptide that distinguishes between homologues and mutants: homologues having, for example, low (e.g., 30-50% sequence identity) sequence identity yet having substantially equivalent functional activities, and mutants, for example sharing 99% sequence identity yet having dramatically different or altered functional activities. Exemplary homologues are set forth in Figure 20 (i.e., CoaA homologues) and in Figure 23 (i.e., CoaX homologues). Exemplary mutants are described in Examples XV and XVIII herein.

VIII. Recombinant Nucleic Acid Molecules and Vectors

The present invention further features recombinant nucleic acid molecules (e.g., recombinant DNA molecules) that include nucleic acid molecules and/or genes described herein (e.g., isolated nucleic acid molecules and/or genes), preferably Bacillus genes, more preferably Bacillus subtilis genes, even more preferably Bacillus subtilis pantothenate kinase genes (e.g., coaX genes or coaA genes), pantothenate biosynthetic genes (e.g., genes encoding pantothenate biosynthetic enzymes, for example, panB genes encoding ketopantoate hydroxymethyltransferase, panE genes encoding ketopantoate reductase, panC genes encoding pantothenate synthetase, and/or panD genes encoding aspartate-α-decarboxylase) and/or isoleucine-valine (ilv) biosynthetic genes (e.g., ilvBN or alsS genes encoding acetohydroxyacid synthetase, ilvC genes encoding acetohydroxyacid isomeroreductase and/or ilvD genes encoding dihydroxyacid dehydratase).

The present invention further features vectors (e.g., recombinant vectors) that include nucleic acid molecules (e.g., isolated or recombinant nucleic acid molecules and/or genes) described herein. In particular, recombinant vectors are featured that include nucleic acid sequences that encode bacterial gene products as described herein, preferably Bacillus gene products, more preferably Bacillus subtilis gene products, even more preferably Bacillus subtilis pantothenate biosynthetic gene products (e.g., pantothenate biosynthetic enzymes, for example, ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase, and/or

aspartate- α -decarboxylase) and/or isoleucine-valine biosynthetic gene products (e.g.,

15

20

25

30

35

acetohydroxyacid synthetase, acetohydroxyacid isomeroreductase and/or dihydroxyacid dehydratase).

- 46 -

The term "recombinant nucleic acid molecule" includes a nucleic acid molecule (e.g., a DNA molecule) that has been altered, modified or engineered such that it differs in nucleotide sequence from the native or natural nucleic acid molecule from which the recombinant nucleic acid molecule was derived (e.g., by addition, deletion or substitution of one or more nucleotides). Preferably, a recombinant nucleic acid molecule (e.g., a recombinant DNA molecule) includes an isolated nucleic acid molecule or gene of the present invention (e.g., an isolated coaX, coaA, pan or ilv gene) operably linked to regulatory sequences.

The term "recombinant vector" includes a vector (e.g., plasmid, phage, phasmid, virus, cosmid or other purified nucleic acid vector) that has been altered, modified or engineered such that it contains greater, fewer or different nucleic acid sequences than those included in the native or natural nucleic acid molecule from which the recombinant vector was derived. Preferably, the recombinant vector includes a coaX, coaA, pan or ilv gene or recombinant nucleic acid molecule including such coaX, coaA, pan or ilv gene, operably linked to regulatory sequences, for example, promoter sequences, terminator sequences and/or artificial ribosome binding sites (RBSs), as defined herein.

The phrase "operably linked to regulatory sequence(s)" means that the nucleotide sequence of the nucleic acid molecule or gene of interest is linked to the regulatory sequence(s) in a manner which allows for expression (e.g., enhanced, increased, constitutive, basal, attenuated, decreased or repressed expression) of the nucleotide sequence, preferably expression of a gene product encoded by the nucleotide sequence (e.g., when the recombinant nucleic acid molecule is included in a recombinant vector, as defined herein, and is introduced into a microorganism).

The term "regulatory sequence" includes nucleic acid sequences which affect (e.g., modulate or regulate) expression of other nucleic acid sequences. In one embodiment, a regulatory sequence is included in a recombinant nucleic acid molecule or recombinant vector in a similar or identical position and/or orientation relative to a particular gene of interest as is observed for the regulatory sequence and gene of interest as it appears in nature, e.g., in a native position and/or orientation. For example, a gene of interest can be included in a recombinant nucleic acid molecule or recombinant vector operably linked to a regulatory sequence which accompanies or is adjacent to the gene of interest in the natural organism (e.g., operably linked to "native" regulatory sequences, for example, to the "native" promoter). Alternatively, a gene of interest can be included in a recombinant nucleic acid molecule or recombinant vector operably

15

20

25

linked to a regulatory sequence which accompanies or is adjacent to another (e.g., a different) gene in the natural organism. Alternatively, a gene of interest can be included in a recombinant nucleic acid molecule or recombinant vector operably linked to a regulatory sequence from another organism. For example, regulatory sequences from other microbes (e.g., other bacterial regulatory sequences, bacteriophage regulatory sequences and the like) can be operably linked to a particular gene of interest.

In one embodiment, a regulatory sequence is a non-native or nonnaturally-occurring sequence (e.g., a sequence which has been modified, mutated, substituted, derivatized, deleted including sequences which are chemically synthesized). Preferred regulatory sequences include promoters, enhancers, termination signals, antitermination signals and other expression control elements (e.g., sequences to which repressors or inducers bind and/or binding sites for transcriptional and/or translational regulatory proteins, for example, in the transcribed mRNA). Such regulatory sequences are described, for example, in Sambrook, J., Fritsh, E. F., and Maniatis, T. Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989. Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in a microorganism (e.g., constitutive promoters and strong constitutive promoters), those which direct inducible expression of a nucleotide sequence in a microorganism (e.g., inducible promoters, for example, xylose inducible promoters) and those which attenuate or repress expression of a nucleotide sequence in a microorganism (e.g., attenuation signals or repressor sequences). It is also within the scope of the present invention to regulate expression of a gene of interest by removing or deleting regulatory sequences. For example, sequences involved in the negative regulation of transcription can be removed such that expression of a gene of interest is enhanced.

In one embodiment, a recombinant nucleic acid molecule or recombinant vector of the present invention includes a nucleic acid sequence or gene that encodes at least one bacterial gene product (e.g., a pantothenate biosynthetic enzyme, an isoleucine-valine biosynthetic enzyme, or a CoaA biosynthetic enzyme, for example CoaA or CoaX) operably linked to a promoter or promoter sequence. Preferred promoters of the present invention include *Bacillus* promoters and/or bacteriophage promoters (e.g., bacteriophage which infect *Bacillus*). In one embodiment, a promoter is a *Bacillus* promoter, preferably a strong *Bacillus* promoter (e.g., a promoter associated with a biochemical housekeeping gene in *Bacillus* or a promoter associated with a glycolytic pathway gene in *Bacillus*). In another embodiment, a promoter is a bacteriophage promoter. In a preferred embodiment, the promoter is from the bacteriophage SP01. In a particularly preferred embodiment, a promoter is selected from the group consisting of

15

20

25

30

 P_{15} , P_{26} or P_{veg} , for example, the promoters set forth in SEQ ID NO:39, SEQ ID NO:40 or SEQ ID NO:41. Additional preferred promoters include tef (the translational elongation factor (TEF) promoter) and pyc (the pyruvate carboxylase (PYC) promoter), which promote high level expression in Bacillus (e.g., Bacillus subtilis). Additional preferred promoters, for example, for use in Gram positive microorganisms include, but are not limited to, the amyE promoter or phage SP02 promoters. Additional preferred promoters, for example, for use in Gram negative microorganisms include, but are not limited to tac, trp, tet, trp-tet, lpp, lac, lpp-lac, lacIq, T7, T5, T3, gal, trc, ara, SP6, λ-P_R or λ - P_L .

In another embodiment, a recombinant nucleic acid molecule or recombinant vector of the present invention includes a terminator sequence or terminator sequences (e.g., transcription terminator sequences). The term "terminator sequences" includes regulatory sequences which serve to terminate transcription of a gene. Terminator sequences (or tandem transcription terminators) can further serve to stabilize mRNA (e.g., by adding structure to mRNA), for example, against nucleases.

In yet another embodiment, a recombinant nucleic acid molecule or recombinant vector of the present invention includes sequences which allow for detection of the vector containing said sequences (i.e., detectable and/or selectable markers), for example, sequences that overcome auxotrophic mutations, for example, ura3 or ilvE, fluorescent markers, and/or colorimetric markers (e.g., lacZ/βgalactosidase), and/or antibiotic resistance genes (e.g., amp or tet).

In yet another embodiment, a recombinant nucleic acid molecule or recombinant vector of the present invention includes an artificial ribosome binding site (RBS). The term "artificial ribosome binding site (RBS)" includes a site within an mRNA molecule (e.g., coded within DNA) to which a ribosome binds (e.g., to initiate translation) which differs from a native RBS (e.g., a RBS found in a naturally-occurring gene) by at least one nucleotide. Preferred artificial RBSs include about 5-6, 7-8, 9-10, 11-12, 13-14, 15-16, 17-18, 19-20, 21-22, 23-24, 25-26, 27-28, 29-30 or more nucleotides of which about 1-2, 3-4, 5-6, 7-8, 9-10, 11-12, 13-15 or more differ from the native RBS (e.g., the native RBS of a gene of interest). Preferably, nucleotides which differ are substituted such that they are identical to one or more nucleotides of an ideal RBS (e.g., SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47 or SEQ ID NO:48), when optimally aligned for comparisons. Artificial RBSs can be used to replace the naturally-occurring or native RBS associated with a particular gene. Artificial RBSs preferably increase translation of a particular gene. Preferred artificial

35 RBSs (e.g., RBSs for increasing the translation of panB, for example, of B. subtilis panB) are depicted in Table IA (e.g., SEQ ID NO:49 and SEQ ID NO:50).

Table 1A: Preferred panB Ribosome Binding Sites

10 20

----AGAAAGGAGTGA ideal RBS (SEQ ID NO:44)

CCCTCT-AG-AAGGAGGAGAAAACATG RBS1 (SEQ ID NO:49)

CCCTCT-AG--AGGAGGAGAAAACATG RBS2 (SEQ ID NO:50)

TAAACAT-G--AGGAGGAGAAAACATG panB native RBS (SEQ ID NO:42)

Additional preferred artificial RBSs (e.g., RBSs for increasing the translation of panD, for example, of B. subtilis panD) are depicted in Table 1B (e.g., SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53 and SEQ ID NO:54).

15 Table 1B: Preferred panD Ribosome Binding Sites

10 20 CTAGAAAAGGAGGAATTTAAATG (SEQ ID NO:88) pAN423 RBS 20 TTAAGAAAGGAGGTGANNNNATG ideal RBS (SEQ ID NO:45) new design A (SEQ ID NO:51) TTAGAAAGGAGGATTTAAATATG ${\tt TTAGAAAGGAGGT} {\bm TT} {\tt AATTAATG}$ new design B (SEQ ID NO:52) TTAGAAAGGAGGTGATTTAAATG new design C1 (SEQ ID NO:53) 25 TTAGAAAGGAGGTGTTTAAAATG new design C2 (SEQ ID NO:54) TTAGAAAGGAGGTGANNNNATG (SEQ ID NO:46) ideal RBS

Additional preferred artificial RBSs (e.g., RBSs for increasing the translation of panD, for example, of B. subtilis panD) are depicted in Table 1C (e.g., SEQ ID NO:55, SEQ ID NO:56 and SEQ ID NO:57). The predicted amino acid sequence at the C-terminus of the PanC protein is shown. The start codon for PanD translation is underlined.

Table 1C: Additional Preferred panD Ribosome Binding Sites

35 10 20 --- --A GAA AGG AGG TGA NNN NNN N ATG ideal RBS (SEQ ID NO:47) ATT CGA GAA ATG GAG AGA ATA TAA T ATG native panD RBS (SEQ ID NO:43) 40 Ile Arg Glu Met Glu Arg Ile SEQ ID NO:89 Met --- --A GAA AGG AGG TGA NNN NNN N ATG ideal RBS (SEQ ID NO:47) ATT CGA GAA AGG AGG TGA ATA TAA T ATG NDI (SEQ ID NO:55) 45 Ile Arg Glu Arg Arg SEQ ID NO:90 Met

25

30

35

```
- 50 -
```

```
ATT CGA GAA AGG AGG TGA ATA ATA - ATG MDII (SEQ ID NO:56)

Ile Arg Glu Arg Arg * Met SEQ ID NO:90

ATT CGT AGA AAG GAG GTG AAT TAA T ATG NDIII (SEQ ID NO:57)

5 Ile Arg Arg Lys Glu Val Asn * Met SEQ ID NO:91

--- --- AGA AAG GAG GTG ANN NNN N ATG ideal RBS (SEQ ID NO:48)
```

Accordingly, in one embodiment, a vector of the present invention includes an artificial RBS as set forth in SEQ ID NO:49 or SEQ ID NO:50. In another embodiment, a vector of the present invention includes an artificial RBS as set forth in SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53 or SEQ ID NO:54. In yet another embodiment, a vector of the present invention includes an artificial RBS as set forth in SEQ ID NO:55, SEQ ID NO:56 or SEQ ID NO:57.

In another embodiment, a recombinant vector of the present invention includes sequences that enhance replication in bacteria (*e.g.*, replication-enhancing sequences). In one embodiment, replication-enhancing sequences are derived from *E. coli*. In another embodiment, replication-enhancing sequences are derived from pBR322 (*e.g.*, sequences included within the pBR322 derived portion of any of the pAN vectors as set forth in the Figures , *i.e.*, the Not I-Not I sequences from about 5.0 kB to 9.0 kB of the vector depicted in Figure 3A).

In yet another embodiment, a recombinant vector of the present invention includes antibiotic resistance genes. The term "antibiotic resistance genes" includes sequences which promote or confer resistance to antibiotics on the host organism (e.g., Bacillus). In one embodiment, the antibiotic resistance genes are selected from the group consisting of cat (chloramphenicol resistance) genes, tet (tetracycline resistance) genes, erm (erythromycin resistance) genes, neo (neomycin resistance) genes and spec (spectinomycin resistance) genes. Recombinant vectors of the present invention can further include homologous recombination sequences (e.g., sequences designed to allow recombination of the gene of interest into the chromosome of the host organism). For example, amyE sequences can be used as homology targets for recombination into the host chromosome.

Preferred vectors of the present invention include, but are not limited to, vectors set forth in Figures 2-15, 17, 19, 22, 25 and 26. It will further be appreciated by one of skill in the art that the design of a vector can be tailored depending on such factors as the choice of microorganism to be genetically engineered, the level of expression of gene product desired and the like.

20

IX. Isolated Proteins

Another aspect of the present invention features isolated proteins (*e.g.*, isolated pantothenate biosynthetic enzymes and/or valine-isoleucine biosynthetic enzymes and/or isolated CoaA or CoaX). In one embodiment, proteins (*e.g.*, isolated pantothenate biosynthetic enzymes and/or valine-isoleucine biosynthetic enzymes and/or isolated CoaA biosynthetic enzymes, for example isolated CoaA or CoaX) are produced by recombinant DNA techniques and can be isolated from microorganisms of the present invention by an appropriate purification scheme using standard protein purification techniques. In another embodiment, proteins are synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" protein (e.g., an isolated or purified biosynthetic enzyme) is substantially free of cellular material or other contaminating proteins from the microorganism from which the protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. In one embodiment, an isolated or purified protein has less than about 30% (by dry weight) of contaminating protein or chemicals, more preferably less than about 20% of contaminating protein or chemicals, still more preferably less than about 10% of contaminating protein or chemicals, and most preferably less than about 5% contaminating protein or chemicals.

In a preferred embodiment, the protein or gene product is derived from Bacillus (e.g., is Bacillus-derived). The term "derived from Bacillus" or "Bacillusderived" includes a protein or gene product which is encoded by a *Bacillus* gene. 25 Preferably, the gene product is derived from a microorganism selected from the group consisting of Bacillus subtilis, Bacillus lentimorbus, Bacillus lentus, Bacillus firmus, Bacillus pantothenticus, Bacillus amyloliquefaciens, Bacillus cereus, Bacillus circulans, Bacillus coagulans, Bacillus licheniformis, Bacillus megaterium, Bacillus pumilus, Bacillus thuringiensis, and other Group 1 Bacillus species, for example, as characterized by 16S rRNA type (Priest, supra). In another preferred embodiment, the protein or gene 30 product is derived from Bacillus brevis or Bacillus stearothermophilus. In another preferred embodiment, the protein or gene product is derived from a microorganism selected from the group consisting of Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus halodurans, Bacillus subtilis, and Bacillus pumilus. In a particularly preferred embodiment, the protein or gene product is derived from Bacillus subtilis (e.g., is 35 Bacillus subtilis-derived). The term "derived from Bacillus subtilis" or "Bacillus subtilis-derived" includes a protein or gene product which is encoded by a Bacillus

15

20

25

subtilis gene. In yet another preferred embodiment, the protein or gene product is encoded by a *Bacillus* gene homologue (e.g., a gene derived from a species distinct from *Bacillus* but having significant homology to a *Bacillus* gene of the present invention, for example, a *Bacillus pan* gene or *Bacillus ilv* gene).

Included within the scope of the present invention are bacterial-derived proteins or gene products and/or Bacillus-derived proteins or gene products (e.g., B. subtilis-derived gene products) that are encoded by naturally-occurring bacterial and/or Bacillus genes (e.g., B. subtilis genes), for example, the genes identified by the present inventors, for example, Bacillus or B. subtilis coaX genes, coaA genes, pan genes and/or ilv genes. Further included within the scope of the present invention are bacterialderived proteins or gene products and/or Bacillus-derived proteins or gene products (e.g., B. subtilis-derived gene products) that are encoded bacterial and/or Bacillus genes (e.g., B. subtilis genes) which differ from naturally-occurring bacterial and/or Bacillus genes (e.g., B. subtilis genes), for example, genes which have nucleic acids that are mutated, inserted or deleted, but which encode proteins substantially similar to the naturally-occurring gene products of the present invention. For example, it is well understood that one of skill in the art can mutate (e.g., substitute) nucleic acids which, due to the degeneracy of the genetic code, encode for an identical amino acid as that encoded by the naturally-occurring gene. Moreover, it is well understood that one of skill in the art can mutate (e.g., substitute) nucleic acids which encode for conservative amino acid substitutions. It is further well understood that one of skill in the art can substitute, add or delete amino acids to a certain degree without substantially affecting the function of a gene product as compared with a naturally-occurring gene product, each instance of which is intended to be included within the scope of the present invention.

In a preferred embodiment, an isolated protein of the present invention (e.g., an isolated pantothenate biosynthetic enzyme and/or an isolated isoleucine-valine biosynthetic enzyme and/or an isolated CoaA biosynthetic enzymes, for example isolated CoaA or CoaX) has an amino acid sequence shown in SEQ ID NO:3, SEQ ID NO:9, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38 or SEQ ID NO:87. In other embodiments, an isolated protein of the present invention is a homologue of the at least one of the proteins set forth as SEQ ID NO:3, SEQ ID NO:9, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38 or SEQ ID NO:87 (e.g., comprises an amino acid sequence at least about 30-40% identical, preferably about 40-50% identical, more preferably about 50-60% identical, and even more preferably about 60-70%, 70-80%, 80-90%, 90-95% or

:3

10

15

more identical to the amino acid sequence of SEQ ID NO:3, SEQ ID NO:9, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38 or SEQ ID NO:87, and has an activity that is substantially similar to that of the protein encoded by the amino acid sequence of SEQ ID NO:3, SEQ ID NO:9, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38 or SEQ ID NO:87, respectively.

To determine the percent homology of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity = # of identical positions/total # of positions x 100), preferably taking into account the number of gaps and size of said gaps necessary to produce an optimal alignment.

The comparison of sequences and determination of percent homology between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison 20 of sequences is the algorithm of Karlin and Altschul (1990) Proc. Natl. Acad. Sci. USA 87:2264-68, modified as in Karlin and Altschul (1993) Proc. Natl. Acad. Sci. USA 90:5873-77. Such an algorithm is incorporated into the NBLAST and XBLAST programs (version 2.0) of Altschul et al. (1990) J. Mol. Biol. 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, 25 wordlength = 12 to obtain nucleotide sequences homologous to nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (1997) Nucleic 30 Acids Research 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See http://www.ncbi.nlm.nih.gov. Another preferred, nonlimiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller (1988) Comput Appl Biosci. 4:11-17. Such an 35 algorithm is incorporated into the ALIGN program available, for example, at the GENESTREAM network server, IGH Montpellier, FRANCE (http://vega.igh.cnrs.fr) or

13

5

10

15

20

25

30

35

at the ISREC server (http://www.ch.embnet.org). When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

In another preferred embodiment, the percent homology between two amino acid sequences can be determined using the GAP program in the GCG software package (available at http://www.gcg.com), using either a Blossom 62 matrix or a PAM250 matrix, and a gap weight of 12, 10, 8, 6, or 4 and a length weight of 2, 3, or 4. In yet another preferred embodiment, the percent homology between two nucleic acid sequences can be accomplished using the GAP program in the GCG software package (available at http://www.gcg.com), using a gap weight of 50 and a length weight of 3.

X. Biotransformations and Bioconversions

Another aspect of the present invention includes biotransformation processes which feature recombinant microorganisms (*e.g.*, mutant microorganisms) and/or isolated CoA, pantothenate or isoleucine-valine biosynthetic enzymes described herein. The term "biotransformation process", also referred to herein as "bioconversion processes", includes biological processes which result in the production (*e.g.*, transformation or conversion) of any compound (*e.g.*, intermediate or product) which is upstream of a CoA, pantothenate or isoleucine-valine biosynthetic enzyme to a compound (*e.g.*, substrate, intermediate or product) which is downstream of said CoA, pantothenate or isoleucine-valine biosynthetic enzyme.

In one embodiment, the invention features a biotransformation process for the production of a panto-compound comprising contacting a microorganism which overexpresses at least one pantothenate biosynthetic enzyme with at least one appropriate substrate or precursor under conditions such that said panto-compound is produced and recovering said panto-compound. In a preferred embodiment, the invention features a biotransformation process for the production of pantoate comprising contacting a microorganism which overexpresses ketopantoate reductase (the panE gene product) with an appropriate substrate (e.g., ketopantoate) under conditions such that pantoate is produced and recovering said pantoate. In another preferred embodiment, the invention features a biotransformation process for the production of pantothenate comprising contacting a microorganism which overexpresses ketopantoate reductase and pantothenate synthetase with appropriate substrates (e.g., ketopantoate and β -alanine) under conditions such that pantothenate is produced and recovering said pantothenate. In yet another preferred embodiment, the invention features a biotransformation process for the production of pantothenate comprising contacting a microorganism which overexpresses ketopantoate hydroxymethyltransferase, ketopantoate reductase and

15

20

25

pantothenate synthetase with appropriate substrates (e.g., α -ketoisovalerate and β -alanine) under conditions such that pantothenate is produced and recovering said pantothenate. Preferred recombinant microorganisms for carrying out the above-described biotransformations include pantothenate kinase mutants. Conditions under which pantoate or pantothenate are produced can include any conditions which result in the desired production of pantoate or pantothenate, respectively.

In yet another embodiment, the present invention includes a method of producing β -alanine that includes culturing a microorganism which overexpresses aspartate- α -decarboxylase under conditions such that β -alanine is produced. Preferably, the aspartate- α -decarboxylase-overexpressing microorganism has a mutation in a nucleic acid sequence encoding a pantothenate biosynthetic enzyme selected from the group consisting of ketopantoate hydroxymethyltransferase, ketopantoate reductase and pantothenate synthetase.

The invention further features a method of producing β -alanine that includes contacting a composition comprising aspartate with an isolated *Bacillus* aspartate- α -decarboxylase enzyme under conditions such that β -alanine is produced (e.g., an *in vitro* synthesis method).

The microorganism(s) and/or enzymes used in the biotransformation reactions are in a form allowing them to perform their intended function (e.g., producing a desired compound). The microorganisms can be whole cells, or can be only those portions of the cells necessary to obtain the desired end result. The microorganisms can be suspended (e.g., in an appropriate solution such as buffered solutions or media), rinsed (e.g., rinsed free of media from culturing the microorganism), acetone-dried, immobilized (e.g., with polyacrylamide gel or k-carrageenan or on synthetic supports, for example, beads, matrices and the like), fixed, cross-linked or permeablized (e.g., have permeablized membranes and/or walls such that compounds, for example, substrates, intermediates or products can more easily pass through said membrane or wall).

Purified or unpurified CoA biosynthetic enzyme(s) (e.g., CoaA and/or CoaX), pantothenate biosynthetic enzyme(s) and/or valine-isoleucine biosynthetic enzyme(s) can also be used in biotransformation reactions. The enzyme can be in a form that allows it to perform its intended function (e.g., obtaining the desired compound). For example, the enzyme can be in free form or immobilized. Purified or unpurified CoA biosynthetic enzyme(s), pantothenate biosynthetic enzyme(s) and/or valine-isoleucine biosynthetic enzyme(s) can be contacted in one or more in vitro reactions with appropriate substrate(s) such that the desired product is produced.

20

25

30

35

With respect to at least the above-described methodologies (e.g., the production methodologies of the present invention), at least one aspect of the invention features the folowing: embodiments is which the methods do not use microorganisms of the genus Corynebacterium and/or microorganisms of the genus Escherichia; embodiments in which the methods do not use microorganisms selected from the group consiting of Escherichia coli and Corynebacterium glutamicum; embodiments in which the methods do not use gram negative microorganisms; embodiments in which the microorganisms utilized do not include, express or produce nucleic acid molecules, genes or proteins (e.g., biosynthetic emzymes) derived from microorganisms of the genus Corynebacterium and/or microorganisms of the genus Escherichia; embodiments in which the microorganisms to not include, express or produce nucleic acid molecules, genes or proteins (e.g., biosynthetic emzymes) derived from microorganisms selected from the group consisting of Escherichia coli and Corynebacterium glutamicum.

15 XI. Screening Assays

Because CoA is an essential factor in bacteria, proteins (e.g., enzymes) involved in the biosynthesis of CoA provide valuable tools in the search for novel antibiotics. In particular, the CoaX protein is a valuable target for identifying bacteriocidal compounds because it bears no resemblance in primary sequence to mammalian pantothenate kinase enzymes. Accordingly, the present invention also provides a method (also referred to herein as a "screening assay") for identifying modulators, i.e., candidate or test compounds or agents (e.g., peptides, peptidomimetics, small molecules or other drugs) which bind to CoaX, or have a stimulatory or inhibitory effect on, for example, coaX expression or CoaX activity.

In one embodiment, the invention provides assays for screening candidate or test compounds which are capable of binding to CoaX proteins or a biologically active portion thereof. In another embodiment, the invention provides assays for screening candidate or test compounds which modulate the activity of CoaX proteins or biologically active portions thereof. The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-compound' library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam, K.S. (1997) *Anticancer Drug Des.* 12:145).

20

25

30

35

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt et al. (1993) Proc. Natl. Acad. Sci. U.S.A. 90:6909; Erb et al. (1994) Proc. Natl. Acad. Sci. USA 91:11422; Zuckermann et al. (1994). J. Med. Chem. 37:2678; Cho et al. (1993) Science 261:1303; Carrell et al. (1994) Angew. Chem. Int. Ed. Engl. 33:2059; Carell et al. (1994) Angew. Chem. Int. Ed. Engl. 33:2061; and in Gallop et al. (1994) J. Med. Chem. 37:1233. Libraries of compounds may be presented in solution (e.g., Houghten (1992) Biotechniques 13:412-421), or on beads (Lam (1991) Nature 354:82-84), chips (Fodor (1993) Nature 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull et al. (1992) Proc Natl Acad Sci USA 89:1865-1869) or on phage (Scott and Smith (1990) Science 249:386-390); (Devlin (1990) Science 249:404-406); (Cwirla et al. (1990) Proc. Natl. Acad. Sci. 87:6378-6382); (Felici (1991) J. Mol. Biol. 222:301-310); (Ladner supra.).

In one embodiment, an assay is a microorganism-based assay in which a recombinant microorganism which expresses a CoaX protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to modulate CoaX activity is determined. Determining the ability of the test compound to modulate CoaX activity can be accomplished by monitoring, for example, intracellular phosphopanthoate or CoA concentrations or secreted pantothenate concentrations (as compounds that inhibit CoaX will result in a buildup of pantothenate in the test microorganism). CoaX substrate can be labeled with a radioisotope or enzymatic label such that modulation of CoaX activity can be determined by detecting a conversion of labeled substrate to intermediate or product. For example, CoaX substrates can be labeled with ³²P, ¹⁴C, or ³H, either directly or indirectly, and the radioisotope detected by direct counting of radioemmission or by scintillation counting. Determining the ability of a compound to modulate CoaX activity can alternatively be determined by detecting the induction of a reporter gene (comprising a CoA-responsive regulatory element operatively linked to a nucleic acid encoding a detectable marker, e.g., luciferase), or detecting a CoA-regulated cellular response.

In yet another embodiment, a screening assay of the present invention is a cell-free assay in which a CoaX protein or biologically active portion thereof is contacted with a test compound *in vitro* and the ability of the test compound to bind to or modulate the activity of the CoaX protein or biologically active portion thereof is determined. In a preferred embodiment, the assay includes contacting the CoaX protein or biologically active portion thereof with known substrates to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to modulate enzymatic activity of the CoaX on its substrates.

Screening assays can be accomplished in any vessel suitable for containing the microorganisms, proteins, and/or reactants. Examples of such vessels include microtiter plates, test tubes, and micro-centrifuge tubes. In more than one embodiment of the above assay methods of the present invention, it may be desirable to immobilize either CoaX protein or a recombinant microorganism expressing CoaX protein to facilitate separation of products and/or substrates, as well as to accommodate automation of the assay. For example, glutathione-S-transferase/CoaX fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtiter plates. Other techniques for immobilizing proteins on matrices (e.g., biotin-conjugation and streptavidin immobilization or antibody conjugation) can also be used in the screening assays of the invention.

In another embodiment, modulators of CoaX expression are identified in a method wherein a cell is contacted with a candidate compound and the expression of coaX mRNA or CoaX polypeptide in the cell is determined. The level of expression in the presence of the candidate compound is compared to the level of expression in the absence of the candidate compound (or to a suitable control, for example, an appropriate buffer control or standard). The candidate compound can then be identified as a modulator of coaX mRNA or CoaX polypeptide expression based on this comparison.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an CoaX modulating agent identified as described herein (e.g., an anti-bactericidal compound) can be used in an infectious animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent.

25

30

5

10

15

20

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are incorporated herein by reference.

- 59 -

EXAMPLES

General Methodology:

Strains. Bacillus subtilis strains of the present invention are generally derived from either of two strains. The first is variously named "168", "1A1", or "RL-1". The genotype is *trpC2*. This strain was derived from the wild type "Marburg" strain by mutagenesis and has been the basis of much of the molecular biology work done on B. subtilis. The second strain is PY79, a prototrophic derivative of 168 that was made Trp⁺ by transduction from the wild type strain W23.

Media. Standard minimal medium for B. subtilis is comprised of 1 x
Spizizen salts and 0.5% glucose. Standard solid "rich medium" is Tryptone Blood Agar Broth (Difco), and standard liquid "rich medium" is VY, a mixture of veal infusion broth and yeast extract. For testing production of pantothenate in liquid test tube cultures, an enriched form of VY, called "Special VY" or "SVY" is used. For batch fermentations, SVYG and PFMG are used. The compositions of these media are given below.

VY, a rich liquid medium: 25 g Difco <u>V</u>eal Infusion Broth, 5 g Difco Yeast Extract, 1L water (autoclave).

TBAB, a rich solid medium: 33 g Difco <u>Tryptone Blood Agar Broth</u>, 1L water (autoclave).

20 MIN, a minimal medium: 100 ml 10 x Spizizen salts; 10 ml 50% glucose; 2 ml 10% arginine HCl*; 10 ml 0.8% tryptophan**; water to 1 liter. (*In some cases, arginine is omitted or replace by sodium glutamate at 0.04% final concentration. In general, B. subtilis grows faster in minimal medium when certain amino acids, such as arginine, glutamine, glutamate, or proline, are added as an auxiliary nitrogen source.)

25 (**For strains that are tryptophan auxotrophs, tryptophan is routinely added to most minimal media.)

10 x Spizizen Salts: 174 g K₂HPO₄·3H₂O; 20 g (NH₄)₂SO₄; 60 g KH₂PO₄; 10 g Na₃Citrate·2H₂O; 2 g MgSO₄·7H₂O; water to 993 mls; then add 3.5 ml FeCl₃ solution and 3.5 ml Trace Elements solution.

30 FeCl₃Solution: 4 g FeCl₃·6H₂O; 197 g Na₅Citrate·2H₂O; water to 1 liter (filter sterilize)

Trace Elements Solution: 0.15 g Na₂MoO₄·2H₂O; 2.5 g H₃BO₃; 0.7 g CoCl₂·6H₂O; 0.25 g CuSO₄·5H₂O; 1.6 g MnCl₂·4H₂O; 0.3 g ZnSO₄·7H₂O; water to 1 liter (filter sterilize).

10

20

35

SVY, Special VY, a supplemented* rich medium for testing pantothenate production in test tube cultures: 25 g Difco Veal Infusion Broth; 5 g Difco yeast extract; 5 g sodium glutamate; 2.7 g ammonium sulfate; 740 ml water (autoclave); add 200 ml 1 M potassium phosphate, pH 7.0; 60 ml 50% glucose. (*For testing pantothenate production in liquid SVY test tube cultures, Na α-ketoisovalerate and/or β-alanine can be added to a concentration of 5 g/L from filter-sterilized stocks.)

PFMG, a yeast extract based medium used in fermentors: 20 g Amberex 1003[™] yeast extract; 5 g sodium glutamate, 2 g ammonium sulfate; 5 g tryptophan; 10 g KH₂PO4; 20 g K₂HPO₄·3H₂O; 1 g MgCl₂·6H₂O; 0.1 g CaCl₂·2H₂O; 1 g sodium citrate; 0.01 g FeSO₄·7H₂O; 1 ml trace elements solution; 20 g glucose; add water to 1 L. Glucose or other sugars are fed as needed. Feed solutions can contain minerals, defined or food grade nutrients.

PF, a chemically defined pantothenate free medium for testing pantothenate auxotrophy: 100 ml 10 x Spizizen Salts; 100 ml 1 x Difco Pantothenate

15 Assay Medium: 10 ml 50% glucose; water to 1 liter.

For pantothenate auxotrophs, 1 mM Na pantothenate is added to both minimal and rich media, since there is generally not enough pantothenate in rich media to support *B. subtilis pan* mutants. Amino acids are at 100 mg per liter, when used. Selection for antibiotic resistance is done with 5 mg/L chloramphenicol, 100 mg/L spectinomycin HCl, 15 mg/L tetracycline HCl, or 1 mg/L erythromycin plus 25 mg/L lincomycin.

Pantothenate Assays: Biological assay. The indicator organism, Lactobacillus plantarum, requires pantothenate for growth, and responds to low concentrations (μg/L).
Thus, using serial dilutions, a wide range of concentrations can be assayed.
Commercially available medium (e.g., Pantothenate Assay Medium (PAM), Difco), can be used. However, Difco PAM supplemented with pantothenate does not support growth to the same level as obtainable using a fresh-mixed version of Pantothenate Assay Medium (FM-PAM), made up of the individual components as specified by
Difco, which is accordingly, routinely used instead of the commercial product.

Before assaying *B. subtilis* culture supernatants, the *B. subtilis* cells must be either removed or killed. *B. subtilis* culture supernatants give approximately the same pantothenate titer when the supernatants are autoclaved as when they are sterile filtered. Accordingly, routine procedures involve autoclaving samples for 5 minutes prior to the biological assay.

15

20

25

30

Pantothenate Assays: HPLC assay. Pantothenic acid production is measured by HPLC with a detector wavelength of 197 nm and a reference at 450 nm. The procedure is a modification of one recommended by Hewlett-Packard for water soluble vitamins. Samples of culture broth are diluted into an equal volume of 60% acetronitrile (ACN), centrifuged and filtered. Typically a further 10-fold dilution before analysis brings the final dilution to 20-fold. Higher concentrations of product are diluted further. Compounds are separated on a C18 Phenomenex 5μ Aqua 250 x 4.6 mm column with 5% acetronitrile (ACN) in 50 mM Na phosphate buffer at pH 2.5. An ACN gradient from 5% to 95% washes the column between every sample. The area of the pantothenate peak is proportional to the concentration between 5 to 1000 mg/L. Other panto-compounds are also separated and quantitated by this method.

Amino Acid Analysis: HPLC assay. Amino acids present in the fermentation medium and throughout the fermentation are measured by HPLC with a detector wavelength of 338 nm and a reference at 390 nm. The procedure is a modification of one recommended by Hewlett-Packard for amino acid analysis. Samples of culture broth are prepared identically as for the panto-compound analysis. Compounds are separated on a C18 Hypersil 5μ ODS 200 x 2.1 mm column. Solvent A is 20 mM Na acetate buffer at pH 7.2. Solvent B contains 40% ACN and 40% methanol. A gradient from 100% Solvent A to 100% Solvent B separates amino acids and washes the column between every sample.

Batch Fermentations. Pantothenate producing strains are grown in stirred tank fermentors, for example, in CF3000 Chemap 14 liter vessels with 10 liter working volumes. Computer control and data collection is by commercial software, for example, B. Braun Biotech MFCS software. Fermentations can be batch processes but are preferably sugar-limited, fed batch processes. Some media components (e.g. of SVYG and PMFG) are added to the fermentor and sterilized in place. Portions of the media are sterilized separately and added to the fermentors aseptically. This procedure is well known to those familiar with the art. Additional nitrogen sources in feeds are sterilized separately and added to the carbon source after cooling.

The initial sugar in the medium is consumed in approximately 6 hours. Afterwards, glucose or other sugars are fed with the possible addition of minerals, and defined or food grade nutrients. Alternatively, feeds are scheduled based on a consensus profile of nutritional requirements from samples taken from earlier fermentations.

- 62 -

After inoculation, agitation is set at a relatively low speed, *e.g.* 200 rpm. When the dissolved oxygen (pO2) falls to 30%, computer control automatically adjusts the agitation to maintain a dissolved oxygen concentration between 25 and 30% pO2.

5

10

15

20

25

30

35

EXAMPLE I: Enhanced Production of a Panto-Compound Using Bacteria Overexpressing *panBCD* Gene Products.

This Example describes the cloning of the *B. subtilis panBCD* operon and the generation of microorganisms overexpressing the *panBCD* gene products.

To clone the *B. subtilis panBCD* operon, a plasmid library of *B. subtilis* GP275 (a derivative of 168) genomic DNA was transformed in *E. coli* BM4062 (*birA*^{ts}), and temperature resistant clones were selected at 42°C. By comparison of restriction maps to the genome sequence, one particular clone was deduced to contain the *B. subtilis birA* gene and the adjacent *panBCD* genes. This plasmid was named pAN201.

To overexpress the panBCD operon and produce pantothenate, the native promoter of the panBCD operon was replaced by either of two strong, constitutive promoters derived from the B. subtilis bacteriophage SP01. These two promoters are named P_{26} and P_{15} . In addition, either of two artificial ribosome binding sites (RBSs) were used to replace the native panB RBS. These two artificial RBSs (set forth as SEQ ID NO:49 and SEQ ID NO:50) were predicted to increase translation of panBCD; their sequences are shown in Table 1A. Three such engineered panBCD expression cassettes were built into circular plasmids capable of replicating in E. coli. Other features of the plasmids include a strong rho-independent transcription terminator from the E. coli ribosomal RNA transcription unit, called T₁T₂, a Gram-positive chloramphenicol resistance gene (cat), derived from pC194, and a pair of NotI restriction sites at the junctions between the E. coli replicon and the segment intended for integration into B. subtilis. Three plasmids of this series, pAN004, pAN005, and pAN006 were constructed. pAN004 contains the P_{26} promoter, RBS1, and a low copy E. coli replicon. pAN005 contains the P_{15} promoter, which in our experience is not as strong as P_{26} , RBS1, and the low copy replicon. pAN006 contains the P_{26} promoter, RBS2, and a medium copy replicon.

The three *panBCD* expression cassettes contained in the abovementioned three plasmids were all ligated to a DNA fragment consisting of sequences that naturally occur immediately upstream from the native *panB* gene and integrated in single copy by homologous recombination into the *panBCD* locus of *B. subtilis* strains RL-1 and PY79, replacing the wild-type operon. This was accomplished in two steps. First a deletion-substitution that replaced about two thirds of the *panB* coding region with a Gram-positive spectinomycin resistance gene (*spec*) was integrated at *panB* to yield Spec⁻, pantothenate auxotrophs. These intermediate strains were than transformed with the *panBCD* expression cassettes of pAN004, pAN005, and pAN006 after ligating them to a DNA fragment containing chromosomal sequences just upstream of *panB*. Selection of the incoming cassette was for pantothenate prototrophy. The resulting strains were named PA221, PA222 and PA223 (from RL-1), and PA235, PA232 and PA233 (from PY79), respectively. An example of a plasmid that contains the joined upstream sequence that is in the integrated strain in PA221 is pAN240 (see Figure 2). The nucleotide sequence of pAN240 is set forth as SEQ ID NO:76.

Polymerase chain reaction using appropriate primers was used to verify the correct chromosomal structures of these engineered strains. When extracts of strain PA221 were examined by SDS-PAGE, two proteins were found to be overexpressed.

One protein had an apparent molecular weight of 29,000 and the other protein appeared to be 39,000 daltons. The 29,000 dalton bands is presumably PanB (predicted molecular weight of 29,761). The larger protein band presumably represents PanC (predicted size 31,960 daltons).

The ability of these strains to produce pantothenate in test tube cultures
was assessed as follows. Each strain was grown in SVY medium supplemented with
f g/L α-ketoisovalerate (α-KIV) and f g/L β-alanine, to ensure that these precursors
were not limiting. Culture supernatants were autoclaved and assayed using the bioassay.
Relative to the parent strains, RL-1 and PY79, the engineered strains produced about 8to 30-fold more pantothenate, attaining 1 g/L pantothenate in some cases.

10

- 64 -

Table 2. Production of pantothenate by engineered B. subtilis strains in liquid test tube cultures grown in SVY medium with 5 g/L α -KIV and 5 g/L β -alanine.

Expt.	Strain	Promoter	RBS at panB	[pantothenate] mg/L
1	RL-1	Native	Native	30
	PA221	P ₂₆	RBS1	990 790
	PA222	P_{J5}	RBS1	250 250
	PA223	P ₂₆	RBS2	790 790
2	PY79	Native	Native	40
	PA235	P_{26}	RBS1	930 860
	PA221	P_{26}	RBS1	1100 1030

5 The P_{26} promoter was about 3- to 4-fold more effective than the P_{15} promoter, while RBS1 and RBS2 were roughly equivalent. Plasmids such as pAN004, pAN005, pAN006 can also be recombined as circles into the B. subtilis wild type panBCD locus by Campbell-type (single crossover) integration, selecting for chloramphenicol resistance at 5 mg/L. Strains obtained in this fashion produce about the same amount of pantothenate as strains PA221, PA222, and PA223, respectively. 10 pAN004 containing the P_{26} promoter, RBS1 and a low copy E. coli replicon, is depicted schematically in Figure 3A. The nucleotide sequence of plasmid pAN004 is set forth as SEQ ID NO:93. pAN006 containing the P_{26} promoter, RBS2 and a medium copy E. coli replicon, is depicted schematically in Figure 3B. The nucleotide sequence of plasmid pAN006 is set forth as SEQ ID NO:94. The nucleotide sequence of panBCD is 15 set forth as SEQ ID NO:59 and the predicted amino acid sequences of PanB, PanC and PanD are set forth as SEQ ID NO:24, SEQ ID NO:26 and SEQ ID NO:28, respectively. Methods for manipulating Bacilli are described, for example, in Harwood, C.R. and Cutting, S.M. (editors), Molecular Biological Methods for Bacillus (1990) John Wiley & 20 Sons, Ltd., Chichester, England, the content of which is incorporated herein by reference.

10

15

20

25

30

35

EXAMPLE II: Enhanced Production of a Panto-Compound Using Bacteria Overexpressing the *panE1* Gene Product – Ketopantoate Reductase.

This Example describes the cloning of the *B. subtilis panE1* gene and the generation of microorganisms overexpressing the *panE1* gene product.

Pan B. subtilis strains (e.g., B. subtilis mutants blocked in the synthesis of pantothenic acid) had previously been isolated, one of which was reported to be affected in ketopantoate reductase activity (Baigori et al. (1991) J. Bacteriol.

173:4240-4242). However, the mutations in these strains were incorrectly mapped to the purE-tre interval of the B. subtilis genetic map which does not contain the panE or panBCD genes. Furthermore as shown below, a panE mutant does not have a Pan phenotype as the ilvC gene product can substitute for the panE gene product in B. subtilis as in other bacterial strains such as E. coli. More recently, the S. typhimuruim panE gene has been located and determined to be allelic to apbA, a gene required for anaerobic purine biosynthesis (Frodyma et al. (1998) J. Biol. Chem. 273:5572-5576). E. coli carries a highly homologous gene at the same map location. Identification of the panE genes in E. coli and S. typhimurium was complicated by the fact that the ilvC gene product, acetohydroxy acid isomeroreductase, is also capable of carrying out the ketopantoate reductase reaction. As a result, pantothenate auxotrophy is not obtained unless both panE and ilvC are mutated.

To identify the *B. subtilis panE1* gene, the *B. subtilis* genome was searched using the protein sequence of *E. coli* or *S. typhimurium* ApbA (PanE), and two open reading frames were identified having homology to ApbA, named *ylbQ* and *ykpB*. These genes were renamed *panE1* and *panE2*, due to their proposed function in pantothenate biosynthesis. Both *panE1* and *panE2* were cloned as PCR products generated from RL-1 genomic DNA as a template. Both genes were disrupted by either a spectinomycin resistance gene (*spec*) or a chloramphenicol resistance gene (*cat*). The interrupted genes were each integrated by double crossover into PY79 to give PA240 (*ΔpanE1::spec*) and PA241 (*ΔpanE2::cat*). Neither of these strains were pantothenate auxotrophs when tested on pantothenate-free (PF) plates, although PA240 containing *ΔpanE1::spec* grew slightly more slowly on TBAB without added pantothenate than with a 1 mM pantothenate supplement. By comparison, a *ΔpanB::spec* strain does not produce single colonies on TBAB, presumably because *B. subtilis* has no active uptake system for pantothenate.

It was hypothesized that the *B. subtilis* gene, *ilvC*, could function for *panE* as had been shown for *E. coli*. Accordingly, the *panE1* and *panE2* disruptions

:5

15

were introduced into a strain, CU550, which is reported to be *trpC2 ilvC4 leuC124*. Both the single *panE1* and the double *panE1*, *panE2* disruptants were pantothenate auxotrophs on PF medium.

5 Table 3. Phenotypes of various panE1 and panE2 mutants on rich and defined media.

Strain	Medium	Growth*: - pan	+ pan
PY79	TBAB	+++	+++
	PF	++	++
PA240	TBAB spec	+	+++
	PF	++-	++
PA241	TBAB cam	+++	+++
	PF	++	++
CU550	TBAB	+++	+++
	PF	++	++
PA256	TBAB spec	-	+++
	PF	-	++
PA258	TBAB spec, cam	-	+++
	PF	-	++

^{*}Each "+" represents about 1 mm of colony diameter after overnight at 37°C.

Thus, mutating both *panE1* and *ilvC* results in pantothenate auxotrophy, while mutating only *panE1* does not, similar to what has been reported for *E.coli* and *S. typhimurium*.

Next, the quantitative effect of panE1 and panE2 knockouts in a pantothenate overproducing strain (PA235 described herein) was examined. The panE1 and panE2 disruptions were introduced into PA235, either singly or together to produce PA245 ($\Delta panE1$::spec), PA248 ($\Delta panE2$::cat) and PA244 ($\Delta panE1$::cat, $\Delta panE2$::spec). The effect of each mutation on pantothenate production was then tested in liquid test tube cultures.

Table 4. Pantothenate production by PA235 derivatives containing panE1 and panE2 disruptions.

Strain	[pan] mg/L	% of PA235
PA235	990	(100)
PA235	940	95
PA245	59	6
PA245	82	8
PA248	1060	106
PA248	1030	104
PA244	25	3
PA244	50	5

Thus, deletion analysis indicated that the *panE1* gene contributes to over 90% of the pantothenate production, while deletion of *panE2* did not have a significant effect on pantothenate production. It is therefore concluded that *panE1* accounts for most, but not necessarily all, of the ketopantoate reductase activity in *B. subtilis*. The rest of the ketopantoate reductase activity is predicted to be supplied by *ilvC*.

Having identified *panE1* as an important gene for pantothenate production, increased *panE1* expression was tested to determine whether it could enhance pantothenate production in strains such as PA221 or PA235. The *panE1* coding sequence was installed downstream of the *P*₂₆ promoter and RBS2 in a vector, pOTP61, designed to integrate and amplify at either the *bpr* locus (a non-essential protease gene) or at the locus of the cloned insert. The resulting plasmid, pAN236 (Figure 4) was transformed into PA221, selecting for resistance to tetracycline at 15 mg/L. The nucleotide sequence of pAN236 is set forth as SEQ ID NO:77. One transformant, named PA236 was chosen for further study.

PA236 was shown to overexpress a protein of about 31,000 daltons, which is close to the expected molecular weight of 33,290 daltons for *panE1* protein.

Briefly, whole cell extracts were prepared from PY79, RL-1, PA221,
PA221/pOTP61and PA236 (2 samples). Cell extracts were separated by gel electrophoresis and the gels were coomassie stained to visualize proteins. In cells engineered to overexpress *panE* (PA236-1 and PA236-2), a band was visible having an approximate molecular weight of ~31,000 daltons (as compared to molecular weight markers). Moreover, PA221 and PA236 expressed increased levels of a ~29,000 dalton

, **, ,**

5

15

25

band, corresponding to the *panB* gene product, and a ~39,000 dalton band, presumably corresponding the *panC* gene product. Furthermore, *E. coli* transformed with pAN006 (Figure 3B) expressed bands correlating to the *panB* and *panC* gene products and *E. coli* transfected with PAN236 expressed a ~31,000 dalton band corresponding to the *panE* gene product.

Next, PA236 was compared to PA221 carrying the empty vector pOTP61 for pantothenate production in liquid test tube cultures supplemented with 5 g/L β -alanine and 5 g/L α -KIV.

10 Table 5. Effect of overexpression of panE1 and panE2 on pantothenate production by engineered strains in liquid test tube cultures.

Strain	Additional Plasmid	Gene Overexpressed	[Pantothenate] mg/L
PA221	pOTP61	none	1,000
			940
PA236	pAN236	panE1	2,030
			2,050
PA238	pAN238	panE2	530
			680

Overexpression of *panE1* caused a two-fold increase in pantothenate production when compared to the parent strain (*e.g.*, to slightly over 2 g/L) whereas overexpression of *panE2* resulted in a strain that produced about 35% less pantothenate than the parent strain. The *panE1* nucleotide sequence and predicted amino acid sequence are set forth as SEQ ID NO:29 and SEQ ID NO:30.

EXAMPLE III: Enhanced Production of a Panto-Compound by Culturing Bacteria Overexpressing *panE1* or *panBCD* in the Presence of Valine.

The ability of valine to function as a media supplement (e.g., as a substitute for α -KIV) in strains engineered to overexpress the panBCD operon and panE1 was evaluated. Valine is closely related to α -KIV by transamination, is less expensive than α -KIV, and is commercially available in kilogram quantities. Valine was substituted for α -KIV in the standard liquid test tube cultures in SVY medium. The concentration of valine was varied from 5 to 50 g/L. Although valine at 5 g/L was

slightly less effective than α -KIV in promoting pantothenate production, valine at 10 or 20 g/L equaled or surpassed 5 g/L α -KIV in promoting pantothenate production.

EXAMPLES IV-X Generation of Microorganisms Capable of Producing Pantothenate in a Precursor-Independent Manner

B. subtilis strains such as PA221 and PA235 (engineered to overexpress panBCD) and PA236 (engineered to overexpress panBCD and panE1) need to be fed α-ketoisovalerate (α-KIV) (or valine) and aspartate (or β-alanine) to achieve maximal pantothenate production, as both these precursors are limiting for pantothenate synthesis. Accordingly, manipulated microorganisms were designed to eliminate the need to feed limiting precursors of pantothenate biosynthesis in the production of pantothenate. These strains are also useful in the production of various pantothenate biosynthetic pathway intermediates.

15 EXAMPLE IV: Generation of Microorganisms Capable of Producing Pantothenate in an Aspartate- (or β-Alanine) Independent Manner

The *panD* gene was cloned into *B. subtilis* expression vector pOTP61 to construct pAN423 (Figure 5). The nucleotide sequence of pAN423 is set forth as SEQ ID NO:78. The *Not*I restriction fragment containing *panD* was isolated from pAN423, self ligated and used to transform PA221. Transformants resistant to Tet¹⁵, Tet³⁰, and Tet⁶⁰ were isolated and saved for further analysis.

Six of the pAN423 transformants plus two control transformants were grown in SVY containing 5 g/l α -KIV with and without 10 g/l aspartate and then assayed for pantothenate production (Table 6).

Table 6. Effect of overproducing PanD on pantothenate production with and without added aspartate.

Culture* (PA221 transformants)	Asp (10 g/L)	TetR** (µg/ml)	OD550	[pan] (mg/L)
pOTP61-1	 	60	8.0	76
pOTP61-2	-	60	7.7	91
423#1-1	-	15	8.5	180
423#1-2	-	15	8.0	150
423#1-3	-	30	8.3	220
423#1-4	-	30	8.5	280
423#1-5	-	60	8.9	580
423#1-6		60	8.8	280

25

20

5

10

pOTP61-1	+	60	7.5	380
pOTP61-2	+	60	6.9	560
423#1-1	+	15	8.5	1200
423#1-2	+	15	8.6	1000
423#1-3	+	30	8.8	1200
423#1-4	+	30	9.0	1200
423#1-5	+	60	9.0	1200
423#1-6	+	60	9.0	1200

^{*}Test tubes cultures were grown in SVY + α -KIV (5 g/L) with Asp (10 g/L) where indicated.

The pAN423 transformants produced at least twice the amount of

pantothenate as the controls (*i.e.*, to a level at or near that which was obtained in earlier experiments by the addition of β-alanine to the culture medium). The data also show that in the absence of added aspartate, transformants containing additional copies of the *panD* gene expression cassette produce more pantothenate than the control transformants. One of the transformants, 423#1-5, produced about five times as much pantothenate as the controls. These results indicated that increased levels of PanD protein "pull" the conversion of available aspartate towards β-alanine, and that increasing *panD* gene expression can result in enhancement of pantothenate production both in the presence and absence of added aspartate.

Transformant 423#1-5 was re-named strain PA401 and studied further in shake flask fermentations. The shake flask medium was SVY with maltose instead of SVY with glucose. Results of shake flask experiments agreed well with test tube experiments during the first 24 hours. In shake flask experiments without the addition of β-alanine, PA401 produced approximately 1.5 g/l of pantothenate in 24 hours. Addition of β-alanine to the culture medium did not further improve pantothenate titers (Table 7), indicating that with this strain and these fermentation conditions, β-alanine is not limiting pantothenate production. In fact, when no β-alanine is fed, one can observe that PA401 is secreting β-alanine in significant amounts into the medium.

^{**}TetR = Approximate Tet-resistance of transformant

- 71 -

Table 7. Shake flask cultures with strain PA401 (panD) with and without \beta-alanine.

	Amino ao	cids (g/l)		24 hours	
Initial β-ala Added	β-ala	Val	рН	OD ₆₀₀	Pantothenate (g/l)
0	0.7	1.5	7.5	13.7	1.5
5 g/l	7.1	1.4	7.6	12.4	1.5

Each value represents the average of duplicate 250 ml baffled flasks containing 50 ml of medium, incubated at 37°C with shaking (200 rpm).

Base Medium: SVY with 10 g/l α -KIV, 30 g/l maltose

2% Inoculum: SVY with Tet 15 grown 24 hours.

EXAMPLE V: Engineering the *panD* gene for Further Increased Synthesis of Aspartate Decarboxylase and Enhanced Production of Pantothenate

This Example describes the generation of improved ribosome binding sites (RBSs) in the *panD* gene to increase the translation of *panD* mRNA.

Increasing the translation of the panD gene mRNA by generation of synthetic panD RBSs

15 The RBS (SEQ ID NO:88) used to express *panD* in pAN423 is a synthetic RBS and has been used to successfully produce other proteins in *B. subtilis* at a high level. However, it contains six mismatches when aligned to the "ideal" *B. subtilis* RBS (SEQ ID NO:45) (*e.g.*, an RBS having a sequence which is complementary to the 16S RNA sequence within the *B. subtilis* ribosome). (See *e.g.*, Table 1B, mismatches in bold). Two new RBSs were designed to more closely mimic the ideal RBS. These synthetic RBSs, named new design A (NDA) and new design B (NDB) (also referred to herein as RBS3 and RBS4), are set forth as SEQ ID NO:51 and SEQ ID NO:52 and are aligned with the ideal RBS in Table 1B.

Oligonucleotides corresponding to the top and bottom strands of each new RBS were synthesized, annealed, then used to replace the RBS in pAN420, generating plasmids pAN426 and pAN427. These constructions are illustrated in Figure 6. The presence of the NDA and NDB RBS in pAN426 and pAN427 was confirmed by DNA sequence analysis. Next, the *panD* genes from pAN426 and pAN427 were transferred to *B. subtilis* expression vector pOTP61 as shown in Figure 7, creating

pAN428 and pAN429. The nucleotide sequence of pAN429 is set forth as SEQ ID NO:79.

NotI restriction fragments lacking the *E. coli* vector sequences were isolated from pAN428 and pAN429, self-ligated, and used to transform strain PA221 to resistance to Tet¹⁵. Four isolates resistant to Tet⁶⁰ were picked from each transformation and assayed for pantothenate and β-alanine production along with PA221 transformed with the empty vector (pOTP61) and PA221 transformed with pAN423 (strain PA401) (see Table 8).

10 Table 8. Panthothenate production by test tube cultures of PA221 transformed with pAN428 and pAN429

Plasmid	_	Medium Supplements	OD550	Pan g/l	β -Ala g/l
pOTP61		α-KIV ⁵	10	UND	0.04
pAN423		α -KIV ⁵	10	0.4	0.04
pAN428-1	*	α -KIV $_{\epsilon}^{5}$	12	0.6	0.04
pAN428-2		α -KIV ⁵	11	0.5	0.03
pAN428-3		α -KIV ₅	11	0.3	0.03
pAN428-4		α -KIV ⁵	10	0.1	UND
pAN429-1		α -KIV $_{\varepsilon}^{5}$	12	0.6	0.04
pAN429-2		α -KIV ⁵	11	0.5	0.04
pAN429-3		α -KIV ⁵	11	0.6	0.05
pAN429-4	#	α -KIV ⁵	12	0.8	0.10
pOTP61		α -KIV ⁵ + Asp ¹⁰	11	0.5	0.08
pAN423		α -KIV + Asp α -KIV ⁵ + Asp 10	12	0.9	1.32
p/114-25		α-Kiv + Asp	12	0.5	1.52
pAN428-1	*	α -KIV ⁵ + Asp ¹⁰	12	0.8	1.97
pAN428-2		α -KIV ⁵ + Asp ¹⁰	12	0.8	1.51
pAN428-3		α -KIV ⁵ + Asp ¹⁰	12	0.9	1.02
pAN428-4		α -KIV ⁵ + Asp ¹⁰	11	0.8	0.30
pAN429-1		α -KIV $_{5}^{5}$ + Asp $_{10}^{10}$	12	0.8	1.78
pAN429-2		α -KIV ⁵ + Asp ¹⁰	12	0.8	1.66
pAN429-3		α -KIV ⁵ + Asp ¹⁰	12	0.8	1.78
pAN429-4	#	α -KIV ⁵ + Asp ¹⁰	13	0.8	2.28

UND: Below the limits of detection. * Renamed PA402 # Renamed PA403

15

i

When grown in medium supplemented with α -KIV at 5 g/l (α -KIV⁵), the pAN428-1 transformant and all four of the pAN429 transformants produced more pantothenate than did PA401, suggesting that these transformants contain higher levels of aspartate decarboxylase activity. When grown in medium supplemented with α -KIV⁵ and Asp¹⁰ none of the pAN428 or pAN429 transformants produced more pantothenate than PA401. However, the pAN428-1 transformant and all four of the pAN429 transformants produced significantly more β -alanine than did PA401. It is possible that the excess β -alanine produced from added aspartate causes inhibition of pantothenate production. Alternatively, β -alanine may accumulate because pantoate is limiting in these strains.

The strains that produced the highest level of β -alanine, the pAN428-1 and pAN429-4 transformants, were renamed PA402 and PA403, respectively. These two strains were grown in SVY medium supplemented with various intermediates and reassayed for pantothenate and β -alanine production. PA221 and PA401 were included as controls. The results of the assays are presented in Table 9.

Table 9. Pantothenate production of PA402 and PA403 in test tube cultures.

1

1

Strain	Medium Supplements	OD550	Pan g/l	β -Ala g/l	Val g/l
PA221	α-KIV ⁵	7.9	UND	UND	0.9
PA401	α-KIV ⁵	8.7	0.3	0.04	0.9
PA402	α-KIV ⁵	8.5	0.5	0.04	0.9
PA403	α-KIV ⁵	9.4	0.7	0.07	0.9
PA221	$\begin{array}{c} \alpha\text{-KIV}^5 + \text{Asp}^{10} \\ \alpha\text{-KIV}^5 + \text{Asp}^{10} \\ \alpha\text{-KIV}^5 + \text{Asp}^{10} \\ \alpha\text{-KIV}^5 + \text{Asp}^{10} \\ \alpha\text{-KIV}^5 + \text{Asp}^{10} \end{array}$	9.8	0.4	0.11	0.8
PA401		9.1	0.8	1.15	0.8
PA402		9.4	0.8	2.02	0.8
PA403		9.7	0.7	2.40	0.8
PA221	Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5	8.9	UND	UND	0.2
PA401		8.7	0.3	0.02	0.2
PA402		10.6	0.5	0.02	0.2
PA403		10.5	0.7	0.02	0.2
PA221	Pantoate ${}^{5}_{5}$ + Asp ${}^{10}_{10}$	9.5	0.4	0.06	0.2
PA401	Pantoate 5 + Asp	9.2	2.2	0.62	0.2

UND: Below the limits of detection.

When grown in medium supplemented with either α -KIV⁵ or Pantoate⁵, PA402 and PA403 produced significantly more pantothenate than did PA401. As before, even though PA402 and PA403 produced significantly more β -alanine than PA401 when grown in medium supplemented with α -KIV⁵ and Asp¹⁰, they did not produce a proportional increase in pantothenate. However, when grown in medium supplemented with Pantoate⁵ plus Asp¹⁰, both PA402 and PA403 produced significantly more pantothenate than PA401, about a 30% increase.

It can be concluded from these experiments that the improved NDA and NDB *panD* ribosome binding sites, engineered into pAN428 and pAN429, respectively, lead to increased levels of aspartate decarboxylase activity.

Increasing the translation of the panD gene mRNA by generation of synthetic panD RBSs within the panBCD operon

15

The native B. subtilis panD gene ribosome binding site (RBS) (SEQ ID NO:43), which is found in the $P_{26}panBCD$ operon cassette present in PA221 (and in other engineered pantothenate production strains described herein), is shown in Table 1C aligned with the ideal ribosome binding site (SEQ ID NO:47). The alignment shows mismatches between the native B. subtilis panD gene RBS, which is located within the 20 coding sequence for PanC, and the the ideal RBS. Three new RBSs (within the P26 panBCD operon cassette) were generated to increase translation of the panD gene mRNA and to yield increased synthesis of aspartate decarboxylase. These synthetic RBSs (termed NDI, NDII, and NDIII, also referred to herein as RBS5, RBS6 and RBS7, respectively) are set forth as SEQ ID NO:55, SEQ ID NO:56 and SEQ ID NO:57, 25 respectively) and are included in Table 1C. It should be noted that although changes in the panD RBS within the panBCD operon also changes the C-terminal amino acid sequence of the PanC protein encoded by that operon, an alignment of known and suspected PanC protein amino acid sequences showed that the sequence of the last nine amino acids of the B. subtilis PanC protein could be altered without affecting any 30 conserved amino acid residues indicating that such changes should not reduce pantothenate synthetase activity or expression. The new RBSs were synthesized and incorporated into the P_{26} panBCD operon expression cassette as follows.

25

30

35

First, PCR primers were designed to contain the following elements: (1) a nucleic acid sequence encoding the first five amino acids of PanD up to and including a unique *Bsi*WI restriction site that had been previously introduced into *panD* by PCR; (2) a stop codon for *panC*, (3) at least one synthetic RBS; and (4) 30-39 bp of nucleic acid sequence having 100% identity with *panC* upstream of the *panD* RBS. The primers were named TP102, TP103, and TP104 and contain the NDI, NDII, and NDIII ribosome binding sites, respectively. These three primers were used in conjunction with the 5' primer TP101, which hybridizes near the start codon of *panC*, in three independent PCR reactions to generate the NDI, NDII, and NDIII PCR products. The PCR products were purified, digested with *Xba*I, then cloned into plasmid vector pASK-1BA3 which had been digested with *Xba*I and *Sma*I. The resulting plasmids were named pAN431, pAN432, and pAN433. The construction of pAN431 is illustrated in Figure 8 and is representative of all three plasmid constructions. The presence of the desired synthetic *panD* gene RBS in each new plasmid was confirmed by DNA sequencing.

Next, the modified *panC* genes containing the new *panD* RBSs were joined with the *panD* gene utilizing the unique *Bsi*WI restriction site. This was accomplished by isolating the appropriate *Nsi*I-*Bsi*WI restriction fragments from pAN431, pAN432, and pAN433 and ligating them with a 2395 bp *Nsi*I-*Bsi*WI restriction fragment from pAN420, which supplied the *Bsi*WI-modified *panD* gene.

These constructions resulted in plasmids pAN441, pAN442, and pAN443, respectively. A representative construction (pAN441) is illustrated in Figure 9. The nucleotide sequence of pAN443 is set forth as SEQ ID NO:80.

The new *panD* gene RBSs were then substituted into the *P*₂₆*panBCD* operon expression cassette as follows. First, a deletion-insertion mutation which removes the region of *panC* containing the *panD* RBS was created. This was constructed by digesting pAN430 with a mixture of *BspE1* and *BglII* and recovering the 4235 bp fragment which is now missing the 3' end of *panC* and the 5' end of *panD*. This fragment was ligated with an *AvaI-BamHI* restriction fragment from plasmid pECC4, which contains the chloramphenicol acetyl transferase (*cat*) gene. The 5' extension produced by *AvaI* digestion is compatible with that produced by *BspEI* while the *BglIII* and *BamHI* extensions are also compatible. The resulting plasmid was named pAN440, and its construction is illustrated in Figure 10.

The resulting deletion-insertion mutation was crossed into the P_{26} panBCD operon via homologous recombination by transforming PA221 with linearized pAN440 and selecting for resistance to chloramphenicol on Cam⁵ plates containing 1 mM pantothenate. Several transformants were tested, and were all found to require 1

mM pantothenate for growth, as expected. Two of these transformants were remaned PA408A and PA408B and were assayed for pantothenate production. Neither strain synthesized measurable quantities of pantothenate, even when grown in medium containing pantoate and β-alanine at 5 g/l, indicating that the strains are deficient in pantothenate synthetase activity. Next, the new panD RBSs were crossed into the P₂₆ panBCD operon by transforming PA408 with linearized pAN441, pAN442, and pAN443 plasmid DNA and selecting for growth on TBAB plates without pantothenate supplementation. A transformation with linearized pAN430 (including the native panD RBS) was included as a control and was expected to give rise to transformants identical to PA221 described herein. Four isolates from each transformation were assayed for pantothenate and β-alanine production in SVY medium supplemented with various intermediates (Tables 10 and 11).

Table 10. Pantothenate production of PA410 - PA413 in test tube cultures.

Strain	RBS	Medium Supplements	OD ₅₅₀	Pan g/l	β -Ala g/l
PA221	native	Pantoate ⁵	11	UND	UND
PA410-1 PA410-2 PA410-3 PA410-4	native	Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5	12 12 12 12	UND UND UND UND	UND UND UND UND
PA411-1 PA411-2 PA411-3 PA411-4	NDI	Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5	12 12 12 12	0.23 0.20 0.19 UND	UND UND UND UND
PA412-1 PA412-2 PA412-3 PA412-4	NDII	Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5	12 11 13 12	UND UND 0.18 0.18	UND UND UND UND

ď.	114 879
٠,	7
Ĩ.	Ñ
4;	Ų.
Ĭ.	ı
ď.	S. PY
÷,	
51	
1111, 50	72
144 1444 101	11.12 S. 12.13
(1111)	1,200 (1,2) (1,4)
STATE OF STATE OF	The state of the state of
in the thirt that it is the to	1,200 (1,2) (1,4)

10

PA413-1	Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5 Pantoate	12	0.18	UND
PA413-2		12	0.17	UND
PA413-3		12	0.16	UND
PA413-4		12	0.17	UND

UND: Below the limits of detection.

Table 11. Pantothenate production of PA410 - PA413 in test tube cultures.

Strain	RBS	Medium Supplements	OD ₅₅₀	Pan g/l	β- Ala g/l
PA221	native	Pantoate ⁵ + Asp ¹⁰	11	0.3	UND
PA410-1 PA410-2 PA410-3 PA410-4	native	Pantoate 5 + Asp 10 Pantoate 6 + Asp 10	12 12 12 12	0.4 0.4 0.4 0.4	UND UND UND UND
PA411-1 PA411-2 PA411-3 PA411-4	NDI	Pantoate 5_5 + Asp ${}^{10}_{10}$ Pantoate 5_5 + Asp ${}^{10}_{10}$ Pantoate 5_5 + Asp ${}^{10}_{10}$ Pantoate 5_5 + Asp ${}^{10}_5$	13 13 13 13	1.7 1.7 1.8 0.4	0.4 0.4 0.3 UND
PA412-1 PA412-2 PA412-3 PA412-4	NDII	$\begin{array}{c} \operatorname{Pantoate}^5_5 + \operatorname{Asp}^{10}_{10} \\ \operatorname{Pantoate}^5_5 + \operatorname{Asp}^{10}_{10} \\ \operatorname{Pantoate}^5_5 + \operatorname{Asp}^{10}_{10} \\ \operatorname{Pantoate}^5_5 + \operatorname{Asp}^{10}_{10} \end{array}$	13 12 12 12	0.4 0.4 1.6 1.5	UND UND 0.3 0.2
PA413-1 PA413-2 PA413-3 PA413-4	NDIII	$\begin{array}{c} \operatorname{Pantoate}_5^5 + \operatorname{Asp}_{10}^{10} \\ \operatorname{Pantoate}_5^5 + \operatorname{Asp}_{10}^{10} \\ \operatorname{Pantoate}_5^5 + \operatorname{Asp}_{10}^{10} \\ \operatorname{Pantoate}_5^5 + \operatorname{Asp}_{10}^{10} \end{array}$	13 13 13 13	1.6 1.6 1.7 1.7	0.3 0.4 0.4 0.4

UND: Below the limits of detection.

As expected from previous experiments using PA221, none of the transformants that contained the native *panD* RBS produced measurable quantities of pantothenate when grown in medium supplemented with pantoate. However, nine of the twelve transformants expected to contain modified *panD* RBSs produced significant

quantities of pantothenate (160-230 mg/l) under these conditions, indicating that they possess elevated levels of aspartate decarboxylase activity. When grown in medium supplemented with both pantoate and aspartate, these same nine transformants produced approximately four times more pantothenate than those with the native panD RBS. In addition, these nine transformants accumulated measurable quantities of β -alanine (230-410 mg/l). All transformants produced roughly equivalent quantities of pantothenate when grown in medium containing pantoate and β -alanine, demonstrating that each contains a functional pantothenate synthetase.

These data demonstrate that the synthetic *panD* RBSs are about four times more effective than the native *panD* RBS in directing translation of the *panD* gene mRNA and evidence the utility of such synthetic RBSs in enhancing pantothenate production. Additional approaches to increasing pantothenate production can include, for example, increasing the half-life of the *panD* gene mRNA, increasing the strength of the promoter for *panD* transcription and/or increasing the stability of the PanD protein.

15

30

35

10

EXAMPLE VI: Construction of Strains Containing an Integrated P_{26} panE1 Cassette without an Antibiotic Resistance Gene.

Example II describes the identification of the *B. subtilis panE1* gene that encodes the enzyme responsible for the majority of the ketopantoate reductase activity in *B. subtilis*. PA236 (containing the pAN236 plasmid) produced about twice as much pantothenate (2 g/l) as its parent strain, PA221 (1 g/l) in 24 hour SVY test tube cultures. PA236 was presumed to contain an amplified (~3 copies) integrated pAN236 plasmid based on selection for tetracycline resistance (the tetR gene product being encoded on the pAN236 plasmid in addition to the *P*₂₆ *panE1* cassette). Also useful in the methodologies of the present invention are strains that contain a single integrated unamplifiable copy of *P*₂₆ *panE1* at the *panE1* locus, for example, without an antibiotic resistance gene in the strain. Such a strain was generated as follows.

A plasmid named pAN251 was derived from pAN236 by inserting additional chromosomal sequences just upstream and just downstream from the P_{26} panE1 cassette. These additional sequences, which provide homology to allow integration of the P_{26} panE1 cassette at panE1 by double crossover, were obtained by PCR from chromosomal DNA as a template. pAN251 is shown in Figure 11. The nucleotide sequence of pAN251 is set forth as SEQ ID NO:81.

Next, a strain was constructed which allowed selection for the incoming P_{26} panE1 cassette. The strain included the following three components: (1) P_{26} panBCD; (2) ApanE1; and (3) ilvC, since both panE1 and ilvC must be mutated to have a Pan phenotype. The starting strain was CU550 (trpC2, ilvC4, leuC124). The P₂₆ panBCD cassette from PA221 chromosomal DNA was introduced in two steps to create strain PA290. Next, ApanE1::spec was transformed into PA290, using chromosomal DNA from strain PA240, to give strain PA294 (trpC2, ilvC4, leuC124, P26 panBCD, ∆panE1::spec), which is a strict pantothenate auxotroph. Finally, PA294 was transformed with plasmid pAN251, selecting for pantothenate prototrophy, to give strain PA303. This strain was expected to have the genotype trpC2, ilvC4, leuC124, P26 10 panBCD, P26 panE1. PA303 was checked for the correct chromosomal structure at the panE1 locus by PCR using primers that flank the P_{26} insertion just upstream of panE1. The PCR product from PA303 was of the expected size, with a concomitant loss of the PCR product from the wild type panE1 gene, consistent with having obtained the desired double crossover event. Furthermore, PA303 was tetracycline sensitive, which 15 is also consistent with the desired double crossover event, as opposed to a Campbelltype single crossover of the plasmids into the chromosome. The trp, ilv, and leu auxotrophies from the parent strain were all maintained in PA303.

In 24 hour liquid SVY test tube cultures, PA303 produced almost the same level of pantothenate as positive control PA236, and about twice as much as PA221, which does not contain engineered *panE1* as indicated in Table 12.

Table 12. Pantothenate production by 24 hr. test tube cultures of PA303 and controls grown in SVY plus 5 g/l α -KIV and 5 g/l β -alanine.

OD ₆₀₀	[pan] g/l
10.9	0.85
10.5	0.85
9.5	1.74
9.3	1.70
10.8	1.66
10.7	1.61
	10.9 10.5 9.5 9.3

25

15

20

25

30

EXAMPLE VII: Generation of Microorganisms Capable of Producing Pantothenate in an α-KIV (or Valine) Independent Manner

 α -ketoisovalerate (α -KIV) is a rate limiting intermediate for pantothenate production in certain strains deregulated for pantothenate synthesis. Addition of either α -KIV or valine at 5 g/l increases pantothenate production about 5-fold in test tube cultures with strains such as PA221. In order to alleviate the need to feed either α -KIV or valine, strains were engineered that have an increased capacity to synthesize α -KIV.

α-KIV is produced in *B. subtilis* from pyruvate by the sequential action of three enzymes encoded by four genes, *ilvB* and *ilvN*, *ilvC*, and *ilvD*. In a wild type *B. subtilis*, three of the genes (*ilvB*, *ilvN*, and *ilvC*) are the first three genes of the large *ilv-leu* operon. The fourth gene necessary for α-KIV synthesis, *ilvD*, is located by itself elsewhere on the chromosome. The *B. subtilis ilv-leu* operon is thought to be regulated only by leucine levels. Feeding of exogenous leucine reduces transcription of the *ilv-leu* operon by about 13-fold, probably by an attenuation mechanism (Grandoni *et al.* (1992) *J. Bacteriol.* 174: 3212-3219). The only known feedback regulation in the *ilv-leu* pathway is the inhibition of the *leuA* gene product by leucine.

As a first step to deregulate the synthesis of α -KIV, a copy of the *ilvBNC* region from the wild type *B. subtilis ilv-leu* operon was isolated by PCR, and installed adjacent to the P_{26} promoter and RBS2 on a vector, pOLL8, that was designed to integrate a single P_{26} expression cassette by double recombination at the *amyE* locus. The *amyE* gene encodes a nonessential α -amylase, and is a useful locus for installing expression cassettes. The resulting plasmid, pAN267, is illustrated in Figure 12. The nucleotide sequence of pAN267 is set forth as SEQ ID NO:82. pAN267 readily gave stable transformants by double crossover at the *amyE* locus of *B. subtilis* strains, as described in detail below.

Construction of pantothenate overproducing strains that are leucine prototrophs

Initially, a *B. subtilis* strain containing ilvC4 and $\Delta panE1$ was used to introduce a single copy of P_{26} panE1 into the chromosome without using an antibiotic resistance gene. The double mutant was required to select for the incoming P_{26} panE1 cassette because a $\Delta panE1$ mutation alone does not result in pantothenate auxotrophy. A strain named CU550 was obtained containing ilvC4 to be used as a basis for this type of strain construction. However, CU550 also contains a closely linked leuC124 mutation, so all strains derived from CU550 required leucine. Having shown that the combination of P_{26} panBCD and P_{26} panE1 was favorable for pantothenate production,

the next step was to reassemble this combination of two cassettes in a leucine prototroph.

Accordingly, the two cassettes were combined in two different strain backgrounds, RL-1 and PY79. To introduce chromosomal P26 panE1 into the PY79 and RL-1 strain backgrounds without using an antibiotic resistance gene, a strategy was used that did not rely on ilvC4. (The strategy took advantage of the observation that the ΔpanE1 mutation causes a pantothenate bradytrophy, manifested by relatively small colonies on TBAB (rich) plates). First, \(\Delta panB::cat \) and \(\Delta panE::spec \) were introduced into both strain backgrounds. Next, the resulting strains were transformed simultaneously with DNA from two strains, PA221 (P_{26} panBCD) and PA303 (P_{26} 10 panE1), selecting for Pan+ on TBAB plates. Colonies of two distinct sizes grew on the selective plates, with the larger size comprising about 2% of the colonies. The larger colonies were presumed to represented co-transformants that received both P_{26} panBCD and P_{26} panE1, and that the smaller colonies had received only P_{26} panBCD. Consistent with this prediction, the larger colonies had lost both Cam^r and Spec^r, while the smaller 15 colonies had lost only the cat gene, and retained the spec gene. Furthermore, a representative derivative of PY79 named PA327, and a representative derivative of RL-1, named PA328, both produced the elevated levels of pantothenate in test tube cultures which was about 1.6 to 1.7 g/l (Table 13).

Table 13. Pantothenate production of PA327, PA328, and controls from 24 hr test tube cultures grown in SVY plus 5 g/l α -KIV and β -alanine.

Strain	Background	P26 panE1 copy number	[pan] g/l	
PA221-1 PA221-2	RL-1 RL-1	0	0.92 0.95	
PA236-1 PA236-2	RL-1 RL-1	amplified (~3) amplified (~3)	1.60 1.73	
PA327-1 PA327-2	PY79 PY79	1 1	1.66 1.65	
PA328-1 PA328-2	RL-1 RL-1	1 1	1.61 1.91	

20

20

25

Thus, PA327 and PA328 were concluded to contain both P_{26} panBCD and P_{26} panE1, and were used for further constructions as described below. PCR analysis confirmed the presence of the two cassettes.

5 Installation of a stable P26 ilvBNC cassette into two lineages of pantothenate overproducing strains

Having constructed PA327 and PA328, derivatives of PY79 and RL-1 that contain *P26 panBCD* and *P26 panE1*, and that are Leu⁺, the next step was to introduce stable copies of *P26 ilvBNC*. This was accomplished by transforming PA327 and PA328 with plasmid pAN267, selecting for Spec^r. Screening by PCR showed that about 85% of the obtained transformants contain *P26 ilvBNC* integrated at *amyE* by double crossover. One transformant of PA327, named PA340, and one transformant of PA328, named PA342, were chosen for further study.

In test tube cultures grown in SVY medium plus 5 g/l β-alanine but without added α-KIV, both PA340 and PA342 gave the expected increase in pantothenate production over that of PA327 and PA328, to about 1.3 to 2 g/l (Table 14).

Table 14. Pantothenate and valine production by PA340 and PA342, both containing P_{26} ilvBNC in 24 hr test tube cultures grown in SVY with 5 g/l β -alanine and with or without 5 g/l α -KIV

Strain	Back- ground	OD ₆₀₀ - α-KIV	+α-KIV	[pan] g/l - α-KIV	+ α-KIV	[val] g/l - α-KIV	+ α-KIV
PA340-1 PA340-2	PY79 PY79	11.8	7.1 7.5	2.02	2.10 2.03	0.38 0.40	0.90 0.91
PA342-1 PA342-2	RL-1 RL-1	10.2 9.6	8.0 9.2	1.29	1. 89 2.04	0.27 0.21	0.78 0.79

The two new strains also gave a slight increase in valine secretion, indicating that the *ilvBNC* genes had been deregulated. However, when the same strains were grown with 5 g/l α -KIV added, a further increase in pantothenate production occurred from PA342, suggesting that α -KIV was still rate limiting in this strain background. Similar results, only with more growth and hence higher pantothenate levels, were seen in shake flask cultures (Table 15).

10

15

25

Pantothenate and valine production by PA340 and PA342, both Table 15. containing P₂₆ ilvBNC in 24 hour shake flask cultures grown in SVY with 5 g/l β -alanine and with or without 5 g/l α -KIV.

Strain	Back- ground	OD ₆₀₀ - α-KIV	+ α-KIV	[pan] g/l - α-KIV	+ α-KIV	[val] g/l - α-KIV	+ α-KIV
PA327	PY79	21	22	0.6	3.0	0.5	1.3
PA340-1	PY79	20	20	3.5	4.1	1.0	1.9
PA340-2	PY79	22	19	3.0	2.1	0.8	1.4
PA328	RL-1	20	16	1.4	2.7	0.6	1.3
PA342-1	RL-1	17	16	3.3	3.6	0.9	1.6
PA342-2	RL-1	18	18	3.1	4.2	0.8	1.4

EXAMPLE VIII: Increasing panD Copy Number in Strains Engineered to Overproduce panE1 and the ilvBNC Gene Products Enhances Pantothenate **Production**

Experiments where β -alanine was fed to cultures of engineered B. subtilis strains consistently showed that β -alanine was a rate limiting intermediate in pantothenate synthesis. The effect of adding additional copies of panD on pantothenate production in PA340 and PA342 was examined. Strains PA340 and PA342 were transformed with chromosomal DNA isolated from PA401 with selection on plates containing 15 µg/ml of tetracycline (Tet15 plates). Transformants derived from each parent were patched onto Tet⁶⁰ plates to identify those which were likely to contain multiple copies of the expression cassette. Twelve transformants from each transformation which grew on Tet⁶⁰ were streaked for single colonies on this medium and then assayed in SVY medium test tube cultures for pantothenate production. One transformant from each group was found to produce greater than 300 mg/l pantothenate in 24 hours. These two transformants were saved and named PA404 (PA340 strain 20 background) and PA405 (PA342 strain background). Both strains were resistant to spectinomycin, indicating that the P_{26} ilvBNC expression cassette was still present at amyE. PCR analysis of chromosomal DNA isolated from each strain confirmed that the deregulated panE1 gene had also been retained.

Next, PA404 and PA405 were evaluated in shake flask cultures which were grown in SVY medium containing maltose as the carbon source and supplemented with various intermediates. The cultures were grown for 24 and 48 hours and then assayed for pantothenate, β -alanine, and valine production. The results of this

experiment are presented in Table 16. Analogous shake flask culture data for the parent strains (PA340 and PA342) are included in the tables for comparison.

Table 16. Pantothenate production by PA404 and PA405 in shake flask cultures after 24 hours

Strain	Medium Supplements	OD ₆₀₀	Pan g/l	β -Ala g/l	Val g/l
PA340 PA404 PA342 PA405	none none none	20 22 19 19	0.4 1.8 0.3 1.4	<0.1 <0.1 0.2 0.4	1.0 0.7 0.7 0.5
PA340 PA404 PA342* PA405*	β-alanine ⁵ β-alanine ⁵ β-alanine ⁵ β-alanine ⁵	18 18 17 19	3.6 2.8 3.3 1.3	3.2 5.1 3.3 6.5	0.6 0.7 0.5 0.6

Values are the average of duplicate flasks except where indicated by *.

In the absence of any medium supplementation, PA404 and PA405 made four to five times more pantothenate in 24 hours compared to their isogenic parent strains (Table 16). The supply of β-alanine was clearly limiting in the parent strains PA340 and PA342. Addition of amplified *P26 panD* greatly increased the supply of β-alanine.

EXAMPLE IX: Deregulation of the *B. subtilis ilvD* Gene Enhances Pantothenate Production

To deregulate expression of the *ilvD gene*, standard procedures (described above) were used to integrate the constitutive P_{26} promoter and an artificial ribosome binding site, RBS2, just upstream of the *ilvD* coding region. The *ilvD* gene maps by itself, unlinked to the *ilvBNC* operon. First, a 2.4 kb region of the RL-1 chromosome that contains the *ilvD* coding region and 730 bp of upstream sequence was cloned by PCR into a low copy (about 15 per *E. coli* cell) vector called pOK12, to give plasmid pAN257, shown in Figure 13.

15

20

20

Taking advantage of a natural *Eco*RI site just upstream of the native *ilvD* gene promoter, and a natural *Nco*I site at the *ilvD* start codon, an artificial sequence containing *P*₂₆ and RBS2 was inserted into pAN257 to give pAN263 (Figure 14). The nucleotide sequence of pAN263 is set forth as SEQ ID NO:83. In parallel with this construction, the *cat* gene was also inserted into pAN257, between the same upstream *Eco*RI site and a *BgI*II site in the middle of the *ilvD* coding region, to give pAN261, which is deleted for a large portion of the *ilvD* gene (Figure 15). Using pAN261 and pAN263, the *P*₂₆ *ilvD* cassette could then be installed in the *B*. *subtilis* chromosome in two steps. In the first step, pAN261 is introduced by transformation, selecting for chloramphenicol resistance, and then confirming an Ilv phenotype. In the second step, pAN263 is introduced, selecting for Ilv⁺, checking for chloramphenicol sensitivity, and confirming correct local structure by PCR.

pAN261 was first transformed into strain RL-1 (highly competent) to give strain PA343 ($\triangle ilvD::cat$), and then chromosomal DNA from PA343 was used to transform PA340 and PA342 to $\square ilv$ auxotrophy, yielding strains named PA348 and PA349, respectively. Chromosomal DNA is inherently more efficient than monomeric plasmid in transforming *B. subtilis*. Similarly, pAN263 DNA was transformed into PA343 (moderately competent) to give strain PA345 (P_{26} ilvD), and then PA345 chromosomal DNA was used to transform PA348 and PA349 to $\square ilv$ prototrophy, yielding strains PA374 and PA354, respectively.

As predicted, PA374 and PA354 gave further increases in pantothenate production, to about 2.5 to 2.9 g/l, in test tube cultures grown in SVY plus 5 g/l β -alanine (Table 17).

25 Table 17. Pantothenate and valine production by PA374 and PA354, containing P_{26} ilvD, and controls, in 24 hr test tube cultures grown in SVY with 5 g/l β -alanine and with or without 5 g/l α -KIV.

			OD ₆₀₀		[pan] g/l		[val] g/l	
Strain	Back- ground	ilvD status	α-KIV -	+	α-KIV -	+	α-KIV -	+
PA340	PY79	w.t.	9.2	9.0	2.14	2.23	0.38	0.90
PA348	PY79	ilvD::cat	11.7	10.0	0.19	2.23	0.19	0.91
PA374-1 PA374-2	PY79 PY79	P ₂₆ ilvD P ₂₆ ilvD	9.1 8.2	7.3 7.7	2.93 2.99	2.40 2.36	0.58 0.60	0.87 0.95

PA342	RL-1	w.t.	10.2	8.0	1.29	1.89	0.27	0.78
PA349	RL-1	w.t. ilvD::cat	8.1	7.7	0.17	1.87	0.22	0.88
PA354-1 PA354-2	RL-1 RL-1	P ₂₆ ilvD P ₂₆ ilvD	9.6 7.5	9.6 8.2	2.57 2.48	2.03 2.24	0.65 0.64	1.23 0.97

- 86 -

In the absence of added β-alanine, strains PA374 and PA354 produced only about 0.2 g/l pantothenate in test tube cultures, indicating that PanD activity is significantly rate limiting.

To alleviate this limitation, the amplifiable P_{26} panD cassette from strain 5 PA401 was installed. PA401 chromosomal DNA was transformed into PA374 and PA354, selecting for Tet^r at 15 mg/l, to yield strains PA377 and PA365, respectively. After transformants were obtained, the strains were streaked on plates containing 30 and 60 mg/l tetracycline to reamplify the copy number of the P_{26} panD cassette integrated at the bpr locus. In test tube cultures grown in SVY without α -KIV or β -alanine, a 10 substantial improvement in pantothenate titers over those of PA374 and PA354 was obtained (Tables 18 and 19).

Table 18. Pantothenate production by PA365, containing amplified P₂₆ panD, and controls, in 24 and 36 hr test tube cultures grown in SVY-glucose without β -alanine or α -KIV.

		OD ₆₀₀	<u> </u>	[pan] g/l	ı] g/l	
Strain	Relevant genotype	24 hrs.	36 hrs	24 hrs.	36 hrs.	
PA342-1-1	w.t. <i>ilvD</i>	11.7	8.8	b.d.	0.27	
PA342-1-2	w.t. <i>ilvD</i>	12.8	8.8	b.d.	0.26	
PA354-1-1	P ₂₆ ilvD	n.d.	11.0	n.d.	0.19	
PA354-1-2	P ₂₆ ilvD	n.d.	8.4	n.d.	0.20	
PA365-1	P26 ilvD, P26 panD	9.8	10.0	1.01	2.07	
PA365-2	P26 ilvD, P26 panD	9.9	10.4	0.96	2.09	

n.d. = not determined; b.d. = below detection

Table 19. Pantothenate production by PA377, containing amplified P_{26} panD, and controls, in 27 hr test tube cultures grown in SVY-glucose or SVY-maltose, without α -KIV, and with or without β -alanine.

		OD_{600}			
	Relevant genotype	- β-ala	+ β-ala	- β-ala	+ β-ala
Strain		Glucose	Glucose	Maltose	Maltose
PA374-1	P ₂₆ ilvD	9.4	9.8	7.0	6.4
PA374-2	P ₂₆ ilvD	9.2	9.6	6.6	6.3
		1			
PA377-1	P26 ilvD, P26 panD	10.0	7.6	7.2	6.1
PA377-2	P ₂₆ ilvD, P ₂₆ panD	10.5	7.8	9.4	5.4
		•			
		[pan] g/l			
	Relevant genotype	[pan] g/l	+ β-ala	- β-ala	+ β-ala
Strain	Relevant genotype		+ β-ala Glucose	- β-ala Maltose	+ β-ala Maltose
Strain	Relevant genotype	- β-ala	•	•	•
Strain PA374-1	Relevant genotype P26 ilvD	- β-ala	•	•	•
		- β-ala Glucose	Glucose	Maltose	Maltose
PA374-1	P ₂₆ ilvD	- β-ala Glucose	Glucose 2.76	Maltose 0.14	Maltose
PA374-1	P ₂₆ ilvD	- β-ala Glucose	Glucose 2.76	Maltose 0.14	Maltose

In SVY with glucose, an increase in pantothenate production can still be achieved by feeding 5 g/l β -alanine suggesting that increasing *panD* expression further might increase pantothenate production. In SVY with maltose, no further increase in pantothenate was obtained by feeding β -alanine suggesting that β -alanine and/or aspartate synthesis is suppressed by glucose. Strains PA377 and PA365 have been evaluated in 10 liter fermentors, where they typically produce above 20 g/l pantothenate in 48 hours without supplemental β -alanine and α -KIV or valine, described in detail below.

15

20

10

5

EXAMPLE X: 10 liter Fermentations of Pantothenate-Producing Microbes

Engineering of the P_{26} ilvBNC and P_{26} ilvD cassettes to give strains PA342 and PA354 allowed the production of 22 and 26 g/l of pantothenate, respectively, without the addition of valine or α -KIV to the fermentation medium (Table 20). At 48 hours, both strains had secreted about 0.5 g/l of valine into the medium.

Table 20. 10-liter fermentations of five pantothenate overproducing strains.

Strain	Medium	Feed 40% Glucose plus	OD 600 48 hr	Valine 48 hours g/l	β-ala 48 hr g/l	Pa 36 hr	ntothena g/L 48 hr	te 72 hr
PA 236	SVYG	50 g/l β-ala 25 g/l α-KIV	108	added	added	16	19	21
PA 342	SVYG	50 g/l β-ala	92	0.5	added	17	22	
PA 354	SVYG	50 g/l β-ala	90	0.5	added	19	26	
PA 365	SVYG	25g/l YE	77	0.85	0.4	18	21	27
PA 377	SVYG	25g/l YE	85	1.5	0.5	18	22	31
PA 377	PFMG	25g/l YE	96	0.8	0.4	19	25	29
PA377	PFMG	-	71	0.7	0.1	16	21	-

5 Pantothenate synthesis in fermentors

With the addition of the P_{26} panD cassette to strains PA354 and PA374 to create strains PA365 and PA377, neither β -alanine nor α -KIV needed to be added to the fermentors. Strain PA365 produced 21 g/l pantothenate in 48 hours and 27 g/l in 72 hours with no precursors added to the medium (Table 20). PA377 was somewhat better, producing 18 g/l of pantothenate in 36 hours, 22 g/l in 48 hours, and 31 g/l in 72 hours). Valine was measured at 0.85 and 1.5 g/l for strains PA365 and PA377, respectively, at 48 hours in SVYG medium. Strain PA377 maintained valine between 1-1.5 g/l throughout most of the fermentation and β -alanine between 0.2 and 0.5 g/l.

Strain PA377 was further evaluated in 10-liter fermentors in yeast extract based PFMG medium. Pantothenate yields in PFMG and SVYG medium were similar. In PFMG, PA377 produced 19 g/l of pantothenate in 36 hours, 25 g/l in 48 hours, and 29 g/l in 72 hours. In SVYG, PA377 produced 18 g/L pantothenate in 36 hours, 22 g/L in 48 hours and 31 g/L in 72 hours (Table 20).

EXAMPLE XI: Converting Strain PA377 to a Tryptophan Prototroph

PA377 (Trp⁻) was transformed to Trp⁺ using chromosomal DNA from PY79 to give strain PA824. After re-amplification of the $P_{26}panD$ casette, PA824 was compared to PA377 for pantothenate production in test tube cultures grown in SVY glucose with or without 5 g/L β -alanine (Table 21).

Table 21: Trp^+ derivatives of PA377: Pantothenate production in 48 hour test tube cultures grown in SVY glucose, $\pm \beta$ -alanine

		OD_{600}		[pan] g/L	
Strain	trpC donor	- β-alanine	+ β-alanine	- β-alanine	+ β-alanine
PA377-1	RL-1	8	8	1.5	3.4
PA377-2	RL-1	8	9	1.6	3.6
PA824-1 PA824-2	PY79 PY79	12 11	10 11	0.7	3.7 4.9

The Trp+ strains grew to slightly higher densities than PA377. In the absence of exogenous β-alanine, all of the strains produced similar levels of pantothenate, while with the addition of β-alanine, the Trp+ derivatives produced somewhat more pantothenate.

Fermentor studies with PA824

PA824 was evaluated in CF3000 Chemap 14 liter vessels with 10 liter working volumes. Formulations for two of the media used in the fermentors are given in Tables 22 and 23.

Table 22: Formulation for PFMG-5 medium

BATCH

	MATERIAL	g/L (final [])
1	Amberex 1003	10
2	Na Glutamate	5
3	(NH ₄) ₂ SO ₄	8
4	MAZU DF 37C	2.5

Added After Sterilization and Cool Down

1	KH ₂ PO ₄	10
2	K ₂ HPO ₄ ·3H ₂ O	20
1	Glucose	20
2	MgCl ₂ ·6H ₂ O	1
3	CaCl ₂ ·2H ₂ O	0.1
1	Sodium Citrate	1
2	FeSO ₄ ·7H ₂ O	0.01
3	SM-1000X	1.0 ml
	H_2O	qs to 6000 ml

5

FEED

	MATERIAL	g/L
1	Glucose	600
2	CaCl ₂ ·2H ₂ O	0.6
	$_{ m H_2O}$	qs to 3000 ml

Table 23: Formulation for SVY-4 medium

BATCH

MATERIAL	g/L (final [])
Veal Infusion	25
Yeast Extract	5
Na Glutamate	5
(NH ₄) ₂ SO ₄	4
MAZU DF 37C	2.5
	Veal Infusion Yeast Extract Na Glutamate (NH ₄) ₂ SO ₄

Added After Sterilization and Cool Down

1	KH ₂ PO ₄	10
2	K ₂ HPO ₄ ·3H ₂ O	20
1	Glucose	20
		20
2	MgCl ₂ ·6H ₂ O	1
3	CaCl ₂ ·2H ₂ O	0.1
1	Sodium Citrate	1
2	FeSO ₄ ·7H ₂ O	0.01
3	SM-1000X	1.0 ml
	H ₂ O	qs to 6000 ml

5

FEED

	MATERIAL	g/L
1	Glucose	600
2	CaCl ₂ ·2H ₂ O	0.6
	$_{ m H_2O}$	qs to 3000 ml

All fermentations were glucose limited fed batch processes. Immediately after inoculation, agitation was set at 200 rpm. The initial batched 2% glucose was

30

35

consumed during exponential growth. Afterwards, glucose concentrations were maintained between 0.2 and 1.0 g/L by continuous feeding of a 60% glucose solution. The variable rate feed pump was computer controlled and linked to the dissolved oxygen concentration [pO₂] in the tank by an algorithm. When the [pO₂]fell to 30%, computer control began to automatically adjust the agitation rate to maintain a dissolved oxygen concentration between 25 and 30% [pO₂]. Computer control and data recording were by Braun MFCS software.

In one study, PA284 was grown in fermentors at two temperatures (40°C and 43°C) in the medium described in Table 22. Results of two experiments demonstrated that the highest pantothenate titers at early time points were produced at 43°C. The cell mass approached 150 optical density units at OD₆₀₀ and 56 hours at 43°C, and the pantothenate titers were 21 g/L, 28 g/L and 36 g/L at 36, 48 and 72 hours respectively. In the parallel fermentation at 40°C, the cell mass approached 120 optical density units at OD₆₀₀ and 56 hours, and the pantothenate titers were 18 g/L, 26 g/L and 37 g/L at 36, 48 and 72 hours, respectively.

In another study, PA824 was grown in a fermentor at 43°C in the medium described in Table 23. The cell mass exceeded 160 optical density units at OD₆₀₀ and 36 hours, and the pantothenate titers were 23 g/L, 34 g/L, 37 g/L and 40 g/L at 24, 36, 48 and 60 hours, respectively. In other fermentations, increasing the amount of trace elements in the glucose feed (*e.g.*, increasing the concentration of SM from 1X to 2X) resulted in even higher titers of pantothenate.

25 EXAMPLE XII: Identification and characterization of the *B. subtilis coaA* gene product

The annotated version of the *B. subtilis* genome sequence available on the "Subtilist" web site contains no gene labeled as *coaA*. However a homology search using the protein sequence of *E. coli* pantothenate kinase as a query sequence gave a good match with *B. subtilis* gene *yqjS*, which is annotated as "unknown; similar to pantothenate kinase." This gene appears to be the penultimate gene in an operon containing five open reading frames (Figure 18). Two of the open reading frames encode proteins which are similar to D-serine dehydratase and to "ketoacyl reductase"; the other two have no known homologies. For the open reading frame corresponding to *coaA*, there are three possible start codons; each having a possible ribosome-binding site

30

35

(RBS) associated with it. The three potential *coaA* ORFs were named *coaA1*, *coaA2*, and *coaA3*, from longest to shortest.

All three potential coaA open reading frames were cloned along with their respective RBSs by PCR followed by ligation into expression plasmid pAN229. pAN229 is a low copy vector in E. coli that provides expression from the SP01 phage P_{15} promoter and can integrate by single crossover at bpr with tetracycline selection. A representative resulting plasmid, pAN281, is shown in Figure 19.

To determine if the cloned putative *coaA* ORFs actually encode a pantothenate kinase activity, several isolates of all three plasmids were transformed into the *E. coli* strain YH1, that contains the *coaA15(Ts)* allele. Transformants were streaked to plates incubated at 30° and 43°C to test for complementation of the temperature sensitive allele. All isolates of all three *coaA* variants, except for one isolate of pAN282, complemented well at 43°C, indicating that all three plasmid constructs encode an active pantothenate kinase. Accordingly, it can be concluded that the *B. subtilis yqjS* open reading frame codes for an active pantothenate kinase.

EXAMPLE XIII: Deletion of the coaA gene from the B. subtilis genome

The *coaA* gene of *B. subtilis* (*yqjS*) was deleted from the chromosome of a *B. subtilis* strain by conventional means. The majority of the *coaA* coding sequence was deleted from a plasmid clone and replaced by a chloramphenicol resistance gene (*cat*), while leaving approximately 1 kb of upstream and downstream sequence to allow homologous recombination within the chromosome, to give plasmid pAN296 (see Figure 17). pAN296 was then used to transform a *B. subtilis* strain (PY79), selecting for chloramphenicol resistance. The majority of transformants result from a double crossover event that effectively substitutes the *cat* gene for the *coaA* gene. The transformed strain containing the *coaA* deletion – *cat* insertion grew normally due the presence of a second *B. subtilis* pantothenate kinase encoding gene described herein.

EXAMPLE XIV: Identification and characterization of a second *B. subtilis* gene encoding pantothenate kinase activity

As described in detail in the instant specification, in order to maximize pantothenate production, it is necessary to restrict the flow of pantothenate toward Coenzyme A (CoA), for example, by reducing the activity of pantothenate kinase, the first enzyme in the pathway from pantothenate to CoA. After finding that deletion of

15

the coaA gene from the chromosome of B. subtilis is not a lethal event (see Example XIII), it was concluded that B. subtilis must contain a second gene that encodes an active pantothenate kinase, since pantothenate kinase is an essential enzyme activity.

A second pantothenate kinase-encoding gene was identified by complementing the E. coli strain YH1 (coaA15(Ts)) with a B. subtilis gene bank and selecting for transformants that were able to grow at 43°C. Found among the transformants were two families of plasmids that had overlapping restriction maps within each family, but not between the families. As expected, the restriction map of one family was identical to that predicted from the B. subtilis genome sequence for the homologue of the E. coli coaA gene (which we named coaA also, see above) and surrounding sequences. The other family had a restriction map that was completely non-overlapping with the first.

DNA sequencing of the ends of the cloned inserts from the second family showed that the clones came from a region of the B. subtilis chromosome that includes the 3' end of the ftsH gene, the 5' end of the sul gene, and all of the yacB, yacC, yacD, cvsK, pabB, pabA and pabC genes. None of the open reading frames of these cloned inserts showed homology to any known pantothenate kinase sequences, either prokaryotic or eukaryotic.

Several deletions were created through the B. subtilis genomic sequences in the cloned inserts. Each deletion was tested for complementation of the E. coli 20 temperature sensitive pantothenate kinase. In particular, a deletion that removed all DNA between a Stu I site in the cloning vector and a Swa I site in the yacC gene, leaves vacB as the only intact open reading frame in the cloned insert (see Figure 21). This deleted plasmid still complemented the E. coli pantothenate kinase mutant. However, another deletion that removed DNA from the Swa I site in yacC through a Bst1107I site 25 in the (already truncated) ftsH gene, could not complement the E. coli pantothenate kinase mutant. From these results, it was concluded that the yacB open reading frame was responsible for the complementation activity. To confirm that yacB is a pantothenate kinase gene, the yacB ORF plus 112 base pairs of downstream flanking sequence was amplified by PCR in two independent reactions and cloned downstream of 30 a constitutive promote to give plasmids pAN341 and pAN342 (Figure 22). Both pAN341 and pAN342 complemented the defect in YH1 at 44°C, while a control plasmid, which has the same backbone, but expresses panBCD instead of yacB did not. This confirmed that the yacB open reading frame was responsible for the

35 complementation of YH1.

15

20

As such, a novel gene that encodes pantothenate kinase activity in *B. subtilis* has been discovered that is not related by homology to any previously known pantothenate kinase gene. This gene has been renamed *coaX*, as a second, alternative gene that encodes an enzyme that catalyzes the first step in the pathway from pantothenate to CoaA. Deletion of *coaX* by methods described above for deleting *coaA*, in conjunction with reduction in the activity of the CoaA enzyme, provides a means to reduce pantothenate kinase activity to the desired level.

Several homologues of the *B. subtilis coaX* gene were identified by homology searching of various publically available databases using the published *yacB* (*coaX*) open reading frame sequence and predicted amino acid sequence (as set forth in SEQ ID NOs:84 and 85 respectively). In two cases (*Mycobacterium tuberculosis and Streptomyces coelicolor*) the homologous *coaX* genes are adjacent to, or almost adjacent to, pantothenate biosynthetic genes, consistent with these homologs having a role in pantothenate metabolism. The CoaX proteins show no homology to the CoaA family of pantothenate kinases, nor to the eukaryotic family of pantothenate kinases exemplified by PanK of *Saccharomyces cerevisiae*.

Alignment of the amino acid sequences of several bacterial CoaX homologs with the amino acid sequence predicted from translating the *B. subtilis yacB* ORF described in the published *B. subtilis* genome sequence revealed that the CoaX proteins from other bacteria contained additional amino acid residues at their carboxy-terminal ends. Moreover, these extensions beyond the end of the predicted amino acid for the *B. subtilis* gene product contained two relatively well conserved segments of sequence.

Translation of nucleotide sequences just downstream from the stop codon of the *B. subtilis yacB* ORF in a different reading frame revealed the existence of amino acid sequences very similar to the carboxy-terminal extensions of the other bacterial CoaX proteins. It is thus believed that an error exists in the published DNA sequence of the *B. subtilis yacB* ORF sequence that causes a frame shift leading to an artifactual downstream amino acid sequence and premature termination.

The PCR-generated sequences of *B. subtilis CoaX* in pAN341 and pAN342 (described above) contain enough downstream flanking sequence to encode the putative carboxy-terminal extension described above, which is consistent with the result that the clones were functional in the complementation assay. However when the 3' PCR primer was positioned to include only the shorter *yacB* ORF predicted from the published sequence, but not to include the putative carboxy-terminal extension, then the

resulting plasmids, pAN329 and pAN330 (similar in structure to pAN341 and pAN342; see Figure 22), did not complement the defect in YH1. This result supports the notion that the published yacB coding sequence contains a frame-shift error, and that the carboxy-terminal end of CoaX is necessary for pantothenate kinase activity. The predicted correct nucleotide sequence for B. subtilis coaX is set forth as SEQ ID NO:19 and the translated amino acid sequence is set forth as SEQ ID NO:9. A multiple sequence alignment of the CoaX amino acid sequences of B. subtilis and 11 homologues thereof is set forth in Figure 23.

10

EXAMPLE XV: Generation of mutant *coaA* genes encoding pantothenate kinase having reduced or temperature sensitive activities

This Example describes strategies for modifying the *coaA* gene (*i.e.*, by introducing point mutations) to reduce the activity of pantothenate kinase after *coaX* is deleted from the genome.

Cloning and sequencing of the temperature sensitive allele of the E. coli coaA gene.

Two E. coli strains, each exhibiting a different mutant CoaA phenotype, were obtained from the E. coli Genetic Stock Center. Strain DV62 contains the 20 coaA15(Ts) allele, and DV79 contains the coaA16(Fr) mutation. DV62 is temperature sensitive at 43°C and produces a pantothenate kinase that is temperature sensitive. DV79 was obtained by reversion of DV62 to temperature resistance, and it produces a temperature stable, feedback resistant pantothenate kinase activity. Since the DNA sequences of these alleles are not available in the literature, the coaA genes from the two 25 mutant strains were cloned by PCR and sequenced, in addition to a coaA gene from a strain that is wild type at the coal locus, MM294. The PCR primer at the 5' end was designed to include the start codon plus four bases upstream, and added an arbitrarily chosen ribosome binding site (RBS). The three PCR generated fragments were each ligated between the XbaI and BamHI sites of pAN229 to give pAN284 (from 30 coaA15(Ts)), pAN285 (from wild type coaA), and pAN286 (from coaA16(Fr)). pAN229 is a low copy E. coli vector that provides expression from the P_{15} promoter and that can integrate by single crossover at bpr in B. subtilis with tetracycline selection.

All three plasmids were transformed into the *E. coli* strain YH1 for complementation testing. All three plasmids complemented the temperature sensitive *coaA* mutation in *E. coli* YH1. It is presumed that the *coaA15(Ts)* gene in pAN284 is

20

25

30

35

probably significantly overexpressed relative to the normal chromosomal gene, such that the overproduction compensates for the temperature sensitive defect. Complementation of a defect by overproduction is a well-documented phenomenon in *E. coli*.

The *coaA* coding regions from pAN284, 285, and 286 were subcloned

into pGEM7 to give pAN306, 307, and 308, respectively, for DNA sequencing. As expected, the DNA sequence of the insert in pAN307 (from wild type *coaA*) matched the *coaA* sequence from the *E. coli* genome database (GenBankTM). The sequence from pAN306 contains a single base change that causes a S176L substitution (*i.e.*, a Ser → Leu substitution in the amino acid sequence set forth as SEQ ID NO:2). Interestingly, the DNA sequence of the pAN308 insert, derived from the feedback resistant strain, was identical to that derived from its temperature sensitive parent (represented in pAN306). This is in accord with the genetic data that indicates that the reversion of the temperature sensitive mutation occurred at a second site unlinked to the *coaA* gene.

The S176L mutation, predicted to cause the temperature sensitive defect in *E. coli* pantothenate kinase, changed a serine residue that is conserved in all known or suspected bacterial *coaA* encoded pantothenate kinases, including that of *B. subtilis* (see SEQ ID NO:3 and refer to alignment). Based on this, a serine to leucine change at the homologous residue in the *B. subtilis* pantothenate kinase is predicted to result in either a temperature sensitive enzyme or one which is less active. Accordingly, to produce a mutant *B. subtilis coaA* gene, this specific change was introduced into the *B. subtilis coaA* gene. The mutant version is installed in the chromosome of a *B. subtilis* strain deleted for *coaX*, for example, and the recombinant microorganism is checked for temperature sensitivity (*e.g.*, reduced growth at 43°C). The mutation is then installed into a pantothenate overproducing strain, preferably a strain deleted for the above mentioned *coaX* gene by standard methods to give strains favorable for pantothenate production in *B. subtilis*, *i.e.*, a strain that has reduced pantothenate kinase activity under typical fermentation conditions.

Additional coaA point mutations resulting in reduced pantothenate kinase activity

Of course it is expected that many other point mutations or combinations of more than one point mutation in *B. subtilis coaA* will also lead to reduced activity. Appropriate mutations can be generated by mutagenic polymerase chain reaction and *in vitro* recombination, and identified by screening for alleles that poorly complement the *E. coli coaA15(Ts)* mutant. An example of such a mutation of this type is a tyrosine to histidine substitution at amino acid 181 of B. *subtilis coaA*, generated by mutagenic

15

polymerase chain reaction (see SEQ ID NO:3 and first line of the alignment of Figure 24).

Isolate pAN282A was derived from the middle-sized B. subtilis coaA open reading frame described in Example XII. pAN282A complemented the E. coli coaA15(Ts) mutant very poorly, but nonetheless at a level that was detectable above background. As was done for the E. coli coaA clones, the open reading frame from pAN282A was subcloned into pGEM7 to give pAN303. The DNA sequence of the insert in pAN303 showed a single base change that led to a tyrosine to histidine amino acid change at the tyrosine corresponding to Y181 of SEQ ID NO:3. This tyrosine residue is conserved in all bacterial coaA genes/homologues present in GenBank (Figure 24). This tyrosine residue and the serine that is altered in the E. coli temperature sensitive pantothenate kinase described above are separated by only three amino acid residues in a region which is highly conserved in bacterial pantothenate kinases whereas the DNA sequence of a second isolate of the middle-sized open reading frame, from pAN282B, was identical to the wild type sequence from the B. subtilis genome sequencing project. The single base change found in pAN303 probably occurred during PCR amplification of the coaA gene. If this variant of coaA2 has sufficient residual biological activity in B. subtilis, it may be useful in the future for providing reduced pantothenate kinase activity.

20 A preferred plasmid that can serve as a basis for mutagenizing the *coaA* open reading frame is pAN294 (see *e.g.*, Figure 25 and Example XII). Briefly, mutagenic PCR is performed using pAN294 as a template and variants of *coaA* having reduced pantothenate kinase activity are screened as described above. Alternatively, mutations such as the one isolated in pAN282A can be installed into pAN294. The desired mutation is then introduced into the chromosome of a *B. subtilis* strain by transformation with the appropriate pAN294 derivative and selected for chloramphenicol resistance at 5 mg/L. Among the resulting transformants will be isolates that contain the desired mutation.

In a similar fashion, mutations that reduce the activity of the CoaX enzyme can be generated and identified, and such mutations used for optimizing pantothenate production by reducing CoA production as described above.

EXAMPLE XVI: Deleting the second pantothenate kinase gene, *coaX* gene from *B. subtilis*

With the knowledge gained above concerning the existence and nature of coaX, one can create a deletion of the coaX open reading frame from the B. subtilis chromosome that will remove the encoded activity, and that will not adversely affect the expression of the genes downstream from coaX. In such a deleted strain, the coaA gene will be the only gene that encodes pantothenate kinase.

To delete the *coaX* gene from *B. subtilis*, plasmid pAN336 (SEQ ID NO:92), which contains upstream and downstream homology for double crossover, was constructed with a kanamycin resistance gene replacing most of the *coaX* ORF (Figure 26). Strain PY79 was transformed to kanamycin resistance by pAN336, and an isolate confirmed to have resulted from a double crossover by PCR was named PA876. As predicted, deletion of *coaX* by itself is not lethal for *B. subtilis*. Furthermore,

15 chromosomal DNA from PA876 would not transform competent PA861 (PY79 ΔcoaA ::cat) to kanamycin resistance. These results indicate that it is the combination of Δ*coaA*::*cat* and Δ*coaX* :: *kan* that is lethal for *B. subtilis*, confirming that *B. subtilis* contains two unlinked genes that encode pantothenate kanase, *coaA* and *coaX*, and that either gene alone is capable of supplying sufficient pantothenate kinase for a normal rate of growth.

EXAMPLE XVII: Construction of a plasmid designed to allow directed mutagenesis of the *B. subtilis coaA* gene

In order to easily introduce mutated coaA genes into the B. subtilis chromosome, it was necessary to install an antibiotic resistance gene adjacent to the 25 coaA gene. This was accomplished by joining together in the vector pGEM5 three DNA fragments: (1) a 3.4 kb DNA sequence containing 2.5 kb of genomic sequence upstream from coaA and the coaA open reading frame(s); (2) a 1.1 kb DNA sequence containing a chloramphenicol resistance gene (cat); and (3) a 1.4 kb DNA sequence comprising a region downstream from the operon that contains coaA. The resulting plasmid, named 30 pAN294, effectively replaces the open reading frame yajT (the open reading frame just downstream from coaA) with the cat gene, with enough homology flanking both sides of the cat gene to allow double recombination into the B. subtilis chromosome (Figure 25). pAN294 was transformed into B. subtilis strain PY79, selecting for chloramphenicol resistance at 5 mg/l to give strains PA836 and PA837, which are presumably identical. 35 PA836 and 837 were checked by diagnostic PCR to show that the cat gene had

- 100 -

integrated by double crossover, as opposed to single crossover. PA836 and PA837 grow normally, leading to the conclusion that the open reading frame yqjT is not essential (*i.e.*, the yqjT open reading frame could be deleted from strains PA836 and PA837 with no significant effect on growth or pantothenate production). Thus, variant alleles (*i.e.*, mutations) of the coaA gane can be introduced into pAN294 and the resulting plasmids can be used to introduce the variant alleles into the chromosome of, for example, a B. subtilis strain.

EXAMPLE XVIII: Generation of mutant *coaX* genes encoding pantothenate 10 kinase having reduced or temperature sensitive activities

Mutant *coaX* genes are generated by introducing point mutations into the gene and testing the resulting mutants for the ability to complement the *E. coli* YH1 strain as described in Example XII. Preferred mutations in the *coaX* gene sequences are those that encode a substitution of a residue conserved among CoaX sequences from a variety of bacterial sources (*e.g.*, a conserved residue set forth in Figure 23). Alternatively, random mutations in the *coaX* gene sequence are generated by mutagenic PCR and *in vitro* recombination and identified by screening for alleles that poorly complement the *E. coli coaA15(Ts)* mutant.

Mutants so generated (*i.e.*, mutants having reduced *coaX* activity) can be further engineered such that the endogenous *coaA* gene is deleted (as described in Example XIII). CoaX reduced-activity mutants can also be further engineered to contain reduced-activity CoaA gene products as described in Example XV.

25 EXAMPLE XIX: Enhanced Production of Panto-Compounds Using Bacteria Having Deletions in One or More Pantothenate Biosynthetic Enzymes

If the desired panto-compound is not pantothenate, then an appropriate deletion of one or more of the pantothenate biosynthetic genes from a pantothenate overproducing strain will provide a strain that produces said desired panto-compound. In this example, the desired panto-compound is pantoate. Starting with, for example, strain PA236, PA313 or PA824 either one or both of the *panC* and *panD* genes is deleted. In another example, ketopantoate is the desired panto-compound. Starting with, for example strain PA244, PA245 or PA824 one, two or all of the *ilvC*, *panE1*, *panC* and *panD* genes are deleted from the starting strain. If β-alanine is the desired panto-compound, then *panB* and *panC* can be deleted, preferably in a fashion that leaves

an in frame fusion of a small portion of the 5' end of panB with a small portion of the 3' end of panC, from the strain PA221, PA235, PA245, or PA313. In all of the above-mentioned examples, the panto-compound producing strain will be a pantothenate auxotroph. Accordingly, the growth medium requires sufficient pantothenate for adequate growth. Vectors designed to overexpress panD as described above are then transformed into the above strains to further enhance β -alanine production.

The above-mentioned deletions are accomplished by methods well-known to those skilled in the art, for example, by insertion of an antibiotic resistance gene and removing sufficient sequence from the target gene(s) to inactivate said target gene(s). Alternatively, removal of targeted sequences is accomplished without simultaneous introduction of an antibiotic resistance gene in said target gene and then introduced by congression (co-transformation with any other appropriate selectable DNA sequence) followed by screening for the loss of function of said target gene by replica plating.

15

10

- 102 -

Table 24: Strains (and corresponding phenotypes) for panto-compound production

Name	Pheno type	Drug resist.	<i>panBCD</i> locus	panE locus	<i>ilvD</i> locus	amyE locus	bpr locus	Parent
PA221	Trp-		P26panBCD					
PA222			P ₁₅ panBCD					RL-1
PA235			P26panBCD					
PA236			P_{26} panBCD	P_{26} panE1				PA221
PA327	Trp-		P26panBCD	P26panE1				PA221
PA328	Trp-		P26panBCD	P26panE1				PA235
PA340	Trp-	Spc	P26panBCD	P26panE1		P26ilvBNC		PA327
PA342	Trp-	Spc	P26panBCD	P26panE1		P26ilvBNC		PA328
PA354	Trp-	Spc	P26panBCD	P26panE1	P26ilvD	P26ilvBNC		PA342
PA365	Trp-	Spc, Tet	P26panBCD	P26panE1	P26ilvD	P26ilvBNC	P26panD423	PA354
PA374	Trp-	Spc	P26panBCD	P26panE1	P26ilvD	P26ilvBNC		PA340
PA377	Trp-	Spc, Tet	P26panBCD	P26panE1	P26ilvD	P26ilvBNC	P26panD423	PA374
PA401	Trp-		P26panBCD				P26panD423	PA221
PA402	Trp-		P26panBCD				P26panD428	PA221
PA403	Trp-		P26panBCD				P26panD429	PA221
PA404	Trp-	Spc, Tet	P26panBCD	P26panE1		P26ilvBNC	P26panD423	PA340
PA405	Trp-	Spc, Tet	P26panBCD	P26panE1		P26ilvBNC	P26panD423	PA342
PA651	Trp-	Spc	P26panBC*D	P26panE1	P26ilvD	P26ilvBNC		PA374
PA284		Spc, Tet	P26'panBCD	P26panE1	P26ilvD	P26ilvBNC	P26panD423	PA377

Equivalents Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

- 103 -

What is claimed:

5

- 1. A method of producing a panto-compound comprising culturing a microorganism which overexpresses at least one *Bacillus* pantothenate biosynthetic enzyme under conditions such that the panto-compound is produced.
- 2. The method of claim 1, wherein the microorganism overexpresses at least one *Bacillus subtilis* pantothenate biosynthetic enzyme.
- 3. The method of claim 1 or 2, wherein the pantothenate biosynthetic enzyme is selected from the group consisting of ketopantoate hydroxymethyltransferase, pantothenate synthetase, aspartate-α-decarboxylase and ketopantoate reductase.
- 15 4. The method of any one of claims 1 to 3, wherein the microorganism overexpresses at least two pantothenate biosynthetic enzymes.
 - 5. The method of any one of claims 1 to 3, wherein the microorganism overexpresses at least three pantothenate biosynthetic enzymes.
 - 6. The method of any one of claims 1 to 5, wherein the panto-compound is selected from the group consisting of pantothenate, pantoate, ketopantoate and β -alanine.
- 7. A method of producing a panto-compound comprising culturing a ketopantoate reductase-overexpressing (KPAR-O) microorganism under conditions such that the panto-compound is produced.
- 8. The method of claim 7, wherein the panto-compound is 30 pantothenate or pantoate.
 - 9. The method of claim 7 or 8, wherein the ketopantoate reductase is bacterial-derived.
- The method of claim 7 or 8, wherein the ketopantoate reductase is derived from *Bacillus*.

- 104 -

- 11. The method of claim 7 or 8, wherein the ketopantoate reductase is derived from *Bacillus subtilis*.
- 12. The method of any one of claims 7 to 11, wherein the KPAR-O microorganism further overexpresses at least one pantothenate biosynthetic enzyme in addition to overexpressing ketopantoate reductase.
- 13. The method of claim 12, wherein the KPAR-O microorganism further overexpresses at least one of ketopantoate hydroxymethyltransferase,
 10 pantothenate synthetase and aspartate-α-decarboxylase.
- 14. A method of producing pantothenate in a manner independent of precursor feed comprising culturing an aspartate-α-decarboxylase-overexpressing (AαD-O) microorganism having a deregulated isoleucine-valine (*ilv*) pathway under conditions
 such that pantothenate is produced.
- 15. A method of producing at least 2 g/L pantothenate in a manner independent of aspartate or β-alanine feed comprising culturing an aspartate-α-decarboxylase-overexpressing (AαD-O) microorganism under conditions such that
 20 pantothenate is produced.
- 16. A method of producing at least 2 g/L pantothenate in a manner independent of valine or α-ketoisovalerate feed comprising culturing a microorganism having a deregulated isoleucine-valine (*ilv*) biosynthetic pathway under conditions such that pantothenate is produced.
- 17. A method of producing at least 30 g/L pantothenate in a manner independent of aspartate or β-alanine feed comprising culturing an aspartate-α-decarboxylase-overexpressing (AαD-O) microorganism under conditions such that
 30 pantothenate is produced.
 - 18. A method of producing at least 30 g/L pantothenate in a manner independent of valine or α -ketoisovalerate feed comprising culturing a microorganism having a deregulated isoleucine-valine (ilv) biosynthetic pathway under conditions such that pantothenate is produced.

- 105 -

- 19. A β -alanine independent high yield production method for producing pantothenate comprising culturing a manipulated microorganism under conditions such that pantothenate is produced at a significantly high yield.
- 5 20. The method of any one of claims 14 to 19, wherein the microorganism overexpresses acetohydroxyacid synthetase or is transformed with a vector comprising an *ilvBN* nucleic acid sequence or an *alsS* sequence.
- 21. The method of any one of claims 14 to 19, wherein the microorganism overexpresses acetohydroxyacid isomeroreductase or is transformed with a vector comprising an *ilvC* nucleic acid sequence.
- 22. The method of any one of claims 14 to 19, wherein the microorganism overexpresses dihydroxyacid dehydratase or is transformed with a vector comprising an *ilvD* nucleic acid sequence.
 - 23. The method of any one of claims 19 to 22, wherein the microorganism overexpresses aspartate- α -decarboxylase or is transformed with a vector comprising a *panD* nucleic acid sequence.
 - 24. The method of any one of claims 14 to 23, wherein the microorganism further has a deregulated pantothenate biosynthetic pathway.
- 25. The method of any one of claims 14 to 24, wherein the microorganism further has at least one mutant gene selected from the group consisting of a mutant *avtA* gene, a mutant *ilvE* gene, a mutant *ansB* gene and a mutant *alsD* gene.
- The method of claim 24, wherein the microorganism overexpresses any of ketopantoate hydroxymethyltransferase, ketopantoate reductase,
 pantothenate synthetase and aspartate-α-decarboxylase.
 - 27. The method of claim 24 or 26, wherein the microorganism is transformed with a vector comprising a *panBCD* nucleic acid sequence or a vector comprising a *panE1* nucleic acid sequence.

- 106 -

- 28. The method of any one of claims 14 to 16 and 19 to 27, wherein pantothenate is produced at a level selected from the group consisting of a level greater than 10g/L, a level greater than 20g/L and a level greater than 40g/L.
- 5 29. The method of claim 20, wherein the microorganism overexpresses acetohydroxyacid synthetase derived from *Bacillus* or is transformed with a vector comprising an *ilvBN* nucleic acid sequence or an *alsS* nucleic acid sequence derived from *Bacillus*.
- 30. The method of claim 21, wherein the microorganism overexpresses acetohydroxyacid isomeroreductase derived from *Bacillus* or is transformed with a vector comprising an *ilvC* nucleic acid sequence derived from *Bacillus*.
- 15 31. The method of claim 22, wherein the microorganism overexpresses dihydroxyacid dehydratase derived from *Bacillus* or is transformed with a vector comprising av *ilvD* nucleic acid sequence derived from *Bacillus*.
- 32. The method of claim 23, wherein the microorganism
 20 overexpresses aspartate-α-decarboxylase derived from *Bacillus* or is transformed with a vector comprising a *panD* nucleic acid sequence derived from *Bacillus*.
- The method of claim 24 or 26, wherein the microorganism overexpresses any of ketopantoate hydroxymethyltransferase, ketopantoate reductase,
 pantothenate synthetase and aspartate-α-decarboxylase derived from *Bacillus*.
 - 34. The method of claim 27, wherein the vector comprises a *panBCD* nucleic acid sequence or a *panE1* nucleic acid sequence derived from *Bacillus*.
- 35. A method of producing a panto-compound comprising contacting a composition comprising at least one pantothenate biosynthesis pathway precursor or isoleucine-valine biosynthesis pathway precursor with at least one isolated *Bacillus* enzyme selected from the group consisting of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase and aspartate-α-decarboxylase, under conditions such that the panto-compound is produced.

- 36. A method of producing β -alanine comprising culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism under conditions such that β -alanine is produced.
- 5 37. The method of claim 36, wherein the $A\alpha D$ -O microorganism has a mutation in a nucleic acid sequence encoding a pantothenate biosynthetic enzyme selected from the group consisting of ketopantoate hydroxymethyltransferase, ketopantoate reductase and pantothenate synthetase.
- 38. A method of producing β -alanine comprising contacting a composition comprising aspartate with an isolated *Bacillus* aspartate- α -decarboxylase enzyme under conditions such that β -alanine is produced.
- 39. A method for enhancing production of a panto-compound comprising culturing a mutant microorganism having a mutant *coaX* gene under conditions such that the panto-compound production is enhanced.
 - 40. The method of claim 39, wherein said recombinant microorganism has a mutant *coaA* gene.
 - 41. A method of producing a panto-compound comprising a pantothenate kinase mutant microorganism under conditions such that the panto-compound is produced at a significantly high yield.
- 25 42. The method of claim 41, wherein said mutant microorganism has a mutant *coaA* gene.
 - 43. The method of claim 41, wherein said mutant microorganism has a mutant *coaX* gene.
 - 44. The method of claim 41, where said mutant microorganism has a mutant *coaA* and *coaX* gene.
- 45. The method of any one of claims 39 to 44, wherein said panto-35 compound is selected from the group consisting of ketopantoate, pantoate or pantothenate.

- 46. The method of any one of claims 39 to 44, wherein said panto-compound is pantothenate.
- 47. The method of any one of claims 39 to 44, wherein said panto-5 compound is produced at a level selected from the group consisting of a level greater than 10g/L, a level greater than 20g/L and a level greater than 40g/L.
- 48. The method of any one of claims 39 to 44, wherein said recombinant microorganism further has a deregulated pantothenate biosynthetic pathway or further has a deregulated isoleucine-valine (*ilv*) biosynthetic pathway.
 - 49. The method of claim any one of claims 39 to 44, wherein said recombinant microorganism further overexpresses *panD* and *panE*.
- 15 50. The method of any one of claims 39 to 44, wherein said recombinant microorganism further has at least one mutant gene selected from the group consisting of a mutant *avtA* gene, a mutant *ilvE* gene, a mutant *ansB* gene and a mutant *alsD* gene.
- 51. A method for enhancing production of a panto-compound comprising culturing a microorganism that has a deregulated pantothenate biosynthetic pathway and that also has a mutation that results in reduced pantothenate kinase activity under conditions such that the panto-compound production is enhanced.
- 25 52. A method for identifying compounds which modulate pantothenate kinase activity comprising contacting a recombinant cell expressing pantothenate kinase encoded by the *coaX* gene with a test compound and determining the ability of the test compound to modulate pantothenate kinase activity in said cell.
- 30 53. The method of claim 52, wherein said cell further comprises a mutant *coaA* gene encoding a pantothenate kinase having reduced activity.
 - 54. The method of any one of claims 1 to 51, wherein the microorganism is Gram positive.
 - 55. The method of any one of claims 1 to 51, wherein the microorganism is Gram negative.

20

- 56. The method of any one of claims 1 to 51, wherein the microorganism is a microorganism belonging to a genus selected from the group consisting of *Bacillus*, *Cornyebacterium*, *Lactobacillus*, *Lactococci* and *Streptomyces*.
- 57. The method of any one of claims 1 to 51 and 54 to 56, wherein the microorganism is of the genus *Bacillus*.
- 58. The method of any one of claims 1 to 51 and 54 to 57, wherein the microorganism is *Bacillus subtilis*.
 - 59. The method of any one of claims 1 to 13, 35, 39 to 51 and 54 to 58, further comprising recovering the panto-compound.
- 15 60. The method of any one of claims 14 to 34 and 54 to 58, further comprising recovering the pantothenate.
 - 61. The method of any one of claims 1 to 14, 35, 39 to 46, 48 to 51 and 54 to 59, wherein the panto-compound is produced at a level greater than $2~\rm g/L$.
 - 62. A recombinant microorganism which overexpresses at least one *Bacillus* pantothenate biosynthetic enzyme.
- 63. The recombinant microorganism of claim 62, which overexpresses at least one *Bacillus subtilis* pantothenate biosynthetic enzyme.
 - 64. The recombinant microorganism of claim 62 or 63, wherein the pantothenate biosynthetic enzyme is selected from the group consisting of ketopantoate hydroxymethyltransferase, pantothenate synthetase, aspartate-α-decarboxylase and ketopantoate reductase.
 - 65. The recombinant microorganism of any one of claims 62 to 64, wherein the pantothenate biosynthetic enzyme is ketopantoate reductase.
- 35 66. A recombinant microorganism which overexpresses aspartate-α-decarboxylase and has a deregulated isoleucine-valine (*ilv*) biosynthetic pathway.

20

- 67. A recombinant microorganism having a mutant *coaX* gene, said mutant *coaX* gene encoding reduced pantothenate kinase activity in said microorganism.
- 68. The recombinant microorganism of claim 67 further having a mutant *coaA* gene, said mutant *coaA* gene encoding reduced pantothenate kinase activity in said microorganism.
- 69. A recombinant microorganism having a mutant *coaX* gene and optionally having a mutant *coaA* gene, said mutant microorganism having reduced pantothenate kinase activity as compared to a microorganism having wild-type *coaA* and *coaX* genes.
 - 70. A recombinant microorganism comprising a vector comprising an isolated *coaX* gene.
 - 71. A recombinant microorganism that overproduces a panto-compound, the microorganism having a deregulated pantothenate biosynthetic pathway and having at least one mutation that results in a decrease in the capacity of the microorganism to synthesize Coenzyme A (CoA).
 - 72. The recombinant microorganism of claim 71, having at least one mutation that results in a reduced level of pantothenate kinase activity.
- 73. The recombinant microorganism of claim 72, having a mutation in a *coaA* gene, or homologue thereof, that results in a reduced level of CoaA enzyme activity.
- 74. The recombinant microorganism of claim 72, having a mutation in a *coaX* gene, or homologue thereof, that results in a reduced level of CoaX enzyme 30 activity.
 - 75. The recombinant microorganism of claim 72, having a mutation in a *coaA* gene, or homologue thereof, and having a mutation in a *coaX* gene, or homologue thereof, the mutations resulting in reduced levels of CoaA enzyme activity and reduced CoaX enzyme activity.

- 111 -

- 76. The recombinant microorganism of any one of claims 66 to 70 which further has a deregulated pantothenate biosynthetic pathway.
- 77. The recombinant microorganism of any one of claims 62 to 65 and 67 to 75, further having a deregulated isoleucine-valine (*ilv*) biosynthetic pathway.
 - 78. The recombinant microorganism of any one of claims 62 to 77, which is Gram positive.
- 79. The recombinant microorganism of claim 78 belonging to a genus selected from the group consisting of *Bacillus*, *Cornyebacterium*, *Lactobacillus*, *Lactococci* and *Streptomyces*.
- 80. The recombinant microorganism of claim 79 belonging to the genus *Bacillus*.
 - 81. The recombinant microorganism of claim 80 which is *Bacillus* subtilis.
- 20 82. A recombinant microorganism selected from the group consisting of PA221, PA235, PA236, PA313, PA410, PA402, PA403, PA411, PA412, PA413, PA303, PA327, PA328, PA401, PA340, PA342, PA404, PA405, PA374, PA354, PA365, PA377, PA651 and PA824.
- 25 83. A recombinant vector for use in the production of pantocompounds comprising a nucleic acid sequence which encodes at least one *Bacillus* pantothenate biosynthetic enzyme operably linked to regulatory sequences.
- 84. The vector of claim 83, comprising a nucleic acid sequence which 30 encodes at least one *Bacillus subtilis* pantothenate biosynthetic enzyme.
 - 85. The vector of claim 84, wherein the nucleic acid sequence encodes at least one of ketopantoate hydroxymethyltransferase, pantothenate synthetase, aspartate- α -decarboxylase and ketopantoate reductase.

- 112 -

- 86. A recombinant vector comprising at least one nucleic acid sequence selected from the group consisting of SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29 and SEQ ID NO:59.
- 5 87. The vector of claim 84, wherein the nucleic acid sequence encodes ketopantoate reductase.
 - 88. A vector comprising a mutant *coaX* gene, said mutant encoding a pantothenate kinase enzyme having reduced activity.
 - 89. A vector comprising an isolated *coaX* gene.
 - 90. A vector comprising an isolated *Bacillus coaX* gene.
- 15 91. A vector comprising an isolated *Bacillus subtilis coaX* gene.
 - 92. The vector of any one of claims 86 and 89 to 91, which further comprises regulatory sequences.
- 20 93. The vector of any one of claims 83 to 85, 87 and 92, wherein the regulatory sequences comprise a constitutively active promoter.
- 94. The vector of claim 93, wherein the constitutively active promoter comprises P_{veg} (SEQ ID NO:41), P_{15} (SEQ ID NO:39) or P_{26} (SEQ ID NO:40) sequences.
 - 95. The vector of claim 83, wherein the regulatory sequences comprise at least one artificial ribosome binding site (RBS).
- 30 96. The vector of claim 95, wherein the artificial RBS comprises a nucleotide sequence selected from the group consisting of SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56 and SEQ ID NO:57.
- 35 97. A vector selected from the group consisting of pAN004, pAN005, pAN006, pAN236, pAN423, pAN428, pAN429, pAN441, pAN442, pAN443, pAN251,

25

30

35

pAN267, pAN256, pAN257, pAN263, pAN240, pAN294, pAN296, pAN336, pAN341 and pAN342.

- 98. A recombinant microorganism comprising the vector of claim 86 5 or 93.
 - 99. An isolated nucleic acid molecule which encodes at least one *Bacillus* pantothenate biosynthetic gene.
- 100. The isolated nucleic acid molecule of claim 99 which encodes at least one *Bacillus subtilis* pantothenate biosynthetic gene.
 - 101. The isolated nucleic acid molecule of claim 99 or 100 which encodes ketopantoate reductase.
 - 102. An isolated *Bacillus* pantothenate biosynthetic enzyme polypeptide.
- 103. An isolated *Bacillus subtilis* pantothenate biosynthetic enzyme 20 polypeptide.
 - 104. An isolated *Bacillus* ketopantoate reductase polypeptide.
 - 105. An isolated Bacillus subtilis ketopantoate reductase polypeptide.
 - 106. An isolated *Bacillus* aspartate-α-decarboxylase polypeptide.
 - 107. An isolated *Bacillus subtilis* aspartate- α -decarboxylase polypeptide.
 - 108. An isolated nucleic acid molecule comprising a mutant *coaX* gene.
 - 109. An isolated nucleic acid molecule comprising a *coaX* gene.
 - 110. An isolated pantothenate kinase protein encoded by a *coaX* gene.

- 114 -

METHODS AND MICROORGANISMS FOR PRODUCTION OF PANTO-COMPOUNDS

Abstract of the Disclosure

The present invention features methods of producing panto-compounds (e.g., pantothenate) using microorganisms in which the pantothenate biosynthetic pathway and/or the isoleucine-valine biosynthetic pathway and/or the coenzymeA biosynthetic pathway has been manipulated. Methods featuring ketopantoate reductase overexpressing microorganisms as well as aspartate α-decarboxylase overexpressing microorganisms are provided. Methods of producing panto-compounds in a precursor-independent manner and in high yield are described. Recombinant microorganisms, vectors, isolated nucleic acid molecules, genes and gene products useful in practicing the above methodologies are also provided. The present invention also features a previously unidentified microbial pantothenate kinase gene, coaX, as well as methods of producing panto-compounds utilizing microorganisms having modified pantothenate kinase activity. Recombinant microorganisms, vectors, isolated coaX nucleic acid molecules and purified CoaX proteins are featured. Also featured are methods for identifying pantothenate kinase modulators utilizing the recombinant microorganisms and/or purified CoaX proteins of the present invention.

20

15

5

FIG 1






Figure 2. Plasmid pAN240, containing sequences ligated upstream of the P₂₆panBCD cassette, equivalent to the integrated version in strain PA221.










Figure 4 Plasmid pAN236, containing an integratable and amplifiable P26-RBS2-panE1 expression cassette.






pAN426

[3588 bp]

AmpR

Repressor

f1-IG



BamHI


pAN427


AmpR


Tet

Repressor

[3588 bp] f1-IG










Figure || Structure of pAN251, a plasmid designed to integrate a single copy of P₂₆ panE1 at the panE1 locus by double crossover.




Figure 12 Structure of pAN267, a plasmid designed to stably integrate a P₂₆ ilvBNC cassette at the amyE locus.







Figure 13 Structure of pAN257, a clone of B. subtilis ilvD in a low copy vector.










Figure 17 Structure of pAN296, designed to delete most of the B. subtilis coaA gene and substitute a chloramphenicol resistance gene.




Figure (8 Structure of the B. subtilis chromosome in the region of the coad gene. The scale is in base pairs and the significant open reading frames are shown by the open arrows.




Figure 19 Structure of pAN281, a plasmid for expressing B. subtilis coaA after integration at the bpr locus. pAN282 and pAN283 have similar structures.




FIG.20A

CLUSTAL W (1.7) Multiple Sequence Alignments

	312 aa	329 aa	312 aa	319 aa	311 aa	316 aa
Sequence type explicitly set to Protein Sequence format is Pearson	1: sp 09X795 M.leprae	spl086779 S.coelicolor				
tyk fol	·-	0		\ <u>4</u>	5	9 6
Sequence	Segmence	Segmence	Segmence	Sections	Seguence	Sequence

sp Q9X795 M.leprae sp 053440 M.tuberculosis	spl0867791S.coelicolor	sp P44793 H.influenzae	sp P15044 E.coli	sp P54556 B.subtilis
sp Q9X795 N sp O53440 N	s 622980 ds	sp P44793 I	sp P15044 E	sp P54556 I

sp|Q9X795|M.leprae sp|O53440|M.tuberculosis sp|O86779|S.coelicolor sp|P44793|H.influenzae sp|P15044|E.coli sp|P54556|B.subtilis sp|Q9X795|M.leprae sp|O53440|M.tuberculosis sp|O86779|S.coelicolor sp|P44793|H.influenzae sp|P15044|E.coli


 GLGEQIDLLEVEEVYLPLARLIHLQVAARQRLFAATAEFLGEPQQNPGRP
GLGEQIDLLEVEEVYLPLARLIHLQVAARQRLFAATAEFLGEPQQNPDRP
GLGDVIDLDEVRDIYLPLSRLINLYVGATDGLRGALNTFLGE--QGSQSG
GFNEDLSLDEVSTIYLPLTRLINYYIDENLHRQTVLHRFLGR----NNAK
GINEDLSLEEVAEIYLPLSRLINFYISSNIRRQAVLEQFLGT----NGQR
GLNDYLSVEEVETIYIPLVRLHHLHVKSAAERNKHVNVFLKHP---HSAK
*::: :: ** ::** **:: ::

FIG.20B

sp|Q9X795|M.leprae sp|O53440|M.tuberculosis sp|O86779|S.coelicolor sp|P44793|H.influenzae sp|P15044|E.coli

sp|Q9X795|M.leprae sp|O53440|M.tuberculosis sp|O86779|S.coelicolor sp|P44793|H.influenzae sp|P15044|E.coli sp|P54556|B.subtilis sp|Q9X795|M.leprae sp|O53440|M.tuberculosis sp|O86779|S.coelicolor sp|P44793|H.influenzae sp|P15044|E.coli sp|Q9X795|M.leprae sp|O53440|M.tuberculosis sp|O86779|S.coelicolor sp|P44793|H.influenzae sp|P15044|E.coli sp|P54556|B.subtilis

GRRNLMHRKGFPESYNRRALMRFVTSVKSGADYACAPVYSHLRYDTIPGA QRRNLMHRKGFPESYNRRALMRFVTSVKSGSDYACAPVYSHLHYDIIPGA EARGLMSRKGFPESYDRRALTRFVADIKAGKAEVTAPVYSHLIYDIVPDQ KQDNLLQKKGFPVSYDTPKLIRFLADVKSGKSNVTAPIYSHLTYDIIPDK KERGLMKKKGFPESYDMHRLVKFVSDLKSGVPNVTAPVYSHLIYDVIPDG KKKNMMSRKGFPESYDVKALLEFLNDLKSGKDSVKAPVYSHLIYDNFEGV KKKNMMSRKGFPESYDVKALLEFLNDLKSGKDSVKAPVYSHLIYDREEGV






Figure 22 Structure of pAN341 and pAN342, two independent PCR- derived clones of yacB (renamed coaX).




FIG.23A

CLUSTAL W (1.7) Multiple Sequence Alignments

aa Seq.8: sp O51477 B.burgdorferi 262 aa seq.9: sp P74045 Synechocystis 257 aa aa Seq.10: sp O25533 H.pylori 223 aa seq.11: sp O67753 A.aeolicus 229 aa seq.12: sp Q9RX54 D.radiodurans aa Seq.13: WIT RCA03301 C.acetobutylicum 250 aa aa Seq.13: WIT RRC02473 R.capsulatus 258 aa	NKRAAFMLLLFLRSVLKVILVLDVGNTNIVLGIYNDGKLEYHWRIE
Sequence type explicitly set to Protein Sequence format is Pearson Seq. 1: B.subtilis Coax SEQNO_9 258 aa Seq.2: dbj BAA21476.1 D.vulgaris 212 aa Seq.3: gb AAD35964.1 T.maritima 246 aa Seq.4: pir T36391 S.coelicolor 265 aa Seq.5: sp Q4538 B.pertussis 267 aa Seq.6: sp O06282 M.tuberculosis 272 aa Seq.7: sp O83446 T.pallidum 273 aa	B.subtilis Coax SEQIDNO_9 WIT RCA03301 C.acetobutylicum pir T36391 S.coelicolor sp O06282 M.tuberculosis WIT RRC02473 R.capsulatus dbj BAA21476.1 D.vulgaris sp Q9RX54 D.radiodurans gb AAD35964.1 T.maritima sp O83446 T.pallidum sp O67753 A.aeolicus sp O67753 A.aeolicus sp P74045 Synechocystis sp Q45338 B.pertussis

FIG.23B

B.subtilis|Coax|SEQIDNO_9
WIT|RCA03301|C.acetobutylicum
pir|T36391|S.coelicolor
sp|O06282|M.tuberculosis
WIT|RRC02473|R.capsulatus
dbj|BAA21476.1|D.vulgaris
sp|Q9RX54|D.radiodurans
gp|Q9RX54|D.radiodurans
sp|Q9RX54|D.radiodurans
sp|Q9RX54|D.radiodurans
sp|Q9RX54|D.radiodurans
sp|Q9RX54|D.radiodurans
sp|Q9RX54|D.radiodurans
sp|Q83446|T.pallidum
sp|O51477|B.burgdorferi
sp|O67753|A.aeolicus
sp|P74045|Synechocystis
sp|Q45338|B.pertussis

B.subtilis|Coax|SEQIDNO_9
WIT|RCA03301|C.acetobutylicum
pir|T36391|S.coelicolor
sp|006282|M.tuberculosis
WIT|RRC02473|R.capsulatus
dbj|BAA21476.1|D.vulgaris
sp|Q9RX54|D.radiodurans
gp|Q9RX54|D.radiodurans
sp|Q9RX54|D.radiodurans
sp|Q9RX54|D.radiodurans
sp|Q9RX54|D.radiodurans
sp|Q9RX54|D.radiodurans
sp|Q9RX54|D.radiodurans
sp|Q9RX54|D.radiodurans
sp|Q9RX54|D.radiodurans
sp|Q9RX54|B.burgdorferi
sp|Q7753|A.aeolicus
sp|Q7753|A.aeolicus
sp|Q7753|H.pylori
sp|Q25533|H.pylori

-----ALGISVKQSFSEKVRG -----VLLGKVPLMLASVVPE .-----LGIQKEIFYISVNEE AFDNIDLDALGRWLATLPRRP----Q----RALGVNVAGLARGEAIA TDSRRTADELAVLLQGLMGMHPLLGDELGDGIDGIAICATVPSVLHELRE TDHRRTADEYFVWLNTLMQLK----GLQGRISEAIISSTAPRVVFNLRV TDPGQTTDSIGLRLLEVLRHAG----LGPADVGACVASSVVPGVNPLIRR TNREMLPDDLALQLHGLFTLA----GAP-IPRAAVLSSVAPPVGENYAL TGVFQTEDELFSHLHPLLG-----DAMREIKGIGVASVVPTQNTVIER PDARKTQDEYSLLIHALCERAG----VGRASLRDAFISSVVPVLTKTIAD TNIMIRYDEVYSFFEENFDFN----VN---K-VFISSVVPILNETFKN TDVLRSADEYGIQVMNLFQQD----KLDPTLVEGVIISSVVPNIMYSLEH --GLMFEQIDGIIISSVVPPIMFALER TESEVTADELALTIDGLIG-----EDSERLIGTAALSTVPSVLHEVRI DFLKLSHEEFLKEEFPKLK----SGNAPLQTWVTDYNPKSAQLP--LHFAQNYQLFSSAKEDLKR----TSRHKTEDEFGMILRSLFDHS--

MCTKYFHIEPQIVG-PG-MKTGLNIKYDNPKEVGADRIVNAVAAIHLYGMIRKYFKINPLVVG-PG-IKTGINIKYDNPKEVGADRIVNAVAAHEIYKVTRRYYGDVPAVLVEPG-VKTGVPILTDHPKEVGADRIINAVAAVELYGMLDQYWPSVPHVLIEPG-VRTGIPLLVDNPKEVGADRIINAVAAVELYGLCNRYFDCRPYVVGKPG-CELPVAPRVDPGTTVGPDRLVNTVAGYDRFRLCNRYFDCRPYVVGKPG-CELPVAPRVDPGTTVGPDRLVNTVAGYDRFRACERYL--YRKLLFAPGDIAIPLDNRYERPAEVGADRLVAAYAARRLYPALKRHFMIDAFAVSAEN--LPDVTVELDTPGSVGADRLVAAYAARRLYPAVAQISGVQPVVFGPWAYEHLPVRIPEPVRAEIGTDLVANAVAAYVHFRVIFSFFKIKPLFIGFDLNYDLTFNPYKSDKFLLGSDVFANLVAAIENYSKIPKIK----FLKKEN---FPIQVDYKTPETLGTDRVALAYSAKKFYGQTEVWRVYQPKILTLKN---LPLVNLYP---SFGIDRALAGLGTGLTYGNEKALLNCYPNAKNIAG--FFHLETDYVG---LGIDRQMACLA--VN-ATLRAGGCDIRWLRAQP-LAMGLRNGYRNPDQLGADRWACWVGVLARQPS

*.

FIG.23C

B.subtilis|Coax|SEQIDNO_9
WIT|RCA03301|C.acetobutylicum
pir|T36391|S.coelicolor
sp|O06282|M.tuberculosis
WIT|RRC02473|R.capsulatus
dbj|BAA21476.1|D.vulgaris
sp|Q9RX54|D.radiodurans
gp|AAD35964.1|T.maritima
sp|O8746|T.pallidum
sp|O57477|B.burgdorferi
sp|O67753|A.aeolicus
sp|P74045|Synechocystis
sp|Q25533|H.pylori
sp|Q45338|B.pertussis

B.subtilis|Coax|SEQIDNO_9
WIT|RCA03301|C.acetobutylicum
pir|T36391|S.coelicolor
sp|O06282|M.tuberculosis
WIT|RRC02473|R.capsulatus
dbj|BAA21476.1|D.vulgaris
sp|Q9RX54|D.radiodurans
gp|Q9RX54|D.radiodurans
sp|Q9RX54|D.radiodurans
sp|Q9RX54|F.maritima
sp|O51477|B.burgdorferi
sp|O67753|A.aeolicus
sp|O7753|A.aeolicus
sp|O7753|H.pylori
sp|Q45338|B.pertussis

NP--LIVVDFGTATTYCYIDENKQYMGGAIAPGITISTEALYSRAAKLPR
RS--LIIIDFGTATTFCAVRENGDYLGGAICPGIKVSSEALFEKAAKLPR
GP--AIVVDFGTATTFDAVSARGEYIGGVIAPGIEISVEALGVKGAQLRK
KA--AIVVDFGSSICVDVVSAKGEFLGGAIAPGVQVSSDAAAARSAALRR
GD--LIVVDFGTATTFDVVAPDGAYIGGVIAPGVNLSLEALHMAAAALPH
GPRSLVSVDFGTATTFDVVAPDGAYIGGVIAPGVNLSLEALHMAAAALPH
GPRSLVSVDFGTATTFDCVEG-GAYLGGLICPGVLSSAGALSSRTAKLPR
GLDYAVVVDFGTSTNFDVVGRGRRFLGGILATGAQVSADALFARAAKLPR
KN--GIIIDMGTATTVDLVVN-GSYEGGAILPGFFFMMVHSLFRGTAKLPL
SA--CVVVDCGTALTFTAVDGTGLIQGVAIAPGLRTAVQSLHTGTAQLPL
FEN-VLVVDLGTACTIFAVSRQDGILGGIINSGPLINFNSLLDNAYLIKK
KN--VVVISAGTALVIDLVLE-GKFKGGFITLEGLGKKIKILSDLAEGIPE
FP--CLVVDGGTALTITGFDQDKKLVGGAILPGLGLATLGDRLAALPR
NG---VVVDAGSAITIDLIKE-GKHLGGCILPGPAMMRGALAYGTAHLPL

.. ∗

Here the the term of the term that the term to the term to the term of the term to the ter

FIG.23D

VIATGG	LAPLIANESDCIDIVDPFLTLKGLELI
RTSLVLATGG	
DDVTVIATGG	LAPMVLGESSVIDEHEPWLTLMGLRLV
HDVAIVATGH	TAPLLLPELHTVDHYDQHLTLQGLRLV
MKVIATGG	LASLFDLGFDLFDKVEDDLTMHGLRLI
1111111111	
AVAVATGG	FSRTVQGICQEIDYYDETLTLRGLVEL
NVLTGG	QSKIVK-DMIKHEIFDEDLTIKGVYHF
CAAVITGG	LSRLFS-SEVDFPPIDAQLTLSGLAHI
FNLIITGG	NADLILSLIEIEFIFNIHLTVEGVRIL
FKVVITGG	EGKYFSKFGIYDPLLVHRGMRNL
AMVITGG	DGKILHGFLKEHSPNLSVAWDDNLIFLGMAAI
IATCGG	DAKYLSAFLPHSVCKERLVFDGMEIA
EIYVAGGGWP	EIYVAGGGWPEVRQEAERLLAVTGAAFGATPQPTYLDSPVLDGLAAL

Figure 24 Alignment of a portion of the amino acid sequences of several known or suspected pantothenate kinases. The residues that are mutated in E. coli coa A15(Ts) and B. subtilis coa A from plasmid pAN282A are indicated below and above the alignment, respectively. The coordinate given in the left margin for the B. subtilis protein refers to the coa A1 open reading frame.

Majority	B. subtilis CoaA1 E. coli CoaA H. influenzae CoaA M. leprae CoaA M. tuberculosis CoaA S. coelecolor CoaA
TAPVYSHLIYDIIPGA	H C T Y D R E E G V C T Y D I I P D G C R Y D T I P G A C H Y D I I P G A C I Y D I I P G A
O	000000
ما	E
-	
-	x > - F
a	
>	イベイベイエ
-	FHRH
7	
王	HHHHH
S	~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~
>	×××××
>	>> _ > > >
۵۰	V T A P V V T A P V A C A P V V T A P V V T A P V V T A P V V T A P V V T A P V V T A P V V T A P V V A P V V T A P V V A P V
4	44444
⊢	X F F O O F
>	>>> <
Z	N Z Z > > H
K D N	K X X X X X X X X X X X X X X X X X X X
×	X > X A N X
	168 167 165 169 179

Figure 25 Structure of pAN294, a plasmid for integrating mutagenized B. subtilis coaA at its native locus.





Figure 26 Structure of pAN336, a plasmid designed to delete B. subtilis coaX from the chromosome and replace it with a kanamycin resistance gene.



DECLARATION, PETITION AND POWER OF ATTORNEY FOR CONTINUATION-IN-PART PATENT APPLICATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

METHODS AND MICROORGANISMS FOR PRODUCTION OF PANTO-COMPOUNDS

the specification of which

×	executed by me of even date herewith and about to be filed;				
	was filed on as Application Serial No; and				
	was amended on (if applicable)				
	that I have reviewed and understand the contents of the above-identified including the claims, as amended by any amendment referred to above.				
This application application (s),	on in part discloses and claims subject matter disclosed in my earlier filed as follows:				
E	Serial No. 09/400,494 filed September 21, 1999; as to which I claim priority benefit under Title 35, United States Code, §120.				
×	Serial No. 60/210,072 filed June 7, 2000; as to which I claim priority benefit under Title 35, United States Code, §119(e).				
X	Serial No. 60/221,836 filed July 28, 2000 ; as to which I claim priority benefit under Title 35, United States Code, §119(e).				
×	Serial No. 60/227,860 filed August 24, 2000; as to which I claim priority benefit under Title 35, United States Code, §119(e).				

I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56, including all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application(s) and the national or PCT international filing date of the continuation-in-part application.

 \times

AS TO PARENT APPLICATION

As to the subject matter of this application which is common to said earlier application, I do not know and do not believe that the same was ever known or used in the United States of America before my or our invention thereof or patented or described in any printed publication in any country before my or our invention thereof, or more than one year prior to said earlier application, or in public use or on sale in the United States of America more than one year prior to said earlier application; that the common subject matter has not been patented or made the subject of an inventor's certificate issued before the date of said earlier application in any country foreign to the United States of America on an application filed by me or my legal representatives or assigns more than twelve months prior to said earlier application; and

As to applications for patents or inventor's certificate or PCT international application(s) designating at least one country other than the United States of America, on the common subject matter, filed in or designating any country foreign to the United States of America, prior to said earlier application by me or my legal representatives or assigns,

no such applications have been filed.

	REIGN APPLICATION(S), FOR DESIGN) PRIOR TO Application Number	SAID EARLIER U.S. Date of Filing	APPLICATION Priority Claimed
	 	(month,day,year)	Under 35 USC 119 Per No
			☐ Yes ☐ No
			☐ Yes ☐ No
			☐ Yes ☐ No
			☐ Yes ☐ No
	IGN APPLICATION(S), II S FOR DESIGN) PRIOR T		

AS TO THIS APPLICATION

As to the subject matter of this application which is common to said earlier application, I do not know and do not believe that the same was ever known or used in the United States of America before my or our invention thereof or patented or described in any printed publication in any country before my or our invention thereof, or more than one year prior to said earlier application, or in public use or on sale in the United States of America more than one year prior to said earlier application; that the common subject matter has not been patented or made the subject of an inventor's certificate issued before the date of said earlier application in any country foreign to the United States of America on an application filed by me or my legal representatives or assigns more than twelve months prior to said earlier application; and

As to applications for patents or inventor's certificate or PCT international application(s) designating at least one country other than the United States of America, on the common subject matter, filed in or designating any country foreign to the United States of America, prior to said earlier application by me or my legal representatives or assigns,

X	no such applications have been filed.
	such applications have been filed as follows

EARLIEST FOREIGN APPLICATION(S), IF ANY, FILED WITHIN 12 MONTHS (6 MONTHS FOR DESIGN) PRIOR TO SAID EARLIER U.S. APPLICATION

Country	Application Number	Date of Filing	Priority Claimed	
		(month,day,year)	Under 35 USC 119	
			☐ Yes	□ No
			☐ Yes	□ No
			☐ Yes	□ No
			☐ Yes	□ No
			□ Yes	□ No

ALL FOREIGN APPLICATION(S), IF ANY FILED MORE THAN 12 MONTH (6 MONTHS FOR DESIGN) PRIOR TO THIS U.S. APPLICATION		
İ		

CLAIM FOR BENEFIT OF U.S. PROVISIONAL APPLICATION(S)

I hereby claim the benefit under 35 U.S.C. §119(e) of any United States provisional application(s) listed below.

60/210,072	June 7, 2000	
(Application Serial No.)	(Filing Date)	•
60/221 926	July 28, 2000	
60/221,836	July 28, 2000	•
(Application Serial No.)	(Filing Date)	
60/227,860	August 24, 2000	
(Application Serial No.)	(Filing Date)	•
CLAIM FOR BENI	EFIT OF U.S. PATENT APPLIC	CATION(S)
I hereby claim the benefit under 35 listed below.	U.S.C. §120 of any United States	patent application(s)
09/400,494	September 21, 1999	
(Application Serial No.)	(Filing Date)	-
(Application Serial No.)	(Filing Date)	

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorneys and/or agents to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

James E. Cockfield	Reg. No. 19,162	Nicholas P. Triano III	Reg. No. 36,397
Thomas V. Smurzynski	Reg. No. 24,798	Peter C. Lauro	Reg. No. 32,360
Ralph A. Loren	Reg. No. 29,325	DeAnn F. Smith	Reg. No. 36,683
Giulio A. DeConti, Jr.	Reg. No. 31,503	William D. DeVaul	Reg. No. 42,483
Ann Lamport Hammitte	Reg. No. 34,858	David J. Rikkers	Reg. No. 43,882
Elizabeth A. Hanley	Reg. No. 33,505	Chi Suk Kim	Reg. No. 42,728
Amy E. Mandragouras	Reg. No. 36,207	Maria Laccotripe Zacharakis	Limited Recognition
Anthony A. Laurentano	Reg. No. 38,220		Under 37 C.F.R. § 10.9(b)
Jane E. Remillard	Reg. No. 38,872	Debra J. Milasincic	Reg. No. 46,931
Jeremiah Lynch	Reg. No. 17,425	David R. Burns	Reg. No. 46,590
Kevin J. Canning	Reg. No. 35,470	Sean D. Detweiler	Reg. No. 42,482
Jeanne M. DiGiorgio	Reg. No. 41,710	Peter S. Stecher	Reg. No. 47,259
Megan E. Williams	Reg. No. 43,270		

all of: Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 United States of America

Send Correspondence to Elizabeth A. Hanley, Esq. at Customer No. 000959 whose address is:

Lahive & Cockfield, LLP 28 State Street Boston, MA 02109

Direct telephone calls to Elizabeth A. Hanley, Esq. at (617) 227-7400.

Wherefore I petition that letters patent be granted to me for the invention or discovery described and claimed in the attached specification and claims, and hereby subscribe my name to said specification and claims and to the foregoing declaration, power of attorney, and this petition.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor R. Rogers Yocum	
Inventor's signature	Date
Residence	
4 Orchard Lane, Lexington, Massachusetts 02420 Citizenship	
United States of America	
Post Office Address (if different)	
Same as above	,

Full name of second inventor, if any	
Thomas A. Patterson	
Inventor's signature	Date
Residence	
89 Church Street, North Attleboro, Massachusetts 02760	
Citizenship	
United States of America	
Post Office Address (if different)	
Same as above	

Full name of third inventor, if any Theron Hermann	
Inventor's signature	Date
Residence 18 Chilhowie Drive, Kinnelon, New Jersey 07405	
Citizenship	
United States of America	
Post Office Address (if different)	

Full name of third inventor, if any	
Janice G. Pero	
Inventor's signature	Date
Residence	
20 Solomon Pierce Road, Lexington, Massachusetts 02420	
Citizenship	
United States of America	
Post Office Address (if different)	
Same as above	

SEQUENCE LISTING

<110> Yocum, R. et al.

<120> METHODS AND MICROORGANISMS FOR PRODUCTION OF PANTO-COMPOUNDS

<130> BGI-141CP

<140>

<141>

<150> USSN 09/400,494

<151> 1999-09-21

<150> USSN 60/210,072

<151> 2000-06-07

<150> USSN 60/221,836

<151> 2000-07-28

<150> USSN 60/221,836

<151> 2000-08-24

<160> 94

<170> PatentIn Ver. 2.0

<210> 1

<211> 311

<212> PRT

<213> Haemophilus influenzae

<400> 1

Met Glu Phe Ser Thr Gln Gln Thr Pro Phe Leu Ser Phe Asn Arg Glu 1 5 10 15

Gln Trp Ala Glu Leu Arg Lys Ser Val Pro Leu Lys Leu Thr Glu Gln 20 25 30

Asp Leu Lys Pro Leu Leu Gly Phe Asn Glu Asp Leu Ser Leu Asp Glu 35 40 45

Val Ser Thr Ile Tyr Leu Pro Leu Thr Arg Leu Ile Asn Tyr Tyr Ile 50 55 60

Asp Glu Asn Leu His Arg Gln Thr Val Leu His Arg Phe Leu Gly Arg 65 70 75 80

Asn Asn Ala Lys Thr Pro Tyr Ile Ile Ser Ile Ala Gly Ser Val Ala $85 \hspace{1.5cm} 90 \hspace{1.5cm} 95$

Trp Pro Thr Glu Arg Lys Val Asp Leu Ile Thr Thr Asp Gly Phe Leu 115 120 125

Tyr Pro Leu Asn Lys Leu Lys Gln Asp Asn Leu Leu Gln Lys Lys Gly 130 140

13

ľЦ

1.1

13

BGI-141CP - 2 -

Phe Pro Val Ser Tyr Asp Thr Pro Lys Leu Ile Arg Phe Leu Ala Asp 145 150 155 160

Val Lys Ser Gly Lys Ser Asn Val Thr Ala Pro Ile Tyr Ser His Leu 165 170 175

Thr Tyr Asp Ile Ile Pro Asp Lys Phe Asp Val Val Asp Lys Pro Asp 180 185 190

Ile Leu Ile Leu Glu Gly Leu Asn Val Leu Gln Thr Gly Asn Asn Lys
195 200 205

Thr Asp Gln Thr Phe Val Ser Asp Phe Val Asp Phe Ser Ile Tyr Val 210 215 220

Asp Ala Glu Glu Lys Leu Leu Lys Glu Trp Tyr Ile Lys Arg Phe Leu 225 230 235 240

Lys Phe Arg Glu Ser Ala Phe Asn Asp Pro Asn Ser Tyr Phe Lys His 245 250 255

Tyr Ala Ser Leu Ser Lys Glu Glu Ala Ile Ala Thr Ala Ser Lys Ile
260 265 270

Trp Asp Glu Ile Asn Gly Leu Asn Leu Asn Gln Asn Ile Leu Pro Thr 275 280 285

Arg Glu Arg Ala Asn Leu Ile Leu Lys Lys Gly His Asn His Gln Val 290 295 300

Glu Leu Ile Lys Leu Arg Lys 305 310

<210> 2

<211> 316

<212> PRT

<213> Escherichia coli

<400> 2

Met Ser Ile Lys Glu Gln Thr Leu Met Thr Pro Tyr Leu Gln Phe Asp 1 5 10 15

Arg Asn Gln Trp Ala Ala Leu Arg Asp Ser Val Pro Met Thr Leu Ser 20 25 30

Glu Asp Glu Ile Ala Arg Leu Lys Gly Ile Asn Glu Asp Leu Ser Leu 35 40 45

Glu Glu Val Ala Glu Ile Tyr Leu Pro Leu Ser Arg Leu Leu Asn Phe 50 55 60

Tyr Ile Ser Ser Asn Leu Arg Arg Gln Ala Val Leu Glu Gln Phe Leu 65 70 75 80

Gly Thr Asn Gly Gln Arg Ile Pro Tyr Ile Ile Ser Ile Ala Gly Ser 85 90 95

Val Ala Val Gly Lys Ser Thr Thr Ala Arg Val Leu Gln Ala Leu Leu 100 105 110

Ser Arg Trp Pro Glu His Arg Arg Val Glu Leu Ile Thr Thr Asp Gly 115 120 125

Phe Leu His Pro Asn Gln Val Leu Lys Glu Arg Gly Leu Met Lys Lys 130 135 140

Lys Gly Phe Pro Glu Ser Tyr Asp Met His Arg Leu Val Lys Phe Val 145 150 155 160

Ser Asp Leu Lys Ser Gly Val Pro Asn Val Thr Ala Pro Val Tyr Ser 165 170 175

His Leu Ile Tyr Asp Val Ile Pro Asp Gly Asp Lys Thr Val Val Gln
180 185 190

Pro Asp Ile Leu Ile Leu Glu Gly Leu Asn Val Leu Gln Ser Gly Met 195 200 205

Asp Tyr Pro His Asp Pro His His Val Phe Val Ser Asp Phe Val Asp 210 215 220

Phe Ser Ile Tyr Val Asp Ala Pro Glu Asp Leu Leu Gln Thr Trp Tyr 225 230 235 240

Ile Asn Arg Phe Leu Lys Phe Arg Glu Gly Ala Phe Thr Asp Pro Asp 245 250 255

Ser Tyr Phe His Asn Tyr Ala Lys Leu Thr Lys Glu Glu Ala Ile Lys 260 265 270

Thr Ala Met Thr Leu Trp Lys Glu Ile Asn Trp Leu Asn Leu Lys Gln 275 280 285

Asn Ile Leu Pro Thr Arg Glu Arg Ala Ser Leu Ile Leu Thr Lys Ser 290 295 300

Ala Asn His Ala Val Glu Glu Val Arg Leu Arg Lys 305 310 315

<210> 3

<211> 319

<212> PRT

<213> Bacillus subtilis

<400> 3

Met Lys Asn Lys Glu Leu Asn Leu His Thr Leu Tyr Thr Gln His Asn 1 5 10 15

Arg Glu Ser Trp Ser Gly Phe Gly Gly His Leu Ser Ile Ala Val Ser 20 25 30

Glu Glu Glu Ala Lys Ala Val Glu Gly Leu Asn Asp Tyr Leu Ser Val 35 40 45

Glu Glu Val Glu Thr Ile Tyr Ile Pro Leu Val Arg Leu Leu His Leu 50 55 60

His Val Lys Ser Ala Ala Glu Arg Asn Lys His Val Asn Val Phe Leu 65 70 75 80

Lys His Pro His Ser Ala Lys Ile Pro Phe Ile Ile Gly Ile Ala Gly
85 90 95

Ser Val Ala Val Gly Lys Ser Thr Thr Ala Arg Ile Leu Gln Lys Leu 100 105 110

Leu Ser Arg Leu Pro Asp Arg Pro Lys Val Ser Leu Ile Thr Thr Asp 115 120 125

Gly Phe Leu Phe Pro Thr Ala Glu Leu Lys Lys Lys Asn Met Met Ser 130 135 140

Arg Lys Gly Phe Pro Glu Ser Tyr Asp Val Lys Ala Leu Leu Glu Phe 145 150 155 160

Leu Asn Asp Leu Lys Ser Gly Lys Asp Ser Val Lys Ala Pro Val Tyr
165 170 175

Ser His Leu Thr Tyr Asp Arg Glu Glu Gly Val Phe Glu Val Val Glu
180 185 190

Gln Ala Asp Ile Val Ile Ile Glu Gly Ile Asn Val Leu Gln Ser Pro \$195\$

Thr Leu Glu Asp Asp Arg Glu Asn Pro Arg Ile Phe Val Ser Asp Phe 210 215 220

Phe Asp Phe Ser Ile Tyr Val Asp Ala Glu Glu Ser Arg Ile Phe Thr 225 230 235 240

Trp Tyr Leu Glu Arg Phe Arg Leu Leu Arg Glu Thr Ala Phe Gln Asn 245 250 255

Pro Asp Ser Tyr Phe His Lys Phe Lys Asp Leu Ser Asp Gln Glu Ala 260 265 270

Asp Glu Met Ala Ala Ser Ile Trp Glu Ser Val Asn Arg Pro Asn Leu 275 280 285

Tyr Glu Asn Ile Leu Pro Thr Lys Phe Arg Ser Asp Leu Ile Leu Arg 290 295 300

Lys Gly Asp Gly His Lys Val Glu Glu Val Leu Val Arg Arg Val 305 310 315

<210> 4

<211> 312

<212> PRT

<213> Mycobacterium leprae

<400> 4

Met Pro Arg Leu Ser Glu Pro Ser Pro Tyr Val Glu Phe Asp Arg Lys 1 5 10 15

Gln Trp Arg Ala Leu Arg Met Ser Thr Pro Leu Ala Leu Thr Glu Glu 20 25 30

Glu Leu Ile Gly Leu Arg Gly Leu Gly Glu Gln Ile Asp Leu Leu Glu
35 40 45

Val Glu Glu Val Tyr Leu Pro Leu Ala Arg Leu Ile His Leu Gln Val 50 55 60

Ala Ala Arg Gln Arg Leu Phe Ala Ala Thr Ala Glu Phe Leu Gly Glu 65 70 75 80

Pro Gln Gln Asn Pro Gly Arg Pro Val Pro Phe Ile Ile Gly Val Ala 85 90 95

Gly Ser Val Ala Val Gly Lys Ser Thr Thr Ala Arg Val Leu Gln Ala 100 105 110

Leu Leu Ala Arg Trp Asp His His Thr Arg Val Asp Leu Val Thr Thr 115 120 125

Asp Gly Phe Leu Tyr Pro Asn Ala Glu Leu Gly Arg Arg Asn Leu Met 130 135 140

His Arg Lys Gly Phe Pro Glu Ser Tyr Asn Arg Arg Ala Leu Met Arg 145 150 155 160

Phe Val Thr Ser Val Lys Ser Gly Ala Asp Tyr Ala Cys Ala Pro Val 165 170 175

Tyr Ser His Leu Arg Tyr Asp Thr Ile Pro Gly Ala Lys His Val Val 180 185 190

Arg His Pro Asp Ile Leu Ile Leu Glu Gly Leu Asn Val Leu Gln Thr 195 200 205

Gly Pro Thr Leu Met Val Ser Asp Leu Phe Asp Phe Ser Leu Tyr Val 210 215 220

Asp Ala Arg Ile Gln Asp Ile Glu Gln Trp Tyr Val Ser Arg Phe Leu 225 230 235 240

Ala Met Arg Gly Thr Ala Phe Ala Asp Pro Glu Ser His Phe His His 245 250 255

Tyr Ser Ala Leu Thr Asp Ser Lys Ala Ile Ile Ala Ala Arg Glu Ile 260 265 270

Trp Arg Ser Ile Asn Arg Pro Asn Leu Val Glu Asn Ile Leu Pro Thr 275 280 285

Arg Pro Arg Ala Thr Leu Val Leu Arg Lys Asp Ala Asp His Ser Ile 290 295 300

Asn Arg Leu Arg Leu Arg Lys Leu 305 310

<210> 5

<211> 312

<212> PRT

<213> Mycobacterium tuberculosis

<400> 5

Met Ser Arg Leu Ser Glu Pro Ser Pro Tyr Val Glu Phe Asp Arg Arg 1 5 10 15

Gln Trp Arg Ala Leu Arg Met Ser Thr Pro Leu Ala Leu Thr Glu Glu 20 25 30

Glu Leu Val Gly Leu Arg Gly Leu Gly Glu Gln Ile Asp Leu Leu Glu 35 40 45

Val Glu Glu Val Tyr Leu Pro Leu Ala Arg Leu Ile His Leu Gln Val 50 60

Ala Ala Arg Gln Arg Leu Phe Ala Ala Thr Ala Glu Phe Leu Gly Glu 65 70 75 80

Pro Gln Gln Asn Pro Asp Arg Pro Val Pro Phe Ile Ile Gly Val Ala 85 90 95

Gly Ser Val Ala Val Gly Lys Ser Thr Thr Ala Arg Val Leu Gln Ala 100 105 110

Leu Leu Ala Arg Trp Asp His His Pro Arg Val Asp Leu Val Thr Thr 115 120 125

Asp Gly Phe Leu Tyr Pro Asn Ala Glu Leu Gln Arg Asn Leu Met 130 135 140

His Arg Lys Gly Phe Pro Glu Ser Tyr Asn Arg Arg Ala Leu Met Arg 145 150 155 160

Phe Val Thr Ser Val Lys Ser Gly Ser Asp Tyr Ala Cys Ala Pro Val 165 170 175

Tyr Ser His Leu His Tyr Asp Ile Ile Pro Gly Ala Glu Gln Val Val 180 185 190

Arg His Pro Asp Ile Leu Ile Leu Glu Gly Leu Asn Val Leu Gln Thr 195 200 205

Gly Pro Thr Leu Met Val Ser Asp Leu Phe Asp Phe Ser Leu Tyr Val 210 215 220

Asp Ala Arg Ile Glu Asp Ile Glu Gln Trp Tyr Val Ser Arg Phe Leu 225 230 235 240

Ala Met Arg Thr Thr Ala Phe Ala Asp Pro Glu Ser His Phe His His 245 250 255

Tyr Ala Ala Phe Ser Asp Ser Gln Ala Val Val Ala Ala Arg Glu Ile 260 265 270

Trp Arg Thr Ile Asn Arg Pro Asn Leu Val Glu Asn Ile Leu Pro Thr 275 280 285

Arg Pro Arg Ala Thr Leu Val Leu Arg Lys Asp Ala Asp His Ser Ile 290 295 300

Asn Arg Leu Arg Leu Arg Lys Leu 305 310

<210> 6

<211> 329

<212> PRT

<213> Streptomyces coelicolor

<400> 6
Met Ile Ser Pro Val Pro Ser Ile Pro Arg Ser Ala His Arg Gln Arg

Pro Glu Ala Thr Pro Tyr Val Asp Leu Thr Arg Pro Glu Trp Ser Ala 20 25 30

Leu Arg Asp Lys Thr Pro Leu Pro Leu Thr Ala Glu Glu Val Glu Lys 35 40 45

Leu Arg Gly Leu Gly Asp Val Ile Asp Leu Asp Glu Val Arg Asp Ile 50 55 60

Tyr Leu Pro Leu Ser Arg Leu Leu Asn Leu Tyr Val Gly Ala Thr Asp
65 70 75 80

Gly Leu Arg Gly Ala Leu Asn Thr Phe Leu Gly Glu Gln Gly Ser Gln
85 90 95

Ser Gly Thr Pro Phe Val Ile Gly Val Ala Gly Ser Val Ala Val Gly 100 105 110

Lys Ser Thr Val Ala Arg Leu Leu Gln Ala Leu Leu Ser Arg Trp Pro 115 120 125

Glu His Pro Arg Val Glu Leu Val Thr Thr Asp Gly Phe Leu Leu Pro 130 135 140

Thr Arg Glu Leu Glu Ala Arg Gly Leu Met Ser Arg Lys Gly Phe Pro 145 150 155 160

Glu Ser Tyr Asp Arg Arg Ala Leu Thr Arg Phe Val Ala Asp Ile Lys 165 170 175

Ala Gly Lys Ala Glu Val Thr Ala Pro Val Tyr Ser His Leu Ile Tyr 180 \$185\$

Asp Ile Val Pro Asp Gln Arg Leu Val Val Arg Arg Pro Asp Ile Leu 195 200 205

Ile Val Glu Gly Leu Asn Val Leu Gln Pro Ala Leu Pro Gly Lys Asp 210 215 220

Gly Arg Thr Arg Val Gly Leu Ala Asp Tyr Phe Asp Phe Ser Val Tyr 225 230 235 240

Val Asp Ala Arg Thr Glu Asp Ile Glu Arg Trp Tyr Leu Asn Arg Phe 245 250 255

Arg Lys Leu Arg Ala Thr Ala Phe Gln Asn Pro Ser Ser Tyr Phe Arg 260 265 270

Lys Tyr Thr Gln Val Ser Glu Glu Glu Ala Leu Asp Tyr Ala Arg Thr 275 280 285

Thr Trp Arg Thr Ile Asn Lys Pro Asn Leu Val Glu Asn Val Ala Pro 290 295 300

Thr Arg Gly Arg Ala Thr Leu Val Leu Arg Lys Gly Pro Asp His Lys

305 310 315

Val Gln Arg Leu Ser Leu Arg Lys Leu 325

<210> 7

<211> 265

<212> PRT

<213> Streptomyces coelicolor

<400> 7

Met Leu Thr Ile Asp Val Gly Asn Thr His Thr Val Leu Gly Leu

1 5 10 15

Phe Asp Gly Glu Asp Ile Val Glu His Trp Arg Ile Ser Thr Asp Ser 20 25 30

Arg Arg Thr Ala Asp Glu Leu Ala Val Leu Leu Gln Gly Leu Met Gly 35 40 45

Met His Pro Leu Leu Gly Asp Glu Leu Gly Asp Gly Ile Asp Gly Ile 50 55 60

Ala Ile Cys Ala Thr Val Pro Ser Val Leu His Glu Leu Arg Glu Val 65 70 75 80

Thr Arg Arg Tyr Tyr Gly Asp Val Pro Ala Val Leu Val Glu Pro Gly 85 90 95

Val Lys Thr Gly Val Pro Ile Leu Thr Asp His Pro Lys Glu Val Gly 100 105 110

Ala Asp Arg Ile Ile Asn Ala Val Ala Ala Val Glu Leu Tyr Gly Gly 115 120 120

Pro Ala Ile Val Val Asp Phe Gly Thr Ala Thr Thr Phe Asp Ala Val 130 135 140

Ser Ala Arg Gly Glu Tyr Ile Gly Gly Val Ile Ala Pro Gly Ile Glu 145 150 155 160

Ile Ser Val Glu Ala Leu Gly Val Lys Gly Ala Gln Leu Arg Lys Ile 165 170 175

Glu Val Ala Arg Pro Arg Ser Val Ile Gly Lys Asn Thr Val Glu Ala 180 185 190

Met Gln Ser Gly Ile Val Tyr Gly Phe Ala Gly Gln Val Asp Gly Val 195 200 205

Val Asn Arg Met Ala Arg Glu Leu Ala Asp Asp Pro Asp Asp Val Thr 210 215 220

Val Ile Ala Thr Gly Gly Leu Ala Pro Met Val Leu Gly Glu Ser Ser 225 230 235 240

Val Ile Asp Glu His Glu Pro Trp Leu Thr Leu Met Gly Leu Arg Leu 245 250 255

Val Tyr Glu Arg Asn Val Ser Arg Met

260 265

<210> 8

<211> 272

<212> PRT

<213> Mycobacterium tuberculosis

<400> 8

Met Leu Leu Ala Ile Asp Val Arg Asn Thr His Thr Val Val Gly Leu
1 5 10 15

Leu Ser Gly Met Lys Glu His Ala Lys Val Val Gln Gln Trp Arg Ile 20 25 30

Arg Thr Glu Ser Glu Val Thr Ala Asp Glu Leu Ala Leu Thr Ile Asp 35 40 45

Gly Leu Ile Gly Glu Asp Ser Glu Arg Leu Thr Gly Thr Ala Ala Leu 50 55 60

Ser Thr Val Pro Ser Val Leu His Glu Val Arg Ile Met Leu Asp Gln 65 70 75 80

Tyr Trp Pro Ser Val Pro His Val Leu Ile Glu Pro Gly Val Arg Thr
85 90 95

Gly Ile Pro Leu Leu Val Asp Asn Pro Lys Glu Val Gly Ala Asp Arg \$100\$

Ile Val Asn Cys Leu Ala Ala Tyr Asp Arg Phe Arg Lys Ala Ala Ile 115 120 125

Val Val Asp Phe Gly Ser Ser Ile Cys Val Asp Val Val Ser Ala Lys 130 135 140

Gly Glu Phe Leu Gly Gly Ala Ile Ala Pro Gly Val Gln Val Ser Ser 145 150 155 160

Asp Ala Ala Ala Arg Ser Ala Ala Leu Arg Arg Val Glu Leu Ala 165 170 175

Arg Pro Arg Ser Val Val Gly Lys Asn Thr Val Glu Cys Met Gln Ala 180 185 190

Gly Ala Val Phe Gly Phe Ala Gly Leu Val Asp Gly Leu Val Gly Arg 195 200 205

Ile Arg Glu Asp Val Ser Gly Phe Ser Val Asp His Asp Val Ala Ile 210 215 220

Val Ala Thr Gly His Thr Ala Pro Leu Leu Pro Glu Leu His Thr 225 230 235 240

Val Asp His Tyr Asp Gln His Leu Thr Leu Gln Gly Leu Arg Leu Val
245 250 255

Phe Glu Arg Asn Leu Glu Val Gln Arg Gly Arg Leu Lys Thr Ala Arg 260 265 270

<210> 9

<211> 258

<212> PRT

<213> Bacillus subtilis

<400> 9

Leu Leu Leu Val Ile Asp Val Gly Asn Thr Asn Thr Val Leu Gly Val

Tyr His Asp Gly Lys Leu Glu Tyr His Trp Arg Ile Glu Thr Ser Arg

His Lys Thr Glu Asp Glu Phe Gly Met Ile Leu Arg Ser Leu Phe Asp

His Ser Gly Leu Met Phe Glu Gln Ile Asp Gly Ile Ile Ile Ser Ser

Val Val Pro Pro Ile Met Phe Ala Leu Glu Arg Met Cys Thr Lys Tyr

Phe His Ile Glu Pro Gln Ile Val Gly Pro Gly Met Lys Thr Gly Leu

Asn Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile Val

Asn Ala Val Ala Ala Ile His Leu Tyr Gly Asn Pro Leu Ile Val Val

Asp Phe Gly Thr Ala Thr Thr Tyr Cys Tyr Ile Asp Glu Asn Lys Gln 135

Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala 155

Leu Tyr Ser Arg Ala Ala Lys Leu Pro Arg Ile Glu Ile Thr Arg Pro 170

Asp Asn Ile Ile Gly Lys Asn Thr Val Ser Ala Met Gln Ser Gly Ile

Leu Phe Gly Tyr Val Gly Gln Val Glu Gly Ile Val Lys Arg Met Lys

Trp Gln Ala Lys Gln Asp Leu Lys Val Ile Ala Thr Gly Gly Leu Ala

Pro Leu Ile Ala Asn Glu Ser Asp Cys Ile Asp Ile Val Asp Pro Phe 225

Leu Thr Leu Lys Gly Leu Glu Leu Ile Tyr Glu Arg Asn Arg Val Gly 250

Ser Val

<210> 10

<211> 262

<212> PRT

<213> Deinococcus radiopugnans

<400> 10

Met Pro Ala Phe Pro Leu Leu Ala Val Asp Ile Gly Asn Thr Thr 1 5 10 15

Val Leu Gly Leu Ala Asp Ala Ser Gly Ala Leu Thr His Thr Trp Arg 20 25 30

Ile Arg Thr Asn Arg Glu Met Leu Pro Asp Asp Leu Ala Leu Gln Leu 35 40 45

His Gly Leu Phe Thr Leu Ala Gly Ala Pro Ile Pro Arg Ala Ala Val 50 55 60

Leu Ser Ser Val Ala Pro Pro Val Gly Glu Asn Tyr Ala Leu Ala Leu 65 70 75 80

Lys Arg His Phe Met Ile Asp Ala Phe Ala Val Ser Ala Glu Asn Leu 85 90 95

Pro Asp Val Thr Val Glu Leu Asp Thr Pro Gly Ser Val Gly Ala Asp 100 105 110

Arg Leu Cys Asn Leu Phe Gly Ala Glu Lys Tyr Leu Gly Gly Leu Asp 115 120 125

Tyr Ala Val Val Val Asp Phe Gly Thr Ser Thr Asn Phe Asp Val Val 130 135 140

Gly Arg Gly Arg Arg Phe Leu Gly Gly Ile Leu Ala Thr Gly Ala Gln 145 150 155 160

Val Ser Ala Asp Ala Leu Phe Ala Arg Ala Ala Lys Leu Pro Arg Ile 165 170 175

Thr Leu Gln Ala Pro Glu Thr Ala Ile Gly Lys Asn Thr Val His Ala 180 185 190

Leu Gln Ser Gly Leu Val Phe Gly Tyr Ala Glu Met Val Asp Gly Leu 195 200 205

Leu Arg Arg Ile Arg Ala Glu Leu Pro Gly Glu Ala Val Ala Val Ala 210 215 220

Thr Gly Gly Phe Ser Arg Thr Val Gln Gly Ile Cys Gln Glu Ile Asp 225 230 235 240

Tyr Tyr Asp Glu Thr Leu Thr Leu Arg Gly Leu Val Glu Leu Trp Ala 245 250 255

Ser Arg Ser Glu Val Arg 260

<210> 11

<211> 212

<212> PRT

<213> Desulfovibrio vulgaris

<400> 11

Met Thr Gln His Phe Leu Leu Phe Asp Ile Gly Asn Thr Asn Val Lys
1 5 10 15

Ile Gly Ile Ala Val Glu Thr Ala Val Leu Thr Ser Tyr Val Leu Pro 20 25 30

Thr Asp Pro Gly Gln Thr Thr Asp Ser Ile Gly Leu Arg Leu Leu Glu 35 40 45

Val Leu Arg His Ala Gly Leu Gly Pro Ala Asp Val Gly Ala Cys Val 50 60

Ala Ser Ser Val Val Pro Gly Val Asn Pro Leu Ile Arg Arg Ala Cys 65 70 75 80

Glu Arg Tyr Leu Tyr Arg Lys Leu Leu Phe Ala Pro Gly Asp Ile Ala 85 90 95

Ile Pro Leu Asp Asn Arg Tyr Glu Arg Pro Ala Glu Val Gly Ala Asp
100 105 110

Arg Leu Val Ala Ala Tyr Ala Ala Arg Arg Leu Tyr Pro Gly Pro Arg 115 120 125

Ser Leu Val Ser Val Asp Phe Gly Thr Ala Thr Thr Phe Asp Cys Val 130 135 140

Ser Ala Gly Ala Leu Ser Ser Arg Thr Ala Lys Leu Pro Arg Ile Ser 165 170 175

Leu Glu Val Glu Glu Asp Ser Pro Val Ile Gly Arg Ser Thr Thr Thr 180 185 190

Ser Leu Asn His Gly Phe Ile Phe Gly Phe Ala Ala Met Thr Glu Gly 195 200 205

Val Leu Ala Ala 210

<210> 12

<211> 246

<212> PRT

<213> Thermotoga maritima

<400> 12

Met Tyr Leu Leu Val Asp Val Gly Asn Thr His Ser Val Phe Ser Ile 1 5 15

Thr Glu Asp Gly Lys Thr Phe Arg Arg Trp Arg Leu Ser Thr Gly Val 20 25 30

Phe Gln Thr Glu Asp Glu Leu Phe Ser His Leu His Pro Leu Leu Gly

Asp Ala Met Arg Glu Ile Lys Gly Ile Gly Val Ala Ser Val Val Pro 50 55 60

Thr Gln Asn Thr Val Ile Glu Arg Phe Ser Gln Lys Tyr Phe His Ile 65 70 75 80

Ser Pro Ile Trp Val Lys Ala Lys Asn Gly Cys Val Lys Trp Asn Val 85 90 95

Lys Asn Pro Ser Glu Val Gly Ala Asp Arg Val Ala Asn Val Val Ala 100 105 110

Phe Val Lys Glu Tyr Gly Lys Asn Gly Ile Ile Ile Asp Met Gly Thr 115 120 125

Ala Thr Thr Val Asp Leu Val Val Asn Gly Ser Tyr Glu Gly Gly Ala 130 135 140

Ile Leu Pro Gly Phe Phe Met Met Val His Ser Leu Phe Arg Gly Thr 145 150 150

Ala Lys Leu Pro Leu Val Glu Val Lys Pro Ala Asp Phe Val Val Gly 165 170 175

Lys Asp Thr Glu Glu Asn Ile Arg Leu Gly Val Val Asn Gly Ser Val
180 185 190

Tyr Ala Leu Glu Gly Ile Ile Gly Arg Ile Lys Glu Val Tyr Gly Asp 195 200 205

Leu Pro Val Val Leu Thr Gly Gly Gln Ser Lys Ile Val Lys Asp Met 210 215 220

Ile Lys His Glu Ile Phe Asp Glu Asp Leu Thr Ile Lys Gly Val Tyr 225 230 235 240

His Phe Cys Phe Gly Asp 245

<210> 13

<211> 273

<212> PRT

<213> Treponema pallidum

<400> 13

Met Leu Leu Ile Asp Val Gly Asn Ser His Val Val Phe Gly Ile Gln 1 5 . 10 15

Gly Glu Asn Gly Gly Arg Val Cys Val Arg Glu Leu Phe Arg Leu Ala 20 25 30

Pro Asp Ala Arg Lys Thr Gln Asp Glu Tyr Ser Leu Leu Ile His Ala 35 40 45

Leu Cys Glu Arg Ala Gly Val Gly Arg Ala Ser Leu Arg Asp Ala Phe 50 55 60

Ile Ser Ser Val Val Pro Val Leu Thr Lys Thr Ile Ala Asp Ala Val 65 70 75 80 Ala Gln Ile Ser Gly Val Gln Pro Val Val Phe Gly Pro Trp Ala Tyr 85 90 95

Glu His Leu Pro Val Arg Ile Pro Glu Pro Val Arg Ala Glu Ile Gly
100 105 110

Thr Asp Leu Val Ala Asn Ala Val Ala Ala Tyr Val His Phe Arg Ser 115 120 125

Ala Cys Val Val Val Asp Cys Gly Thr Ala Leu Thr Phe Thr Ala Val 130 135 140

Asp Gly Thr Gly Leu Ile Gln Gly Val Ala Ile Ala Pro Gly Leu Arg 145 150 155 160

Thr Ala Val Gln Ser Leu His Thr Gly Thr Ala Gln Leu Pro Leu Val 165 170 175

Pro Leu Ala Leu Pro Asp Ser Val Leu Gly Lys Asp Thr Thr His Ala 180 185 190

Val Gln Ala Gly Val Val Arg Gly Thr Leu Phe Val Ile Arg Ala Met 195 200 205

Ile Ala Gln Cys Gln Lys Glu Leu Gly Cys Arg Cys Ala Ala Val Ile 210 215 220

Thr Gly Gly Leu Ser Arg Leu Phe Ser Ser Glu Val Asp Phe Pro Pro 225 230 235 240

Ile Asp Ala Gln Leu Thr Leu Ser Gly Leu Ala His Ile Ala Arg Leu 245 250 255

Val Pro Thr Ser Leu Leu Pro Pro Ala Thr Val Ser Gly Ser Ser Gly 260 265 270

Asn

<210> 14

<211> 262

<212> PRT

<213> Borrelia burgdorferi

<400> 14

Met Asn Lys Pro Leu Leu Ser Glu Leu Ile Ile Asp Ile Gly Asn Thr 1 $$ 15

Ser Ile Ala Phe Ala Leu Phe Lys Asp Asn Gln Val Asn Leu Phe Ile 20 25 30

Lys Met Lys Thr Asn Leu Met Leu Arg Tyr Asp Glu Val Tyr Ser Phe 35 40 45

Phe Glu Glu Asn Phe Asp Phe Asn Val Asn Lys Val Phe Ile Ser Ser 50 55 60

Val Val Pro Ile Leu Asn Glu Thr Phe Lys Asn Val Ile Phe Ser Phe 65 70 75 80

Phe Lys Ile Lys Pro Leu Phe Ile Gly Phe Asp Leu Asn Tyr Asp Leu 85 90 95

Thr Phe Asn Pro Tyr Lys Ser Asp Lys Phe Leu Leu Gly Ser Asp Val

Phe Ala Asn Leu Val Ala Ala Ile Glu Asn Tyr Ser Phe Glu Asn Val 115 120 125

Leu Val Val Asp Leu Gly Thr Ala Cys Thr Ile Phe Ala Val Ser Arg 130 135 140

Gln Asp Gly Ile Leu Gly Gly Ile Ile Asn Ser Gly Pro Leu Ile Asn 145 150 155 160

Phe Asn Ser Leu Leu Asp Asn Ala Tyr Leu Ile Lys Lys Phe Pro Ile 165 170 175

Ser Thr Pro Asn Asn Leu Leu Glu Arg Thr Thr Ser Gly Ser Val Asn 180 185 190

Ser Gly Leu Phe Tyr Gln Tyr Lys Tyr Leu Ile Glu Gly Val Tyr Arg 195 200 205

Asp Ile Lys Gln Met Tyr Lys Lys Phe Asn Leu Ile Ile Thr Gly 210 215 220

Gly Asn Ala Asp Leu Ile Leu Ser Leu Ile Glu Ile Glu Phe Ile Phe 225 230 235 240

Asn Ile His Leu Thr Val Glu Gly Val Arg Ile Leu Gly Asn Ser Ile 245 250 255

Asp Phe Lys Phe Val Asn 260

<210> 15

<211> 229

<212> PRT

<213> Aquifex aeolicus

<400> 15

Met Arg Phe Leu Thr Val Asp Val Gly Asn Ser Ser Val Asp Ile Ala 1 5 10 15

Leu Trp Glu Gly Lys Lys Val Lys Asp Phe Leu Lys Leu Ser His Glu 20 25 30

Glu Phe Leu Lys Glu Glu Phe Pro Lys Leu Lys Ala Leu Gly Ile Ser 35 40 45

Val Lys Gln Ser Phe Ser Glu Lys Val Arg Gly Lys Ile Pro Lys Ile 50 55 60

Lys Phe Leu Lys Lys Glu Asn Phe Pro Ile Gln Val Asp Tyr Lys Thr 65 70 75 80

Pro Glu Thr Leu Gly Thr Asp Arg Val Ala Leu Ala Tyr Ser Ala Lys
85 90 95

Lys Phe Tyr Gly Lys Asn Val Val Val Ile Ser Ala Gly Thr Ala Leu 100 105 110

Val Ile Asp Leu Val Leu Glu Gly Lys Phe Lys Gly Gly Phe Ile Thr 115 120 125

Leu Gly Leu Gly Lys Lys Leu Lys Ile Leu Ser Asp Leu Ala Glu Gly 130 135

Ile Pro Glu Phe Phe Pro Glu Glu Val Glu Ile Phe Leu Gly ArgSer145150155160

Thr Arg Glu Cys Val Leu Gly Gly Ala Tyr Arg Glu Ser Thr Glu Phe 165 170 175

Ile Lys Ser Thr Leu Lys Leu Trp Arg Lys Val Phe Lys Arg Lys Phe 180 185 190

Lys Val Val Ile Thr Gly Gly Glu Gly Lys Tyr Phe Ser Lys Phe Gly 195 200 205

Ile Tyr Asp Pro Leu Leu Val His Arg Gly Met Arg Asn Leu Leu Tyr 210 215 220

Leu Tyr His Arg Ile 225

<210> 16

<211> 257

<212> PRT

<213> Synechocystis sp.

<400> 16

Met Glu Thr Ser Lys Pro Gly Cys Gly Leu Ala Leu Asp Asn Asp Lys

1 5 10 15

Gln Lys Pro Trp Leu Gly Leu Met Ile Gly Asn Ser Arg Leu His Trp 20 25 30

Ala Tyr Cys Ser Gly Asn Ala Pro Leu Gln Thr Trp Val Thr Asp Tyr 35 40 45

Asn Pro Lys Ser Ala Gln Leu Pro Val Leu Leu Gly Lys Val Pro Leu 50 55 60

Met Leu Ala Ser Val Val Pro Glu Gln Thr Glu Val Trp Arg Val Tyr 65 70 75 80

Gln Pro Lys Ile Leu Thr Leu Lys Asn Leu Pro Leu Val Asn Leu Tyr 85 90 95

Pro Ser Phe Gly Ile Asp Arg Ala Leu Ala Gly Leu Gly Thr Gly Leu 100 105 110

Thr Tyr Gly Phe Pro Cys Leu Val Val Asp Gly Gly Thr Ala Leu Thr 115 120 125

Ile Thr Gly Phe Asp Gln Asp Lys Lys Leu Val Gly Gly Ala Ile Leu 130 135 140 Pro Gly Leu Gly Leu Gln Leu Ala Thr Leu Gly Asp Arg Leu Ala Ala 145 150 155 160

Leu Pro Lys Leu Glu Met Asp Gln Leu Thr Glu Leu Pro Asp Arg Trp
165 170 175

Ala Leu Asp Thr Pro Ser Ala Ile Phe Ser Gly Val Val Tyr Gly Val
180 185 190

Leu Gly Ala Leu Gln Ser Tyr Leu Gln Asp Trp Gln Lys Leu Phe Pro 195 200 205

Gly Ala Ala Met Val Ile Thr Gly Gly Asp Gly Lys Ile Leu His Gly 210 215 220

Phe Leu Lys Glu His Ser Pro Asn Leu Ser Val Ala Trp Asp Asp Asn 225 230 235 240

Leu Ile Phe Leu Gly Met Ala Ala Ile His His Gly Asp Arg Pro Ile 245 250 255

Cys

<210> 17

<211> 223

<212> PRT

<213> Helicobacter pylori

<400> 17

Met Pro Ala Arg Gln Ser Phe Thr Asp Leu Lys Asn Leu Val Leu Cys 1 5 10 15

Asp Ile Gly Asn Thr Arg Ile His Phe Ala Gln Asn Tyr Gln Leu Phe 20 25 30

Ser Ser Ala Lys Glu Asp Leu Lys Arg Leu Gly Ile Gln Lys Glu Ile 35 40 45

Phe Tyr Ile Ser Val Asn Glu Glu Asn Glu Lys Ala Leu Leu Asn Cys 50 55 60

Tyr Pro Asn Ala Lys Asn Ile Ala Gly Phe Phe His Leu Glu Thr Asp 65 70 75 80

Tyr Val Gly Leu Gly Ile Asp Arg Gln Met Ala Cys Leu Ala Val Asn 85 90 95

Asn Gly Val Val Asp Ala Gly Ser Ala Ile Thr Ile Asp Leu Ile
100 105 110

Lys Glu Gly Lys His Leu Gly Gly Cys Ile Leu Pro Gly Leu Ala Gln
115 120 125

Tyr Ile His Ala Tyr Lys Lys Ser Ala Lys Ile Leu Glu Gln Pro Phe 130 135

Lys Ala Leu Asp Ser Leu Glu Val Leu Pro Lys Ser Thr Arg Asp Ala 145 150 155 160 Val Asn Tyr Gly Met Val Leu Ser Val Ile Ala Cys Ile Gln His Leu 165 170 175

Ala Lys Asn Gln Lys Ile Tyr Leu Cys Gly Gly Asp Ala Lys Tyr Leu 180 185 190

Ser Ala Phe Leu Pro His Ser Val Cys Lys Glu Arg Leu Val Phe Asp 195 200 205

Gly Met Glu Ile Ala Leu Lys Lys Ala Gly Ile Leu Glu Cys Lys 210 215 220

<210> 18

<211> 267

<212> PRT

<213> Bordetella pertussis

<400> 18

Met Ile Ile Leu Ile Asp Ser Gly Asn Ser Arg Leu Lys Val Gly Trp
1 5 10 15

Phe Asp Pro Asp Ala Pro Gln Ala Ala Arg Glu Pro Ala Pro Val Ala 20 25 30

Phe Asp Asn Leu Asp Leu Asp Ala Leu Gly Arg Trp Leu Ala Thr Leu 35 40 45

Pro Arg Arg Pro Gln Arg Ala Leu Gly Val Asn Val Ala Gly Leu Ala 50 55 60

Arg Gly Glu Ala Ile Ala Ala Thr Leu Arg Ala Gly Gly Cys Asp Ile 65 70 75 80

Arg Trp Leu Arg Ala Gln Pro Leu Ala Met Gly Leu Arg Asn Gly Tyr 85 90 95

Arg Asn Pro Asp Gln Leu Gly Ala Asp Arg Trp Ala Cys Met Val Gly
100 105 110

Val Leu Ala Arg Gln Pro Ser Val His Pro Pro Leu Leu Val Ala Ser 115 120 125

Phe Gly Thr Ala Thr Thr Leu Asp Thr Ile Gly Pro Asp Asn Val Phe 130 135 140

Pro Gly Gly Leu Ile Leu Pro Gly Pro Ala Met Met Arg Gly Ala Leu 145 150 155 160

Ala Tyr Gly Thr Ala His Leu Pro Leu Ala Asp Gly Leu Val Ala Asp 165 170 175

Tyr Pro Ile Asp Thr His Gln Ala Ile Ala Ser Gly Ile Ala Ala Ala 180 185 190

Gln Ala Gly Ala Ile Val Arg Gln Trp Leu Ala Gly Arg Gln Arg Tyr 195 200 205

Gly Gln Ala Pro Glu Ile Tyr Val Ala Gly Gly Gly Trp Pro Glu Val 210 215 220 Arg Gln Glu Ala Glu Arg Leu Leu Ala Val Thr Gly Ala Ala Phe Gly 225 230 235 240

Ala Thr Pro Gln Pro Thr Tyr Leu Asp Ser Pro Val Leu Asp Gly Leu 245 250 255

Ala Ala Leu Ala Ala Gln Gly Ala Pro Thr Ala 260 265

<210> 19

<211> 777

<212> DNA

<213> Bacillus subtilis

100

<220>

<221> CDS

<222> (1)..(774)

<400> 19

ttg tta ctg gtt atc gat gtg ggg aac acc aat act gta ctt ggt gta 48 Leu Leu Leu Val Ile Asp Val Gly Asn Thr Asn Thr Val Leu Gly Val 1 5 10

tat cat gat gga aaa tta gaa tat cac tgg cgt ata gaa aca agc agg 96 Tyr His Asp Gly Lys Leu Glu Tyr His Trp Arg Ile Glu Thr Ser Arg 20 25 30

cat aaa aca gaa gat gag ttt ggg atg att ttg cgc tcc tta ttt gat 144
His Lys Thr Glu Asp Glu Phe Gly Met Ile Leu Arg Ser Leu Phe Asp
35 40 45

cac tcc ggg ctt atg ttt gaa cag ata ggt att att att tcg tca 192
His Ser Gly Leu Met Phe Glu Gln Ile Asp Gly Ile Ile Ile Ser Ser
50 60

gta gtg ccg cca atc atg ttt gcg tta gaa aga atg tgc aca aaa tac 240 Val Val Pro Pro Ile Met Phe Ala Leu Glu Arg Met Cys Thr Lys Tyr 65 70 75 80

ttt cat atc gag cct caa att gtt ggt cca ggt atg aaa acc ggt tta 288
Phe His Ile Glu Pro Gln Ile Val Gly Pro Gly Met Lys Thr Gly Leu
85 90 95

aat ata aaa tat gac aat ccg aaa gaa gta ggg gca gac aga atc gta 336 Asn Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile Val

110

105

aat get get geg ata cac ttg tac ggc aat cca tta att gtt gtc 384
Asn Ala Val Ala Ala Ile His Leu Tyr Gly Asn Pro Leu Ile Val Val
115 120 125

gat ttc gga acc gcc aca acg tac tgc tat att gat gaa aac aaa caa 432 Asp Phe Gly Thr Ala Thr Thr Tyr Cys Tyr Ile Asp Glu Asn Lys Gln 130 135 140

tac atg ggc ggg gcg att gcc cct ggg att aca att tcg aca gag gcg 480 Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala 145 150 155 160

ctt tac tcg cgt gca gca aag ctt cct cgt atc gaa atc acc cgg ccc 528

BGI-141CP

Leu Tyr Ser Arg Ala Ala Lys Leu Pro Arg Ile Glu Ile Thr Arg Pro 170 gac aat att atc gga aaa aac act gtt agc gcg atg caa tct gga att 576 Asp Asn Ile Ile Gly Lys Asn Thr Val Ser Ala Met Gln Ser Gly Ile 185 624 tta ttt qqc tat qtc qqc caa gtg gaa gga atc gtt aag cga atg aaa Leu Phe Gly Tyr Val Gly Gln Val Glu Gly Ile Val Lys Arg Met Lys 200 tgg cag gca aaa cag gac ctc aag gtc att gcg aca gga ggc ctg gcg 672 Trp Gln Ala Lys Gln Asp Leu Lys Val Ile Ala Thr Gly Gly Leu Ala ccg ctc att gcg aac gaa tca gat tgt ata gac atc gtt gat cca ttc 720 Pro Leu Ile Ala Asn Glu Ser Asp Cys Ile Asp Ile Val Asp Pro Phe 225 230 235 tta acc cta aaa ggg ctg gaa ttg att tat gaa aga aac cgc gta gga 768 Leu Thr Leu Lys Gly Leu Glu Leu Ile Tyr Glu Arg Asn Arg Val Gly 245 250 777 agt gta tag Ser Val <210> 20 <211> 960 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(957) <400> 20 gtg aaa aat aaa gaa ctt aac cta cat act tta tat aca cag cac aat 48 Met Lys Asn Lys Glu Leu Asn Leu His Thr Leu Tyr Thr Gln His Asn cgg gag tct tgg tct ggt ttt ggg ggg cat ttg tcg att gct gta tct 96 Arg Glu Ser Trp Ser Gly Phe Gly Gly His Leu Ser Ile Ala Val Ser 2.0 25 gaa gaa gag gca aaa gct gtg gaa gga ttg aat gat tat cta tct gtt 144 Glu Glu Glu Ala Lys Ala Val Glu Gly Leu Asn Asp Tyr Leu Ser Val 40 gaa gaa gtg gag acg atc tat att ccg ctt gtt cgc ttg ctt cat tta 192 Glu Glu Val Glu Thr Ile Tyr Ile Pro Leu Val Arg Leu Leu His Leu 50 55 cat gtc aag tct gcg gct gaa cgc aat aag cat gtc aat gtt ttt ttg 240 His Val Lys Ser Ala Ala Glu Arg Asn Lys His Val Asn Val Phe Leu 65 70 75

aag cac cca cat tca gcc aaa att ccg ttt att atc ggc att gcc ggc

Lys His Pro His Ser Ala Lys Ile Pro Phe Ile Ile Gly Ile Ala Gly

90

85

288

- 20 -

~	~	_	_	gga Gly		-	_					_	_		_	336
				cct Pro												384
				cct Pro												432
-				cct Pro	-	_		_	-	-		-			_	480
_		-		aaa Lys 165			_	-	_	_	_	_	_			528
				tat Tyr												576
_	-	_		gtg Val			_				-		_	-		624
	_		_	gac Asp		-		-								672
	-			att Ile												720
				cgt Arg 245		_	_			_		-				768
	_			ttt Phe					_	_		-	_			816
		_		gcc Ala	_				-	_			_			864
	-			ttg Leu							_			~	-	912
_		_		cat His	_	_				_	-	_			tga	960

<210> 21

<211> 882

<212> DNA

<213> Bacillus subtilis

<220> <221> CDS <222> (1)..(879)

<400> 21 ttg tcg att gct gta tct gaa gag gca aaa gct gtg gaa gga ttg 48 Met Ser Ile Ala Val Ser Glu Glu Glu Ala Lys Ala Val Glu Gly Leu aat gat tat cta tct gtt gaa gaa gtg gag acg atc tat att ccg ctt 96 Asn Asp Tyr Leu Ser Val Glu Glu Val Glu Thr Ile Tyr Ile Pro Leu 2.0 gtt cgc ttg ctt cat tta cat gtc aag tct gcg gct gaa cgc aat aag 144 Val Arg Leu His Leu His Val Lys Ser Ala Ala Glu Arg Asn Lys 35 cat gtc aat gtt ttt ttg aag cac cca cat tca gcc aaa att ccg ttt 192 His Val Asn Val Phe Leu Lys His Pro His Ser Ala Lys Ile Pro Phe 50 att atc ggc att gcc ggc agt gtc gca gtc gga aaa agc acg acg gcg 240 Ile Ile Gly Ile Ala Gly Ser Val Ala Val Gly Lys Ser Thr Thr Ala cgg atc ttg cag aag ctg ctt tcg cgt ttg cct gac cgt cca aaa gtg 288 Arg Ile Leu Gln Lys Leu Leu Ser Arg Leu Pro Asp Arg Pro Lys Val agc ctt atc acg aca gat ggt ttt tta ttt cct act gcc gag ctg aaa 336 Ser Leu Ile Thr Thr Asp Gly Phe Leu Phe Pro Thr Ala Glu Leu Lys 100 105 aag aaa aat atg atg tca aga aaa gga ttt cct gaa agc tat gat gta Lys Lys Asn Met Met Ser Arg Lys Gly Phe Pro Glu Ser Tyr Asp Val 115 120 aag gcg ctg ctc gaa ttt ttg aat gac tta aaa tca gga aag gac agc 432 Lys Ala Leu Leu Glu Phe Leu Asn Asp Leu Lys Ser Gly Lys Asp Ser 130 135 gta aag gcc ccg gtg tat tcc cat cta acc tat gac cgc gag gaa ggt 480 Val Lys Ala Pro Val Tyr Ser His Leu Thr Tyr Asp Arg Glu Glu Gly 145 155 150 gtg ttc gag gtt gta gaa cag gcg gat att gtg att att gaa ggc att 528 Val Phe Glu Val Val Glu Gln Ala Asp Ile Val Ile Ile Glu Gly Ile 175 165 170 aat gtt ctt cag tcg ccc acc ttg gag gat gac cgg gaa aac ccg cgt 576 Asn Val Leu Gln Ser Pro Thr Leu Glu Asp Asp Arg Glu Asn Pro Arg 180 185 att ttt gtt tcc gat ttc ttt gat ttt tcg att tat gtg gat gcg gag 624 Ile Phe Val Ser Asp Phe Phe Asp Phe Ser Ile Tyr Val Asp Ala Glu 195 200 gaa agc cgg att ttc act tgg tat tta gag cgt ttt cgc ctg ctt cgg 672 Glu Ser Arg Ile Phe Thr Trp Tyr Leu Glu Arg Phe Arg Leu Leu Arg

215

gaa aca gct ttt caa aat cct gat tca tat ttt cat aaa Glu Thr Ala Phe Gln Asn Pro Asp Ser Tyr Phe His Lys 225 230 235	
ttg tcc gat cag gag gct gac gag atg gca gcc tcg att Leu Ser Asp Gln Glu Ala Asp Glu Met Ala Ala Ser Ile ' 245 250	
gtc aac cgg ccg aat tta tat gaa aat att ttg cca act Val Asn Arg Pro Asn Leu Tyr Glu Asn Ile Leu Pro Thr 260 265	
tca gat ctc att ttg cgt aag gga gac ggg cat aag gtc Ser Asp Leu Ile Leu Arg Lys Gly Asp Gly His Lys Val 275 280 285	
ttg gta agg agg gta tga Leu Val Arg Arg Val 290	882
<210> 22 <211> 846 <212> DNA <213> Bacillus subtilis	
<220> <221> CDS <222> (1)(843)	
<pre><400> 22 gtg gaa gga ttg aat gat tat cta tct gtt gaa gaa gtg Met Glu Gly Leu Asn Asp Tyr Leu Ser Val Glu Glu Val 1 5 10</pre>	
tat att ccg ctt gtt cgc ttg ctt cat tta cat gtc aag Tyr Ile Pro Leu Val Arg Leu Leu His Leu His Val Lys 20 25	
gaa cgc aat aag cat gtc aat gtt ttt ttg aag cac cca Glu Arg Asn Lys His Val Asn Val Phe Leu Lys His Pro 35 40 45	•
aaa att ccg ttt att atc ggc att gcc ggc agt gtc gca Lys Ile Pro Phe Ile Ile Gly Ile Ala Gly Ser Val Ala 50 55 60	
agc acg acg gcg cgg atc ttg cag aag ctg ctt tcg cgt Ser Thr Thr Ala Arg Ile Leu Gln Lys Leu Leu Ser Arg 65 70 75	
cgt cca aaa gtg agc ctt atc acg aca gat ggt ttt tta Arg Pro Lys Val Ser Leu Ile Thr Thr Asp Gly Phe Leu 85 90	
gcc gag ctg aaa aag aaa aat atg atg tca aga aaa gga Ala Glu Leu Lys Lys Lys Asn Met Met Ser Arg Lys Gly 100 105	=

Ser	Tyr	Asp 115	Val	Lys	Ala	Leu	Leu 120	Glu	Phe	Leu	Asn	Asp 125	Leu	Lys	Ser	
gga Gly	aag Lys 130	gac Asp	agc Ser	gta Val	aag Lys	gcc Ala 135	ccg Pro	gtg Val	tat Tyr	tcc Ser	cat His 140	cta Leu	acc Thr	tat Tyr	gac Asp	432
cgc Arg 145	gag Glu	gaa Glu	ggt Gly	gtg Val	ttc Phe 150	gag Glu	gtt Val	gta Val	gaa Glu	cag Gln 155	gcg Ala	gat Asp	att Ile	gtg Val	att Ile 160	480
att Ile	gaa Glu	ggc Gly	att Ile	aat Asn 165	gtt Val	ctt Leu	cag Gln	tcg Ser	ccc Pro 170	acc Thr	ttg Leu	gag Glu	gat Asp	gac Asp 175	cgg Arg	528
gaa Glu	aac Asn	ccg Pro	cgt Arg 180	att Ile	ttt Phe	gtt Val	tcc Ser	gat Asp 185	ttc Phe	ttt Phe	gat Asp	ttt Phe	tcg Ser 190	att Ile	tat Tyr	576
gtg Val	gat Asp	gcg Ala 195	gag Glu	gaa Glu	agc Ser	cgg Arg	att Ile 200	ttc Phe	act Thr	tgg Trp	tat Tyr	tta Leu 205	gag Glu	cgt Arg	ttt Phe	624
cgc Arg	ctg Leu 210	ctt Leu	cgg Arg	gaa Glu	aca Thr	gct Ala 215	ttt Phe	caa Gln	aat Asn	cct Pro	gat Asp 220	tca Ser	tat Tyr	ttt Phe	cat His	672
aaa Lys 225	ttt Phe	aaa Lys	gac Asp	ttg Leu	tcc Ser 230	gat Asp	cag Gln	gag Glu	gct Ala	gac Asp 235	Glu	atg Met	gca Ala	gcc Ala	tcg Ser 240	720
att Ile	tgg Trp	gag Glu	agt Ser	gtc Val 245	aac Asn	cgg Arg	ccg Pro	aat Asn	tta Leu 250	Tyr	gaa Glu	aat Asn	att Ile	ttg Leu 255	Pro	768
act Thr	aaa Lys	ttc Phe	agg Arg 260	Ser	gat Asp	ctc Leu	att Ile	ttg Leu 265	Arg	aag Lys	gga Gly	gac Asp	ggg Gly 270	His	aag Lys	816
gtc Val	gag Glu	gaa Glu 275	. Val	, ttg . Leu	gta Val	agg Arg	agg Arg 280	Val	ı tga	ı						846
<21 <21	0> 2 1> 8 2> I 3> B	331 DNA	.lus	subt	ilis	3										
	1> 0	CDS (1)	. (83:	1)												
ato	Ly:	a aca	a aaa c Lya	s Lei	g gat ı Ası	ttt p Phe	t cta e Lei	a aaa 1 Ly:	a atq s Met	t Ly:	g gag s Gl	g tc u Se:	t gaa r Glu	a gaa 1 Gl	a ccg u Pro 5	48
att Ile	gto Vai	c ato	g cto Leo 2	u Thi	c gct	t tai	t gat r Asj	ta o Ty 2	r Pro	g gc	a gc a Al	t aaa a Ly	a cti s Lei 3	ı Al	t gaa a Glu	96

caa Gln	gcg Ala	gga Gly 35	gtt Val	gac Asp	atg Met	att Ile	tta Leu 40	gtc Val	ggt Gly	gat Asp	tca Ser	ctt Leu 45	gga Gly	atg Met	gtc Val	144
gtc Val	ctc Leu 50	ggc Gly	ctt Leu	gat Asp	tca Ser	act Thr 55	gtc Val	ggt Gly	gtg Val	aca Thr	gtt Val 60	gcg Ala	gac Asp	atg Met	atc Ile	192
cat His 65	cat His	aca Thr	aaa Lys	gcc Ala	gtt Val 70	aaa Lys	agg Arg	ggt Gly	gcg Ala	ccg Pro 75	aat Asn	acc Thr	ttt Phe	att Ile	gtg Val 80	240
aca Thr	gat Asp	atg Met	ccg Pro	ttt Phe 85	atg Met	tct Ser	tat Tyr	cac His	ctg Leu 90	tct Ser	aag Lys	gaa Glu	gat Asp	acg Thr 95	ctg Leu	288
aaa Lys	aat Asn	gca Ala	gcg Ala 100	gct Ala	atc Ile	gtt Val	cag Gln	gaa Glu 105	agc Ser	gga Gly	gct Ala	gac Asp	gca Ala 110	ctg Leu	aag Lys	336
ctt Leu	gag Glu	ggc Gly 115	gga Gly	gaa Glu	ggc Gly	gtg Val	ttt Phe 120	gaa Glu	tcc Ser	att Ile	cgc Arg	gca Ala 125	ttg Leu	acg Thr	ctt Leu	384
gga Gly	ggc Gly 130	att Ile	cca Pro	gta Val	gtc Val	agt Ser 135	cac His	tta Leu	ggt Gly	ttg Leu	aca Thr 140	ccg Pro	cag Gln	tca Ser	gtc Val	432
ggc Gly 145	gta Val	ctg Leu	ggc	ggc	tat Tyr 150	aaa Lys	gta Val	cag Gln	ggc Gly	aaa Lys 155	gac Asp	gaa Glu	caa Gln	agc Ser	gcc Ala 160	480
aaa Lys	aaa Lys	tta Leu	ata Ile	gaa Glu 165	Asp	agt Ser	ata Ile	aaa Lys	tgc Cys 170	Glu	gaa Glu	gca Ala	gga Gly	gct Ala 175	atg Met	528
atg Met	ctt Leu	gtg Val	ctg Leu 180	Glu	tgt Cys	gtg Val	ccg Pro	gca Ala 185	. Glu	ctc Leu	aca Thr	gcc Ala	aaa Lys 190	Il∈	gcc Ala	576
gag Glu	acg Thr	cta Leu 195	. Ser	ata Ile	ccg Pro	gto Val	att 11e 200	e Gly	ato Ile	e Gly	gct Ala	ggt Gly 205	v Val	aaa Lys	a gcg s Ala	624
gac Asp	gga Gly 210	Glr.	ı gtt ı Val	cto Leu	: gtt . Val	tat Tyr 215	: His	gat Asp	att Ile	ato e Ile	ggc Gl ₂ 220	/ His	ggt Gly	gtt Val	gag L Glu	672
aga Arg 225	J Thr	cct Pro	aaa Lys	a ttt s Phe	gta Val 230	Lys	g caa s Glr	a tat n Tyr	acg Thi	g cgo r Arg 235	g Ile	gat Asp	gaa Glu	a aco	atc Ile 240	720
gaa Glu	a aca ı Thi	a gca : Ala	a ato	c ago e Sei 245	Gl3	a tat y Ty:	c gtt r Val	cag l Glr	g gat n Asp 250	o Val	a aga l Arg	a cat g His	cgt Arg	g Ala 25!	t ttc a Phe	768
cct Pro	gaa Glu	a caa ı Glı	a aag n Lys 26	s His	t tco	tti Pho	t caa e Gli	a ato n Mei 26!	t Ası	c caq n Gli	g aca	a gto r Vai	g cti Let 27	ı As	c ggc p Gly	816

ttg tac ggg gga aaa Leu Tyr Gly Gly Lys 275 831

<210> 24

<211> 277

<212> PRT

<213> Bacillus subtilis

<400> 24

Met Lys Thr Lys Leu Asp Phe Leu Lys Met Lys Glu Ser Glu Glu Pro 1 5 10

Ile Val Met Leu Thr Ala Tyr Asp Tyr Pro Ala Ala Lys Leu Ala Glu 20 25 30

Gln Ala Gly Val Asp Met Ile Leu Val Gly Asp Ser Leu Gly Met Val \$35\$

Val Leu Gly Leu Asp Ser Thr Val Gly Val Thr Val Ala Asp Met Ile 50 60

His His Thr Lys Ala Val Lys Arg Gly Ala Pro Asn Thr Phe Ile Val 65 70 75 80

Thr Asp Met Pro Phe Met Ser Tyr His Leu Ser Lys Glu Asp Thr Leu 85 90 95

Lys Asn Ala Ala Ala Ile Val Gln Glu Ser Gly Ala Asp Ala Leu Lys
100 105 110

Leu Glu Gly Glu Gly Val Phe Glu Ser Ile Arg Ala Leu Thr Leu 115 120 125

Gly Gly Ile Pro Val Val Ser His Leu Gly Leu Thr Pro Gln Ser Val 130 135 140

Gly Val Leu Gly Gly Tyr Lys Val Gln Gly Lys Asp Glu Gln Ser Ala 145 150 155 160

Lys Lys Leu Ile Glu Asp Ser Ile Lys Cys Glu Glu Ala Gly Ala Met 165 170 175

Met Leu Val Leu Glu Cys Val Pro Ala Glu Leu Thr Ala Lys Ile Ala 180 185 190

Glu Thr Leu Ser Ile Pro Val Ile Gly Ile Gly Ala Gly Val Lys Ala 195 200 205

Asp Gly Gln Val Leu Val Tyr His Asp Ile Ile Gly His Gly Val Glu 210 215 220

Arg Thr Pro Lys Phe Val Lys Gln Tyr Thr Arg Ile Asp Glu Thr Ile 225 230 235 240

Glu Thr Ala Ile Ser Gly Tyr Val Gln Asp Val Arg His Arg Ala Phe 245 250 255

Pro Glu Gln Lys His Ser Phe Gln Met Asn Gln Thr Val Leu Asp Gly 260 265 270

Leu Tyr Gly Gly Lys 275

<210> 25 <211> 858 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(858) <400> 25 atg aga cag att act gat att tca cag ctg aaa gaa gcc ata aaa caa Met Arg Gln Ile Thr Asp Ile Ser Gln Leu Lys Glu Ala Ile Lys Gln tac cat tca gag ggc aag tca atc gga ttt gtt ccg acg atg ggg ttt 96 Tyr His Ser Glu Gly Lys Ser Ile Gly Phe Val Pro Thr Met Gly Phe ctg cat gag ggg cat tta acc tta gca gac aaa gca aga caa gaa aac 144 Leu His Glu Gly His Leu Thr Leu Ala Asp Lys Ala Arg Gln Glu Asn 192 gac gcc gtt att atg agt att ttt gtg aat cct gca caa ttc ggc cct Asp Ala Val Ile Met Ser Ile Phe Val Asn Pro Ala Gln Phe Gly Pro 55 aat gaa gat ttt gaa gca tat ccg cgc gat att gag cgg gat gca gct 240 Asn Glu Asp Phe Glu Ala Tyr Pro Arg Asp Ile Glu Arg Asp Ala Ala ctt gca gaa aac gcc gga gtc gat att ctt ttt acg cca gat gct cat 288 Leu Ala Glu Asn Ala Gly Val Asp Ile Leu Phe Thr Pro Asp Ala His 85 gat atg tat ccc ggt gaa aag aat gtc acg att cat gta gaa aga cgc 336 Asp Met Tyr Pro Gly Glu Lys Asn Val Thr Ile His Val Glu Arg Arg 105 aca gac gtg tta tgc ggg cgc tca aga gaa gga cat ttt gac ggg gtc 384 Thr Asp Val Leu Cys Gly Arg Ser Arg Glu Gly His Phe Asp Gly Val 120 115 gcg atc gta ctg acg aag ctt ttc aat cta gtc aag ccg act cgt gcc 432 Ala Ile Val Leu Thr Lys Leu Phe Asn Leu Val Lys Pro Thr Arg Ala 135 130 tat ttc ggt tta aaa gat gcg cag cag gta gct gtt gtt gat ggg tta 480 Tyr Phe Gly Leu Lys Asp Ala Gln Gln Val Ala Val Asp Gly Leu 155 145 528 atc agc gac ttc ttc atg gat att gaa ttg gtt cct gtc gat acg gtc Ile Ser Asp Phe Phe Met Asp Ile Glu Leu Val Pro Val Asp Thr Val 170 165 576 aga gag gaa gac ggc tta gcc aaa agc tct cgc aat gta tac tta aca

Arg Glu Glu Asp Gly Leu Ala Lys Ser Ser Arg Asn Val Tyr Leu Thr

190 185 180 gct gag gaa aga aaa gaa gcg cct aag ctg tat cgg gcc ctt caa aca 624 Ala Glu Glu Arg Lys Glu Ala Pro Lys Leu Tyr Arg Ala Leu Gln Thr 200 195 agt gcg gaa ctt gtc caa gcc ggt gaa aga gat cct gaa gcg gtg ata 672 Ser Ala Glu Leu Val Gln Ala Gly Glu Arg Asp Pro Glu Ala Val Ile 210 215 aaa gct gca aaa gat atc att gaa acg act agc gga acc ata gac tat 720 Lys Ala Ala Lys Asp Ile Ile Glu Thr Thr Ser Gly Thr Ile Asp Tyr 235 230 225 gta gag ctt tat tcc tat ccg gaa ctc gag cct gtg aat gaa att gct Val Glu Leu Tyr Ser Tyr Pro Glu Leu Glu Pro Val Asn Glu Ile Ala 816 gga aag atg att ctc gct gtt gca gtt gct ttt tca aaa gcg cgt tta Gly Lys Met Ile Leu Ala Val Ala Val Ala Phe Ser Lys Ala Arg Leu 260 ata gat aat atc att att gat att cga gaa atg gag aga ata 858 Ile Asp Asn Ile Ile Ile Asp Ile Arg Glu Met Glu Arg Ile <210> 26 <211> 286 <212> PRT <213> Bacillus subtilis <400> 26 Met Arg Gln Ile Thr Asp Ile Ser Gln Leu Lys Glu Ala Ile Lys Gln Tyr His Ser Glu Gly Lys Ser Ile Gly Phe Val Pro Thr Met Gly Phe Leu His Glu Gly His Leu Thr Leu Ala Asp Lys Ala Arg Gln Glu Asn Asp Ala Val Ile Met Ser Ile Phe Val Asn Pro Ala Gln Phe Gly Pro Asn Glu Asp Phe Glu Ala Tyr Pro Arg Asp Ile Glu Arg Asp Ala Ala Leu Ala Glu Asn Ala Gly Val Asp Ile Leu Phe Thr Pro Asp Ala His Asp Met Tyr Pro Gly Glu Lys Asn Val Thr Ile His Val Glu Arg Arg 105 Thr Asp Val Leu Cys Gly Arg Ser Arg Glu Gly His Phe Asp Gly Val 115 Ala Ile Val Leu Thr Lys Leu Phe Asn Leu Val Lys Pro Thr Arg Ala 135

Tyr Phe Gly Leu Lys Asp Ala Gln Gln Val Ala Val Asp Gly Leu

145					150					155					160	
Ile	Ser	Asp	Phe	Phe 165	Met	Asp	Ile	Glu	Leu 170	Val	Pro	Val	Asp	Thr 175	Val	
Arg	Glu	Glu	Asp 180	Gly	Leu	Ala	Lys	Ser 185	Ser	Arg	Asn	Val	Туr 190	Leu	Thr	
Ala	Glu	Glu 195	Arg	Lys	Glu	Ala	Pro 200	Lys	Leu	Tyr	Arg	Ala 205	Leu	Gln	Thr	
Ser	Ala 210	Glu	Leu	Val	Gln	Ala 215	Gly	Glu	Arg	Asp	Pro 220	Glu	Ala	Val	Ile	
Lys 225	Ala	Ala	Lys	Asp	Ile 230	Ile	Glu	Thr	Thr	Ser 235	Gly	Thr	Ile	Asp	Tyr 240	
Val	Glu	Leu	Tyr	Ser 245	Tyr	Pro	Glu	Leu	Glu 250	Pro	Val	Asn	Glu	Ile 255	Ala	
Gly	Lys	Met	Ile 260	Leu	Ala	Val	Ala	Val 265	Ala	Phe	Ser	Lys	Ala 270	Arg	Leu	
Ile	Asp	Asn 275	Ile	Ile	Ile	Asp	Ile 280	Arg	Glu	Met	Glu	Arg 285	Ile			
<21 <21 <21 <22 <22	0> 1> C	81 NA acil DS	lus (381		ilis											
atq	Tyr	cga	aca Thr	atg Met	Met	agc Ser	ggc	aaa Lys	ctt Leu 10	His	agg Arg	gca Ala	act Thr	gtt Val 15	acg Thr	48
gaa Glu	gca Ala	aac Asn	ctg Leu 20	. Asn	tat Tyr	gtg Val	gga Gly	ago Ser 25	Ile	aca Thr	att : Ile	gat Asp	gaa Glu 30	gat Asp	ctc Leu	96
att Ile	gat Asp	gct Ala 35	ı Val	. Gly	atg Met	g ctt Leu	cct Pro 40	Asr	gaa Glu	ı aaa ı Lys	a gta s Val	caa Gln 45	ıIl∈	gtg Val	aat Asn	144
aat Asr	aat Asr 50	ı Asr	gga Gly	a gca 7 Ala	a cgt a Arg	ctt g Leu 55	ı Glu	a acg ı Thi	tat Tyr	att	att Ile	Pro	ggt Gl	aaa y Lys	cgg Arg	192
gga Gly 65	ser Ser	: ggd	gto Val	c ata L Ile	a tgo e Cys 70	s Lev	a aad 1 Asi	c ggt n Gly	gca Ala	a gco a Ala 75	a Ala	a cgc	ctt Lei	gtg 1 Val	g cag Gln 80	240
gaa Glu	a gga ı Gly	a gat 7 Asp	aag Lys	g gto s Val	l Ile	t att e Ile	ati	t tco e Sei	c tac Ty:	r Lys	a atg s Met	g atg : Met	tct Sei	t gat r Asp 95	caa Gln	288

gaa Glu	gcg Ala	gca Ala	agc Ser 100	cat His	gag Glu	ccg Pro	aaa Lys	gtg Val 105	gct Ala	gtt Val	ctg Leu	Asn	gat Asp 110	caa Gln	aac Asn	336
aaa Lys	att Ile	gaa Glu 115	caa Gln	atg Met	ctg Leu	GJA aaa	aac Asn 120	gaa Glu	cca Pro	gcc Ala	cgt Arg	aca Thr 125	att Ile	ttg Leu		381
<213 <213	0> 28 1> 12 2> PF 3> Ba	27 RT	lus :	subti	ilis											
<40 Met	0> 28 Tyr	3 Arg	Thr	Met 5	Met	Ser	Gly	Lys	Leu 10	His	Arg	Ala	Thr	Val 15	Thr	
Glu	Ala	Asn	Leu 20	Asn	Tyr	Val	Gly	Ser 25	Ile	Thr	Ile	Asp	G1u 30	Asp	Leu	
Ile	Asp	Ala 35	Val	Gly	Met	Leu	Pro 40	Asn	Glu	Lys	Val	Gln 45	Ile	Val	Asn	
Asn	Asn 50	Asn	Gly	Ala	Arg	Leu 55	Glu	Thr	Tyr	Ile	Ile 60	Pro	Gly	Lys	Arg	
Gly 65		Gly	Val	Ile	Cys 70	Leu	Asn	Gly	Ala	Ala 75	Ala	Arg	Leu	Val	Gln 80	
Glu	Gly	Asp	Lys	Val 85	Ile	Ile	Ile	Ser	Туr 90	Lys	Met	Met	Ser	Asp 95	Gln	
Glu	Ala	Ala	Ser 100		Glu	Pro	Lys	Val 105		Val	Leu	Asn	Asp 110		Asn	
Lys	Ile	Glu 115		. Met	Leu	Gly	Asn 120		Pro	Ala	Arg	Thr 125	Ile	Leu		
<21 <21	.0> 2 .1> 8 .2> D	94 NA	.lus	subt	ilis											
	20> 21> C 22> (. (894	1)												
ato Met	00> 2 g aaa : Lys l	att	: gga e Gly	a att 7 Ile	: Ile	ggc Gly	gga Gly	ggo Gly	tco Ser 10	· Val	ggt L Gly	ctt Leu	. tta . Lev	tgc Cys 15	Ala	48
tai Ty:	tat r Tyi	t ttg Lei	g tca 1 Sei 20	r Let	tat ı Tyr	cac His	c gad s Asp	gtg Val	Thi	gtt Val	t gtg l Val	g acg Thr	agg Arg 30	, Arg	caa Gln	96
ga: Gl:	a cag u Glr	g gct n Ala	a Ala	g gco a Ala	c att	caç e Glr	g tct n Sei 40	c Gli	a gga ı Gly	a ato	c cgg e Arg	g ctt g Leu 45	ι Туз	aaa Lys	ggc Gly	144

Gly ggg	gag Glu 50	gaa Glu	ttc Phe	agg Arg	gct Ala	gat Asp 55	tgc Cys	agt Ser	gcg Ala	gac Asp	acg Thr 60	agt Ser	atc Ile	aat Asn	tcg Ser	192
gac Asp 65	ttt Phe	gac Asp	ctg Leu	ctt Leu	gtc Val 70	gtg Val	aca Thr	gtg Val	aag Lys	cag Gln 75	cat His	cag Gln	ctt Leu	caa Gln	tct Ser 80	240
gtt Val	ttt Phe	tcg Ser	tcg Ser	ctt Leu 85	gaa Glu	cga Arg	atc Ile	ggg Gly	aag Lys 90	acg Thr	aat Asn	ata Ile	tta Leu	ttt Phe 95	ttg Leu	288
caa Gln	aac Asn	ggc Gly	atg Met 100	gly ggg	cat His	atc Ile	cac His	gac Asp 105	cta Leu	aaa Lys	gac Asp	tgg Trp	cac His 110	gtt Val	ggc Gly	336
cat His	tcc Ser	att Ile 115	tat Tyr	gtt Val	gga Gly	atc Ile	gtt Val 120	gag Glu	cac His	gga Gly	gct Ala	gta Val 125	aga Arg	aaa Lys	tcg Ser	384
gat Asp	aca Thr 130	gct Ala	gtt Val	gat Asp	cat His	aca Thr 135	ggc Gly	cta Leu	ggt Gly	gcg Ala	ata Ile 140	aaa Lys	tgg Trp	agc Ser	gcg Ala	432
ttc Phe 145	gac Asp	gat Asp	gct Ala	gaa Glu	cca Pro 150	gac Asp	cgg Arg	ctg Leu	aac Asn	atc Ile 155	ttg Leu	ttt Phe	cag Gln	cat His	aac Asn 160	480
cat His	tcg Ser	gat Asp	ttt Phe	ccg Pro 165	att Ile	tat Tyr	tat Tyr	gag Glu	acg Thr 170	gat Asp	tgg Trp	tac Tyr	cgt Arg	ctg Leu 175	ctg Leu	528
acg Thr	ggc Gly	aag Lys	ctg Leu 180	att Ile	gta Val	aat Asn	gcg Ala	tgt Cys 185	Ile	aat Asn	cct Pro	tta Leu	act Thr 190	Ala	tta Leu	576
ttg Leu	caa Gln	gtg Val 195	aaa Lys	aat Asn	gga Gly	gaa Glu	ctg Leu 200	ctg Leu	aca Thr	acg Thr	cca Pro	gct Ala 205	tat Tyr	ctg Leu	gct Ala	624
ttt Phe	atg Met 210	. Lys	ctg Leu	gta Val	ttt Phe	cag Gln 215	Glu	gca Ala	tgc Cys	cgc Arg	att ; Ile 220	tta Leu	aaa Lys	ctt Leu	gaa Glu	672
aat Asr 225	ı Glu	ı gaa ı Glu	aag Lys	gct Ala	tgg Trp 230	Glu	cgg Arg	gtt Val	cag Gln	gcc Ala 235	ı Val	tgt Cys	ggg	g caa g Gln	acg Thr 240	720
aaa Lys	ı gaç Glu	ı aatı ı Asn	. cgt . Arg	tca Ser 245	Ser	atg Met	ctg Leu	gtt Val	gac Asp 250	Va]	c att L Ile	gga Gly	ggc Gly	c cgg 7 Arg 255	cag Gln	768
acg Thi	g gaa Glu	a gct ı Ala	gac As <u>r</u> 260	Ala	att a Ile	ato E Ile	gga Gly	tac Tyr 265	: Let	ı ttg ı Lei	g aag 1 Lys	g gaa s Glu	a gca a Ala 270	a Ser	ctt Leu	816
caa Glr	a ggt n Gly	ctt Leu 275	ı Ası	gco Ala	gto a Val	c cac L His	cta Leu 280	ı Glı	g ttt 1 Phe	tta e Lei	a tat u Tyi	ggc Gl ₃ 285	y Sei	c ato	aaa Lys	864

gca ttg gag cga aat aca aac aaa gtc ttt Ala Leu Glu Arg Asn Thr Asn Lys Val Phe 290 295

894

<210> 30

<211> 298

<212> PRT

<213> Bacillus subtilis

<400> 30

Met Lys Ile Gly Ile Ile Gly Gly Gly Ser Val Gly Leu Leu Cys Ala 1 5 10 15

Tyr Tyr Leu Ser Leu Tyr His Asp Val Thr Val Val Thr Arg Arg Gln $20 \hspace{1cm} 25 \hspace{1cm} 30 \hspace{1cm}$

Glu Gln Ala Ala Ile Gln Ser Glu Gly Ile Arg Leu Tyr Lys Gly 35 40 45

Gly Glu Glu Phe Arg Ala Asp Cys Ser Ala Asp Thr Ser Ile Asn Ser 50 55 60

Asp Phe Asp Leu Leu Val Val Thr Val Lys Gln His Gln Leu Gln Ser 65 70 75 80

Val Phe Ser Ser Leu Glu Arg Ile Gly Lys Thr Asn Ile Leu Phe Leu 85 90 95

Gln Asn Gly Met Gly His Ile His Asp Leu Lys Asp Trp His Val Gly 100 105 110

His Ser Ile Tyr Val Gly Ile Val Glu His Gly Ala Val Arg Lys Ser 115 120 125

Asp Thr Ala Val Asp His Thr Gly Leu Gly Ala Ile Lys Trp Ser Ala 130 135 140

Phe Asp Asp Ala Glu Pro Asp Arg Leu Asn Ile Leu Phe Gln His Asn 145 150 155 160

His Ser Asp Phe Pro Ile Tyr Tyr Glu Thr Asp Trp Tyr Arg Leu Leu 165 170 175

Thr Gly Lys Leu Ile Val Asn Ala Cys Ile Asn Pro Leu Thr Ala Leu 180 185 190

Leu Gln Val Lys Asn Gly Glu Leu Leu Thr Thr Pro Ala Tyr Leu Ala 195 200 205

Phe Met Lys Leu Val Phe Gln Glu Ala Cys Arg Ile Leu Lys Leu Glu 210 215 220

Asn Glu Glu Lys Ala Trp Glu Arg Val Gln Ala Val Cys Gly Gln Thr 225 230 235 240

Lys Glu Asn Arg Ser Ser Met Leu Val Asp Val Ile Gly Gly Arg Gln 245 250 255

Thr Glu Ala Asp Ala Ile Ile Gly Tyr Leu Leu Lys Glu Ala Ser Leu 260 265 270

Gln Gly Leu Asp Ala Val His Leu Glu Phe Leu Tyr Gly Ser Ile Lys 275 280 285

Ala Leu Glu Arg Asn Thr Asn Lys Val Phe 290 295

<210> 31

<211> 1725

<212> DNA

<213> Bacillus subtilis

<220>

<221> CDS

<222> (1)..(1722)

<400> 31

atg ggg act aat gta cag gtg gat tca gca tct gcc gaa tgt aca cag

Met Gly Thr Asn Val Gln Val Asp Ser Ala Ser Ala Glu Cys Thr Gln

10 15

acg atg agc gga gca tta atg ctg att gaa tca tta aaa aaa gag aaa 96
Thr Met Ser Gly Ala Leu Met Leu Ile Glu Ser Leu Lys Lys Glu Lys
20 25 30

gta gaa atg atc ttc ggt tat ccg ggc ggg gct gtg ctt ccg att tac 144 Val Glu Met Ile Phe Gly Tyr Pro Gly Gly Ala Val Leu Pro Ile Tyr 40 45

gat aag cta tac aat tca ggg ttg gta cat atc ctt ccc cgt cac gaa 192
Asp Lys Leu Tyr Asn Ser Gly Leu Val His Ile Leu Pro Arg His Glu
60

caa gga gca att cat gca gcg gag gga tac gca agg gtc tcc gga aaa 240 Gln Gly Ala Ile His Ala Ala Glu Gly Tyr Ala Arg Val Ser Gly Lys
65 70 75 80

ccg ggt gtc gtc att gcc acg tca ggg ccg gga gcg aca aac ctt gtt 288
Pro Gly Val Val Ile Ala Thr Ser Gly Pro Gly Ala Thr Asn Leu Val
85
90
95

aca ggc ctt gct gat gcc atg att gat tca ttg ccg tta gtc gtc ttt 336
Thr Gly Leu Ala Asp Ala Met Ile Asp Ser Leu Pro Leu Val Val Phe
100 105 110

aca ggg cag gta gca acc tct gta atc ggg agc gat gca ttt cag gaa 384
Thr Gly Gln Val Ala Thr Ser Val Ile Gly Ser Asp Ala Phe Gln Glu
115 120 125

gca gac att tta ggg att acg atg cca gta aca aaa cac agc tac cag
Ala Asp Ile Leu Gly Ile Thr Met Pro Val Thr Lys His Ser Tyr Gln
130 135 140

gtt cgc cag ccg gaa gat ctg ccg cgc atc att aaa gaa gcg ttc cat
Val Arg Gln Pro Glu Asp Leu Pro Arg Ile Ile Lys Glu Ala Phe His
145 150 155 160

att gca aca act gga aga ccc gga cct gta ttg att gat att ccg aaa 528

Ile Ala Thr Thr Gly Arg Pro Gly Pro Val Leu Ile Asp Ile Pro Lys
165 170 175

gat Asp	gta Val	gca Ala	aca Thr 180	att Ile	gaa Glu	gga Gly	gaa Glu	ttc Phe 185	agc Ser	tac Tyr	gat Asp	cat His	gag Glu 190	atg Met	aat Asn	576
ctc Leu	ccg Pro	gga Gly 195	tac Tyr	cag Gln	ccg Pro	aca Thr	aca Thr 200	gag Glu	ccg Pro	aat Asn	tat Tyr	ttg Leu 205	cag Gln	atc Ile	cgc Arg	624
aag Lys	ctt Leu 210	gtg Val	gaa Glu	gcc Ala	gtg Val	agc Ser 215	agt Ser	gcg Ala	aaa Lys	aaa Lys	ccg Pro 220	gtg Val	atc Ile	ctg Leu	gcg Ala	672
ggt Gly 225	gcg Ala	ggc Gly	gta Val	ctg Leu	cac His 230	gga Gly	aaa Lys	gcg Ala	tca Ser	gaa Glu 235	gaa Glu	tta Leu	aaa Lys	aat Asn	tat Tyr 240	720
gct Ala	gaa Glu	cag Gln	cag Gln	caa Gln 245	atc Ile	cct Pro	gtg Val	gca Ala	cac His 250	acc Thr	ctt Leu	ttg Leu	Gly	ctc Leu 255	gga Gly	768
ggc	ttc Phe	ccg Pro	gct Ala 260	gac Asp	cat His	ccg Pro	ctt Leu	ttc Phe 265	cta Leu	ggg Gly	atg Met	gcg Ala	gga Gly 270	atg Met	cac His	816
ggt Gly	act Thr	tat Tyr 275	aca Thr	gcc Ala	aat Asn	atg Met	gcc Ala 280	ctt Leu	cat His	gaa Glu	tgt Cys	gat Asp 285	cta Leu	tta Leu	atc Ile	864
agt Ser	atc Ile 290	ggc Gly	gcc Ala	cgt Arg	ttt Phe	gat Asp 295	gac Asp	cgt Arg	gtc Val	aca Thr	gga Gly 300	aac Asn	ctg Leu	aaa Lys	cac His	912
ttt Phe 305	gcc Ala	aga Arg	aac Asn	gca Ala	aag Lys 310	ata Ile	gcc Ala	cac His	atc Ile	gat Asp 315	att Ile	gat Asp	cca Pro	gct Ala	gaa Glu 320	960
		aaa Lys														1008
att Ile	gtc Val	ctg Leu	cag Gln 340	gag Glu	ctg Leu	atc Ile	aaa Lys	caa Gln 345	gac Asp	ggc	aaa Lys	caa Gln	agc Ser 350	gat Asp	tca Ser	1056
agc Ser	gaa Glu	tgg Trp 355	aaa Lys	aaa Lys	cag Gln	ctc Leu	gca Ala 360	gaa Glu	tgg Trp	aaa Lys	gaa Glu	gag Glu 365	tat Tyr	ccg Pro	ctc Leu	1104
tgg Trp	tat Tyr 370	Val	gat Asp	aat Asn	gaa Glu	gaa Glu 375	. Glu	ggt Gly	ttt Phe	aaa Lys	cct Pro	Gln	aaa Lys	ttg Leu	att Ile	1152
gaa Glu 385	Tyr	att Ile	cat His	caa Gln	ttt Phe 390	Thr	aaa Lys	gga Gly	gag Glu	gcc Ala 395	ı Ile	gto Val	gca Ala	acg Thr	gat Asp 400	1200
gta Val	. ggc	cag Gln	cat His	caa Gln 405	. Met	tgg Trp	tca Ser	gcg Ala	caa Gln 410	. Phe	tat Tyr	ccg Pro	tto Phe	caa Glr 415	aaa Lys	1248

gca Ala	gat Asp	aaa Lys	tgg Trp 420	gtc Val	acg Thr	tca Ser	ggc Gly	gga Gly 425	ctt Leu	gga Gly	acg Thr	atg Met	gga Gly 430	ttc Phe	ggt Gly	1296
ctt Leu	ccg Pro	gcg Ala 435	gcg Ala	atc Ile	ggc Gly	gca Ala	cag Gln 440	ctg Leu	gcc Ala	gaa Glu	aaa Lys	gat Asp 445	gct Ala	act Thr	gtt Val	1344
gtc Val	gcg Ala 450	gtt Val	gtc Val	gga Gly	gac Asp	ggc Gly 455	gga Gly	ttc Phe	caa Gln	atg Met	acg Thr 460	ctt Leu	caa Gln	gaa Glu	ctc Leu	1392
gat Asp 465	gtt Val	att Ile	cgc Arg	gaa Glu	tta Leu 470	aat Asn	ctt Leu	ccg Pro	gtc Val	aag Lys 475	gta Val	gtg Val	att Ile	tta Leu	aat Asn 480	1440
aac Asn	gct Ala	tgt Cys	ctc Leu	gga Gly 485	atg Met	gtc Val	aga Arg	cag Gln	tgg Trp 490	cag Gln	gaa Glu	att Ile	ttc Phe	tat Tyr 495	gaa Glu	1488
gaa Glu	cgt Arg	tat Tyr	tca Ser 500	gaa Glu	tct Ser	aaa Lys	ttc Phe	gct Ala 505	tct Ser	cag Gln	cct Pro	gac Asp	ttc Phe 510	gtc Val	aaa Lys	1536
ttg Leu	tcc Ser	gaa Glu 515	gca Ala	tac Tyr	ggc	att Ile	aaa Lys 520	ggc	atc Ile	aga Arg	att Ile	tca Ser 525	tca Ser	gaa Glu	gcg Ala	1584
gaa Glu	gca Ala 530	aag Lys	gaa Glu	aag Lys	ctg Leu	gaa Glu 535	gag Glu	gca Ala	tta Leu	aca Thr	tca Ser 540	aga Arg	gaa Glu	cct Pro	gtt Val	1632
gtc Val 545	Ile	gac Asp	gtg Val	cgg Arg	gtt Val 550	gcc Ala	agc Ser	gaa Glu	gaa Glu	aaa Lys 555	gta Val	ttc Phe	ccg Pro	atg Met	gtg Val 560	1680
gct Ala	ccg Pro	Gly	aaa Lys	ggg Gly 565	ctg Leu	cat His	gaa Glu	atg Met	gtg Val 570	Gly	gtg Val	aaa Lys	cct Pro	tga		1725

<210> 32

<211> 574

<212> PRT

<213> Bacillus subtilis

<400> 32

Met Gly Thr Asn Val Gln Val Asp Ser Ala Ser Ala Glu Cys Thr Gln $1 \hspace{1cm} 5 \hspace{1cm} 10 \hspace{1cm} 15$

Thr Met Ser Gly Ala Leu Met Leu Ile Glu Ser Leu Lys Lys Glu Lys 20 25 30

Val Glu Met Ile Phe Gly Tyr Pro Gly Gly Ala Val Leu Pro Ile Tyr 35 40 45

Asp Lys Leu Tyr Asn Ser Gly Leu Val His Ile Leu Pro Arg His Glu 50 55 60

Gln Gly Ala Ile His Ala Ala Glu Gly Tyr Ala Arg Val Ser Gly Lys 65 70 75 80

Pro Gly Val Val Ile Ala Thr Ser Gly Pro Gly Ala Thr Asn Leu Val 85 90 95

Thr Gly Leu Ala Asp Ala Met Ile Asp Ser Leu Pro Leu Val Val Phe 100 105 110

Thr Gly Gln Val Ala Thr Ser Val Ile Gly Ser Asp Ala Phe Gln Glu 115 120 125

Ala Asp Ile Leu Gly Ile Thr Met Pro Val Thr Lys His Ser Tyr Gln 130 135 140

Val Arg Gln Pro Glu Asp Leu Pro Arg Ile Ile Lys Glu Ala Phe His 145 150 155 160

Ile Ala Thr Thr Gly Arg Pro Gly Pro Val Leu Ile Asp Ile Pro Lys 165 170 175

Asp Val Ala Thr Ile Glu Gly Glu Phe Ser Tyr Asp His Glu Met Asn 180 185 190

Leu Pro Gly Tyr Gln Pro Thr Thr Glu Pro Asn Tyr Leu Gln Ile Arg 195 200 205

Lys Leu Val Glu Ala Val Ser Ser Ala Lys Lys Pro Val Ile Leu Ala 210 215 220

Gly Ala Gly Val Leu His Gly Lys Ala Ser Glu Glu Leu Lys Asn Tyr 225 230 235 240

Ala Glu Gln Gln Ile Pro Val Ala His Thr Leu Leu Gly Leu Gly 245 250 255

Gly Phe Pro Ala Asp His Pro Leu Phe Leu Gly Met Ala Gly Met His 260 265 270

Gly Thr Tyr Thr Ala Asn Met Ala Leu His Glu Cys Asp Leu Leu Ile 275 280 285

Ser Ile Gly Ala Arg Phe Asp Asp Arg Val Thr Gly Asn Leu Lys His 290 295 300

Phe Ala Arg Asn Ala Lys Ile Ala His Ile Asp Ile Asp Pro Ala Glu 305 310 315 320

Ile Gly Lys Ile Met Lys Thr Gln Ile Pro Val Val Gly Asp Ser Lys 325 330 335

Ile Val Leu Gln Glu Leu Ile Lys Gln Asp Gly Lys Gln Ser Asp Ser 340 345 350

Ser Glu Trp Lys Lys Gln Leu Ala Glu Trp Lys Glu Glu Tyr Pro Leu 355 360 365

Trp Tyr Val Asp Asn Glu Glu Glu Gly Phe Lys Pro Gln Lys Leu Ile 370 375 380

Glu Tyr Ile His Gln Phe Thr Lys Gly Glu Ala Ile Val Ala Thr Asp 385 390 395 400 Val Gly Gln His Gln Met Trp Ser Ala Gln Phe Tyr Pro Phe Gln Lys 405 410 415

Ala Asp Lys Trp Val Thr Ser Gly Gly Leu Gly Thr Met Gly Phe Gly 420 425 430

Leu Pro Ala Ala Ile Gly Ala Gln Leu Ala Glu Lys Asp Ala Thr Val 435 440 445

Val Ala Val Val Gly Asp Gly Gly Phe Gln Met Thr Leu Gln Glu Leu 450 455 460

Asp Val Ile Arg Glu Leu Asn Leu Pro Val Lys Val Val Ile Leu Asn 465 470 475 480

Asn Ala Cys Leu Gly Met Val Arg Gln Trp Gln Glu Ile Phe Tyr Glu 485 490 495

Glu Arg Tyr Ser Glu Ser Lys Phe Ala Ser Gln Pro Asp Phe Val Lys 500 505 510

Leu Ser Glu Ala Tyr Gly Ile Lys Gly Ile Arg Ile Ser Ser Glu Ala 515 520 525

Glu Ala Lys Glu Lys Leu Glu Glu Ala Leu Thr Ser Arg Glu Pro Val 530 540

Val Ile Asp Val Arg Val Ala Ser Glu Glu Lys Val Phe Pro Met Val 545 550 555 560

Ala Pro Gly Lys Gly Leu His Glu Met Val Gly Val Lys Pro 565 570

<210> 33

<211> 525

<212> DNA

<213> Bacillus subtilis

<220>

<221> CDS

<222> (1)..(522)

<400> 33

ttg aaa aga att atc aca ttg act gtg gtg aac cgc tcc ggg gtg tta 48
Met Lys Arg Ile Ile Thr Leu Thr Val Val Asn Arg Ser Gly Val Leu

1 5 10 15

aac cgg atc acc ggt cta ttc aca aaa agg cat tac aac att gaa agc 96
Asn Arg Ile Thr Gly Leu Phe Thr Lys Arg His Tyr Asn Ile Glu Ser
20 25 30

att aca gtt gga cac aca gaa aca gcc ggc gtt tcc aga atc acc ttc 144

Ile Thr Val Gly His Thr Glu Thr Ala Gly Val Ser Arg Ile Thr Phe

35 40 45

gtc gtt cat gtt gaa ggt gaa aat gat gtt gaa cag tta acg aaa cag 192 Val Val His Val Glu Gly Glu Asn Asp Val Glu Gln Leu Thr Lys Gln 50 55 60

ctc aac aaa cag att gat gtg ctg aaa gtc aca gac atc aca aat caa 240

Leu 65	Asn	Lys	Gln	Ile	Asp 70	Val	Leu	Lys	Val	Thr 75	Asp	Ile	Thr	Asn	Gln 80	
tcg Ser	att Ile	gtc Val	cag Gln	agg Arg 85	gag Glu	ctg Leu	gcc Ala	tta Leu	atc Ile 90	aag Lys	gtt Val	gtc Val	tcc Ser	gca Ala 95	cct Pro	288
tca Ser	aca Thr	aga Arg	aca Thr 100	gag Glu	att Ile	aat Asn	gga Gly	atc Ile 105	ata Ile	gaa Glu	ccg Pro	ttt Phe	aga Arg 110	gcc Ala	tct Ser	336
gtc Val	gtt Val	gat Asp 115	gtc Val	agc Ser	aga Arg	gac Asp	agc Ser 120	atc Ile	gtt Val	gtt Val	cag Gln	gtg Val 125	aca Thr	ggt Gly	gaa Glu	384
tct Ser	aac Asn 130	aaa Lys	att Ile	gaa Glu	gcg Ala	ctt Leu 135	att Ile	gag Glu	tta Leu	tta Leu	aaa Lys 140	cct Pro	tat Tyr	ggc Gly	att Ile	432
aaa Lys 145	gaa Glu	atc Ile	gcg Ala	aga Arg	aca Thr 150	ggt Gly	aca Thr	acg Thr	gct Ala	ttt Phe 155	gcg Ala	agg Arg	gga Gly	acc Thr	agc Ser 160	480
aaa Lys	agg Arg	cgt Arg	cat His	cca Pro 165	ata Ile	aaa Lys	caa Gln	tat Tyr	cta Leu 170	ttg Leu	tat Tyr	aaa Lys	aca Thr	taa		525
<21	0> 3	4														
<21 <21 <21		RT acil	lus	subt	ilis											
<21 <21 <21	2> P 3> B 0> 3	RT acil 4			Thr	Leu	Thr	Val	Val		Arg	· Ser	Gly	Val	Leu	
<21 <21 <21 <40 Met	2> P 3> B 0> 3 Lys	RT acil 4 Arg	·Ile	· Ile 5 · Gly	Thr				10 Arg					15 Glu	Leu Ser	
<21 <21 <21 <40 Met 1	2> P 3> B 0> 3 Lys	RT acil 4 Arg	Thr 20	: Ile 5 : Gly	Thr Leu	Phe	Thr	Lys 25	10 Arg	His	Tyr	· Asn	Ile 30	15 Glu		
<21 <21 <40 Met 1 Asr	2> P 3> B 0> 3 Lys Arg	RT acil 4 Arg Ile Val 35	Thr 20	e Ile 5 Gly His	Thr Leu	Phe Glu	Thr Thr 40	Lys 25 Ala	10 Arg	His Val	Tyr Ser	Asn Arg 45	Ile 30	Is Glu	Ser	
<21 <21 <40 Met 1 Asr	2> P 3> B 0> 3 Lys Arg Thr Val Val	RT acil 4 Arg Ile Val 35 His	Thr 20 Gly	e Ile 5 Gly His	Thr Leu Thr	Phe Glu Glu 55 Val	Thr Thr 40	Lys 25 Ala Asp	Arg Gly	His Val	Tyr Ser Glr 60	Asn Arg 45	Ile 30 Ile Thr	Glu Thr	Ser Phe	
<21 <21 <40 Met 1 Asr Ile Val	2> P 3> B 0> 3 Lys Arg Thr Val 50 1 Asr	RT acil 4 Arg Val 35 His	Thr 20 Gly	Gly His	Thr Leu Thr Gly Asp 70	Phe Glu Glu 55 Val	Thr 40 Asr	Lys 25 Ala Asp	Arg Gly Val	His Val	Ser Glr 60	Asn Arg 45 Leu D Ile	Ile 30 Ile Thr	Glu Thr Lys	Ser Phe Gln Gln 80	
<21 <21 <40 Met 1 Asr Val Let 6! Se:	2> P 3> B 0> 3 Lys Arg Thr 1 Val 50 1 Asr	RT acil 4 Arg Ile Val 35 His	Thr 20 Gly Val	Gly His Glu Arg 85	Thr Leu Thr Gly 70 Glu Glu	Phe Glu 55 Val	Thr 40 Asr	Lys 25 Ala Asp Lys	Arg Arg Val Val	His Val	Ser Glr 60	Asn Arg 45 Leu D Ile	Ile 30 Ile Thr	Glu Thr Lys Asr Ala 9!	Ser Phe Gln Gln 80	
<21 <21 <40 Met 1 Asr Val Let 6: Se:	2> P 3> B 0> 3 Lys Arg Arg Arg I Val 50 I Asr II Thi	RT acil 4 Arg Ile Val His	Thr 20 Gly Glr Glr Thr 100 Value Value	Gly His Glu Arg 85	Thr Leu Thr Asp 70 Glu	Phe Glu 55 Val Leu Asr	Thr 40 Asn Leu Ala	Lys 25 Ala Asp Let 109	Arg Arg Val Val	His Val	Tyr Ser Glr 60 Asr	Asn Arg 45 Leu D Ile	Ile 30 Ile Thr E Thr Ser 110	Glu Thr Lys Asr Als 99	Ser Phe Gln R0 Pro	

Lys Glu Ile Ala Arg Thr Gly Thr Thr Ala Phe Ala Arg Gly Thr Ser 145 150 155 160

Lys Arg Arg His Pro Ile Lys Gln Tyr Leu Leu Tyr Lys Thr 165 170

<210> 35 <211> 1029

<212> DNA

<213> Bacillus subtilis

<220>

<221> CDS

<222> (1)..(1026)

<400> 35

atg gta aaa gta tat tat aac ggt gat atc aaa gag aac gta ttg gct 48
Met Val Lys Val Tyr Tyr Asn Gly Asp Ile Lys Glu Asn Val Leu Ala

1 5 10 15

gga aaa aca gta gcg gtt atc ggg tac ggt tcg caa ggc cac gca cat 96 Gly Lys Thr Val Ala Val Ile Gly Tyr Gly Ser Gln Gly His Ala His 20 25 30

gcc ctg aac ctt aaa gaa agc gga gta gac gtg atc gtc ggt gtt aga 144
Ala Leu Asn Leu Lys Glu Ser Gly Val Asp Val Ile Val Gly Val Arg
40 45

caa gga aaa tct ttc act caa gcc caa gaa gac gga cat aaa gta ttt 192 Gln Gly Lys Ser Phe Thr Gln Ala Gln Glu Asp Gly His Lys Val Phe 50 55 60

tca gta aaa gaa gcg gca gcc caa gcc gaa atc atc atg gtt ctg ctt 240 Ser Val Lys Glu Ala Ala Ala Gln Ala Glu Ile Ile Met Val Leu Leu 65 70 75 80

ccg gat gag cag cag caa aaa gta tac gaa gct gaa atc aaa gat gaa 288
Pro Asp Glu Gln Gln Lys Val Tyr Glu Ala Glu Ile Lys Asp Glu
85 90 95

ttg aca gca gga aaa tca tta gta ttc gct cat gga ttt aac gtg cat
Leu Thr Ala Gly Lys Ser Leu Val Phe Ala His Gly Phe Asn Val His
100 105 110

ttc cat caa att gtt cct ccg gcg gat gta gat gta ttc tta gtg gcc 384
Phe His Gln Ile Val Pro Pro Ala Asp Val Asp Val Phe Leu Val Ala
115 120 125

cct aaa ggc ccg gga cac ttg gta aga aga aca tat gag caa gga gct 432
Pro Lys Gly Pro Gly His Leu Val Arg Arg Thr Tyr Glu Gln Gly Ala
130 135 140

ggc gta cct gca ttg ttc gca atc tat caa gat gtg act gga gaa gca 480 Gly Val Pro Ala Leu Phe Ala Ile Tyr Gln Asp Val Thr Gly Glu Ala 145 150 155 160

aga gac aaa gcc ctc gct tat gct aaa gga atc ggc ggc gca aga gcg 528 Arg Asp Lys Ala Leu Ala Tyr Ala Lys Gly Ile Gly Gly Ala Arg Ala 165 170 175 BGI-141CP - 40 -

	gta Val		-	_				_							576
	gag Glu														624
	ttt Phe 210														672
	gag Glu	~				_			_	-		_		_	720
_	gga Gly		_		_	_								_	768
	gat Asp								-	-					816
	atg Met		-	_			_						~		864
	tgg Trp 290		-	_			_	-		_			_		912
	gca Ala	_	-					_	-			-	_		960
_	gaa Glu	_	_	_					_	_	_	-			1008
-	tcc Ser	-													1029
<21 <21	0> 3 1> 3 2> P 3> B	42 RT	lus :	subt	ilis										

<400> 36

Met Val Lys Val Tyr Tyr Asn Gly Asp Ile Lys Glu Asn Val Leu Ala 1 5 10 15

Ala Leu Asn Leu Lys Glu Ser Gly Val Asp Val Ile Val Gly Val Arg 35 40 45

Gln Gly Lys Ser Phe Thr Gln Ala Gln Glu Asp Gly His Lys Val Phe

	50	ı				55					60				
Ser 65	Val	Lys	Glu	Ala	Ala 70	Ala	Gln	Ala	Glu	Ile 75	Ile	Met	Val	Leu	Leu 80
Pro	Asp	Glu	Gln	Gln 85	Gln	Lys	Val	Tyr	Glu 90	Ala	Glu	Ile	Lys	Asp 95	Glu
Leu	Thr	Ala	Gly 100	Lys	Ser	Leu	Val	Phe 105	Ala	His	Gly	Phe	Asn 110	Val	His
Phe	His	Gln 115	Ile	Val	Pro	Pro	Ala 120	Asp	Val	Asp	Val	Phe 125	Leu	Val	Ala
Pro	Lys 130	Gly	Pro	Gly	His	Leu 135	Val	Arg	Arg	Thr	Tyr 140	Glu	Gln	Gly	Ala
Gly 145	Val	Pro	Ala	Leu	Phe 150	Ala	Ile	Tyr	Gln	Asp 155	Val	Thr	Gly	Glu	Ala 160
Arg	Asp	Lys	Ala	Leu 165	Ala	Tyr	Ala	Lys	Gly 170	Ile	Gly	Gly	Ala	Arg 175	Ala
Gly	Val	Leu	Glu 180	Thr	Thr	Phe	Lys	Glu 185	Glu	Thr	Glu	Thr	Asp 190	Leu	Phe
Gly	Glu	Gln 195	Ala	Val	Leu	Суѕ	Gly 200	Gly	Leu	Ser	Ala	Leu 205	Val	Lys	Ala
Gly	Phe 210	Glu	Thr	Leu	Thr	Glu 215	Ala	Gly	Tyr	Gln	Pro 220	Glu	Leu	Ala	Tyr
Phe 225	Glu	Cys	Leu	His	Glu 230	Leu	Lys	Leu	Ile	Val 235	Asp	Leu	Met	Tyr	Glu 240
Glu	Gly	Leu	Ala	Gly 245	Met	Arg	Tyr	Ser	Ile 250	Ser	Asp	Thr	Ala	Gln 255	Trp
Gly	Asp	Phe	Val 260	Ser	Gly	Pro	Arg	Val 265	Val	Asp	Ala	Lys	Val 270	Lys	Glu
Ser	Met	Lys 275		Val	Leu		Asp 280		Gln	Asn	Gly	Thr 285	Phe	Ala	Lys
Glu	Trp 290	Ile	Val	Glu	Asn	Gln 295	Val	Asn	Arg	Pro	Arg 300	Phe	Asn	Ala	Ile
Asn 305	Ala	Ser	Glu	Asn	Glu 310	His	Gln	Ile	Glu	Val 315	Val	Gly	Arg	Lys	Leu 320
Arg	Glu	Met	Met	Pro 325	Phe	Val	Lys	Gln	Gly 330	Lys	Lys	Lys	Glu	Ala 335	Val

Val Ser Val Ala Gln Asn 340

<210> 37 <211> 1674

<212> DNA <213> Bacillus subtilis

<220> <221> CDS <222> (1)..(1674) <400> 37 48 atg gca gaa tta cgc agt aat atg atc aca caa gga atc gat aga gct Met Ala Glu Leu Arg Ser Asn Met Ile Thr Gln Gly Ile Asp Arg Ala ccg cac cgc agt ttg ctt cgt gca gcg gta aaa gaa gag gat ttc 96 Pro His Arg Ser Leu Leu Arg Ala Ala Gly Val Lys Glu Glu Asp Phe 20 ggc aag ccg ttt att gcg gtg tgt aat tca tac att gat atc gtt ccc 144 Gly Lys Pro Phe Ile Ala Val Cys Asn Ser Tyr Ile Asp Ile Val Pro ggt cat gtt cac ttg cag gag ttt ggg aaa atc gta aaa gaa gca atc 192 Gly His Val His Leu Gln Glu Phe Gly Lys Ile Val Lys Glu Ala Ile aga gaa gca ggg ggc gtt ccg ttt gaa ttt aat acc att ggg gta gat 240 Arg Glu Ala Gly Gly Val Pro Phe Glu Phe Asn Thr Ile Gly Val Asp 65 gat ggc atc gca atg ggg cat atc ggt atg aga tat tcg ctg cca agc 288 Asp Gly Ile Ala Met Gly His Ile Gly Met Arg Tyr Ser Leu Pro Ser cgt gaa att atc gca gac tct gtg gaa acg gtt gta tcc gca cac tgg 336 Arg Glu Ile Ile Ala Asp Ser Val Glu Thr Val Val Ser Ala His Trp ttt gac gga atg gtc tgt att ccg aac tgc gac aaa atc aca ccg gga 384 Phe Asp Gly Met Val Cys Ile Pro Asn Cys Asp Lys Ile Thr Pro Gly 115 120 atg ctt atg gcg gca atg cgc atc aac att ccg acg att ttt gtc agc 432 Met Leu Met Ala Ala Met Arg Ile Asn Ile Pro Thr Ile Phe Val Ser 130 135 ggc gga ccg atg gcg gca gga aga aca agt tac ggg cga aaa atc tcc 480 Gly Gly Pro Met Ala Ala Gly Arg Thr Ser Tyr Gly Arg Lys Ile Ser 145 150 ctt tcc tca gta ttc gaa ggg gta ggc gcc tac caa gca ggg aaa atc 528 Leu Ser Ser Val Phe Glu Gly Val Gly Ala Tyr Gln Ala Gly Lys Ile 170 175 165 aac gaa aac gag ctt caa gaa cta gag cag ttc gga tgc cca acg tgc 576 Asn Glu Asn Glu Leu Gln Glu Leu Gln Phe Gly Cys Pro Thr Cys 180 185 ggg tct tgc tca ggc atg ttt acg gcg aac tca atg aac tgt ctg tca 624 Gly Ser Cys Ser Gly Met Phe Thr Ala Asn Ser Met Asn Cys Leu Ser 195 200 205 gaa gca ctt ggt ctt gct ttg ccg ggt aat gga acc att ctg gca aca 672 Glu Ala Leu Gly Leu Ala Leu Pro Gly Asn Gly Thr Ile Leu Ala Thr 215

	ccg Pro	_	_												720
	acg Thr														768
	att Ile														816
	acc Thr														864
	tct Ser 290														912
_	aag Lys	_												_	960
	ggc Gly													_	1008
	ctt Leu													_	1056
	att Ile	_		_	_	-	_	_		_		_		_	1104
~	caa Gln 370				_	_				_	-				1152
	gct Ala														1200
	aca Thr														1248
	ctt Leu	_							-		-		_		1296
	atc Ile														1344
_	gcg Ala 450						-		_						1392

gca ttg att acg gac gga cgt ttt tcc gga gcc tcc cgt ggc ctc tca Ala Leu Ile Thr Asp Gly Arg Phe Ser Gly Ala Ser Arg Gly Leu Ser 465 470 480	1440
atc ggc cac gta tca cct gag gcc gct gag ggc ggg ccg ctt gcc ttt Ile Gly His Val Ser Pro Glu Ala Ala Glu Gly Gly Pro Leu Ala Phe 485 490 495	1488
gtt gaa aac gga gac cat att atc gtt gat att gaa aaa cgc atc ttg Val Glu Asn Gly Asp His Ile Ile Val Asp Ile Glu Lys Arg Ile Leu 500 505 510	1536
gat gta caa gtg cca gaa gaa gag tgg gaa aaa cga aaa gcg aac tgg Asp Val Gln Val Pro Glu Glu Glu Trp Glu Lys Arg Lys Ala Asn Trp 515 520 525	1584
aaa ggt ttt gaa ccg aaa gtg aaa acc ggc tac ctg gca cgt tat tct Lys Gly Phe Glu Pro Lys Val Lys Thr Gly Tyr Leu Ala Arg Tyr Ser 530 540	1632
aaa ctt gtg aca agt gcc aac acc ggc ggt att atg aaa atc Lys Leu Val Thr Ser Ala Asn Thr Gly Gly Ile Met Lys Ile 545 550 555	1674
<210> 38 <211> 558 <212> PRT <213> Bacillus subtilis	
<400> 38	
Met Ala Glu Leu Arg Ser Asn Met Ile Thr Gln Gly Ile Asp Arg Ala 1 5 10 15	V
Met Ala Glu Leu Arg Ser Asn Met Ile Thr Gln Gly Ile Asp Arg Ala	,
Met Ala Glu Leu Arg Ser Asn Met Ile Thr Gln Gly Ile Asp Arg Ala 1 5 10 15 Pro His Arg Ser Leu Leu Arg Ala Ala Gly Val Lys Glu Glu Asp Phe	
Met Ala Glu Leu Arg Ser Asn Met Ile Thr Gln Gly Ile Asp Arg Ala 1 5 10 15 Pro His Arg Ser Leu Leu Arg Ala Ala Gly Val Lys Glu Glu Asp Phe 20 25 30 Gly Lys Pro Phe Ile Ala Val Cys Asn Ser Tyr Ile Asp Ile Val Pro	
Met Ala Glu Leu Arg Ser Asn Met Ile Thr Gln Gly Ile Asp Arg Ala 1 10 5	
Met Ala Glu Leu Arg Ser Asn Met Ile Thr Gln Gly Ile Asp Arg Ala 10 Thr Gln Gly Ile Asp Arg Ala 15 Thr Gly Ile Ala Ala Gly Ile Asp Ile Ala Ala Gly Ile Ala Ile Ala Val Cys Asn Ser Tyr Ile Asp Ile Val Pro Asp Gly His Val His Leu Gln Glu Phe Gly Lys Ile Val Lys Glu Ala Ile 50 Thr Gly Ile Ala Gly Val Pro Phe Gly Phe Asn Thr Ile Gly Val Asp	
Met Ala Glu Leu Arg Ser Asn Met Ile Thr Gln Gly Ile Asp Arg Ala 15 Pro His Arg Ser Leu Leu Arg Ala Ala Gly Val Lys Glu Glu Asp Phe 20 Gly Lys Pro Phe Ile Ala Val Cys Asn Ser Tyr Ile Asp Ile Val Pro 35 Gly His Val His Leu Gln Glu Phe Gly Lys Ile Val Lys Glu Ala Ile 50 Arg Glu Ala Gly Gly Val Pro Phe Glu Phe Asn Thr Ile Gly Val Asp 65 Asp Gly Ile Ala Met Gly His Ile Gly Met Arg Tyr Ser Leu Pro Ser	
Met Ala Glu Leu Arg Ser Asn Met Ile Thr Gln Gly Ile Asp Arg Ala 15 Pro His Arg Ser Leu Leu Arg Ala Ala Gly Val Lys Glu Glu Asp Phe 20 Gly Lys Pro Phe Ile Ala Val Cys Asn Ser Tyr Ile Asp Ile Val Pro 45 Gly His Val His Leu Gln Glu Phe Gly Lys Ile Val Lys Glu Ala Ile 50 Arg Glu Ala Gly Gly Val Pro Phe Glu Phe Asn Thr Ile Gly Val Asp 80 Asp Gly Ile Ala Met Gly His Ile Gly Met Arg Tyr Ser Leu Pro Ser 90 Arg Glu Ile Ile Ala Asp Ser Val Glu Thr Val Val Ser Ala His Trp	
Met Ala Glu Leu Arg Ser Asn Met Ile Thr Gln Gly Ile Asp Arg Ala 15 Pro His Arg Ser Leu Leu Arg Ala Ala Gly Val Lys Glu Glu Asp Phe 25 Gly Lys Pro Phe Ile Ala Val Cys Asn Ser Tyr Ile Asp Ile Val Pro 45 Gly His Val His Leu Gln Glu Phe Gly Lys Ile Val Lys Glu Ala Ile 55 Arg Glu Ala Gly Gly Val Pro Phe Glu Phe Asn Thr Ile Gly Val Asp 80 Asp Gly Ile Ala Met Gly His Ile Gly Met Arg Tyr Ser Leu Pro Ser 95 Arg Glu Ile Ile Ala Asp Ser Val Glu Thr Val Val Ser Ala His Trp 100 Phe Asp Gly Met Val Cys Ile Pro Asn Cys Asp Lys Ile Thr Pro Gly	

- Leu Ser Ser Val Phe Glu Gly Val Gly Ala Tyr Gln Ala Gly Lys Ile 165 170 175
- Asn Glu Asn Glu Leu Gln Glu Leu Glu Gln Phe Gly Cys Pro Thr Cys 180 185 190
- Gly Ser Cys Ser Gly Met Phe Thr Ala Asn Ser Met Asn Cys Leu Ser 195 200 205
- Glu Ala Leu Gly Leu Ala Leu Pro Gly Asn Gly Thr Ile Leu Ala Thr 210 215 220
- Ser Pro Glu Arg Lys Glu Phe Val Arg Lys Ser Ala Ala Gln Leu Met 225 230 235 240
- Glu Thr Ile Arg Lys Asp Ile Lys Pro Arg Asp Ile Val Thr Val Lys 245 250 255
- Ala Ile Asp Asn Ala Phe Ala Leu Asp Met Ala Leu Gly Gly Ser Thr 260 265 270
- Asn Thr Val Leu His Thr Leu Ala Leu Ala Asn Glu Ala Gly Val Glu 275 280 285
- Tyr Ser Leu Glu Arg Ile Asn Glu Val Ala Glu Arg Val Pro His Leu 290 295 300
- Ala Lys Leu Ala Pro Ala Ser Asp Val Phe Ile Glu Asp Leu His Glu 305 310 315 320
- Ala Gly Gly Val Ser Ala Ala Leu Asn Glu Leu Ser Lys Lys Glu Gly 325 330 335
- Ala Leu His Leu Asp Ala Leu Thr Val Thr Gly Lys Thr Leu Gly Glu 340 345 350
- Thr Ile Ala Gly His Glu Val Lys Asp Tyr Asp Val Ile His Pro Leu 355 360 365
- Asp Gln Pro Phe Thr Glu Lys Gly Gly Leu Ala Val Leu Phe Gly Asn 370 375 380
- Leu Ala Pro Asp Gly Ala Ile Ile Lys Thr Gly Gly Val Gln Asn Gly 385 390 395 400
- Ile Thr Arg His Glu Gly Pro Ala Val Val Phe Asp Ser Gln Asp Glu 405 410 415
- Ala Leu Asp Gly Ile Ile Asn Arg Lys Val Lys Glu Gly Asp Val Val
 420 425 430
- Ile Ile Arg Tyr Glu Gly Pro Lys Gly Gly Pro Gly Met Pro Glu Met 435 $$ 440 $$ $$ Pro Gly Met Pro Glu Met
- Leu Ala Pro Thr Ser Gln Ile Val Gly Met Gly Leu Gly Pro Lys Val 450 450 460
- Ala Leu Ile Thr Asp Gly Arg Phe Ser Gly Ala Ser Arg Gly Leu Ser 465 470 475 480
- Ile Gly His Val Ser Pro Glu Ala Ala Glu Gly Gly Pro Leu Ala Phe

495 490 485 Val Glu Asn Gly Asp His Ile Ile Val Asp Ile Glu Lys Arg Ile Leu 505 500 Asp Val Gln Val Pro Glu Glu Glu Trp Glu Lys Arg Lys Ala Asn Trp 520 515 Lys Gly Phe Glu Pro Lys Val Lys Thr Gly Tyr Leu Ala Arg Tyr Ser 535 Lys Leu Val Thr Ser Ala Asn Thr Gly Gly Ile Met Lys Ile 550 <210> 39 <211> 194 <212> DNA <213> Artificial Sequence <220> <223> Description of Artificial Sequence:promoter sequence <220> <221> -35_signal <222> (136)..(141) <220> <221> -10_signal <222> (159)..(164) <400> 39 gctattgacg acagctatgg ttcactgtcc accaaccaaa actgtgctca gtaccgccaa 60 tatttctccc ttgaggggta caaagaggtg tccctagaag agatccacgc tgtgtaaaaa 120 194 gcaaccccgc ctgt <210> 40 <211> 163 <212> DNA <213> Artificial Sequence <223> Description of Artificial Sequence:promoter sequence <220> <221> -35_signal <222> (113)..(118) <220> <221> -10_signal <222> (136)..(141) <400> 40 gcctacctag cttccaagaa agatatccta acagcacaag agcggaaaga tgttttgttc 60

tacatccaga acaacctctg ctaaaattcc tgaaaaattt tgcaaaaagt tgttgacttt 120 163 atctacaagg tgtggtataa taatcttaac aacagcagga cgc <210> 41 <211> 127 <212> DNA <213> Artificial Sequence <220> <223> Description of Artificial Sequence:promoter sequence <220> <221> -35_signal <222> (34)..(39) <220> <221> -10_signal <222> (58)..(63) <220> <221> -35_signal <222> (75)..(80) <220> <221> -10_signal <222> (98)..(103) <400> 41 gaggaatcat agaattttgt caaaataatt ttattgacaa cgtcttatta acgttgatat 60 aatttaaatt ttatttgaca aaaatgggct cgtgttgtac aataaatgta gtgaggtgga 120 127 tgcaatg <210> 42 <211> 24 <212> DNA <213> Artificial Sequence <220> <223> Description of Artificial Sequence: ribosome binding site <400> 42 24 taaacatgag gaggagaaaa catg <210> 43 <211> 28 <212> DNA <213> Artificial Sequence <220> <223> Description of Artificial Sequence: ribosome binding site <400> 43

attcgagaaa tggagagaat ataatatg	28
<210> 44 <211> 13 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 44 agaaaggagg tga	13
<210> 45 <211> 23 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 45 ttaagaaagg aggtgannnn atg	23
<210> 46 <211> 23 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 46 ttagaaagga ggtgannnnn atg	23
<210> 47 <211> 23 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 47 agaaaggagg tgannnnnn atg	23
<210> 48 <211> 22 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	

<400> 48 agaaaggagg tgannnnna tg	22
<210> 49 <211> 25 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 49 ccctctagaa ggaggagaaa acatg	25
<210> 50 <211> 24 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 50 ccctctagag gaggagaaaa catg	24
<210> 51 <211> 23 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 51 ttagaaagga ggatttaaat atg	23
<210> 52 <211> 23 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 52 ttagaaagga ggtttaatta atg	23
<210> 53 <211> 23 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 53	

BGI-141CP - 50 -

ttagaaagga ggtgatttaa atg	23
<210> 54 <211> 23 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 54 ttagaaagga ggtgtttaaa atg	23
<210> 55 <211> 28 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 55 attcgagaaa ggaggtgaat ataatatg	28
<210> 56 <211> 27 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 56 attcgagaaa ggaggtgaat aataatg	27
<210> 57 <211> 28 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 57 attcgtagaa aggaggtgaa ttaatatg	28
<210> 58 <211> 3291 <212> DNA <213> Bacillus subtilis	
<400> 58 atggggacta atgtacaggt ggattcagca tctgccgaat gtacacagac gatgagcgga	60
gcattaatgc tgattgaatc attaaaaaaa gagaaagtag aaatgatctt cggttatccg	120

ggcggggctg tgcttccgat ttacgataag ctatacaatt cagggttggt acatatcctt 180 ccccgtcacg aacaaggagc aattcatgca gcggagggat acgcaagggt ctccggaaaa 240 cegggtgteg teattgecae gteagggeeg ggagegaeaa acettgttae aggeettget 300 gatgccatga ttgattcatt gccgttagtc gtctttacag ggcaggtagc aacctctgta 360 atcgggagcg atgcatttca ggaagcagac attttaggga ttacgatgcc agtaacaaaa 420 cacagetace aggttegeca geeggaagat etgeeggea teattaaaga agegtteeat 480 attgcaacaa ctggaagacc cggacctgta ttgattgata ttccgaaaga tgtagcaaca 540 attgaaggag aattcagcta cgatcatgag atgaatctcc cgggatacca gccgacaaca 600 gagccgaatt atttgcagat ccgcaagctt gtggaagccg tgagcagtgc gaaaaaaccg 660 gtgatcctgg cgggtgcggg cgtactgcac ggaaaagcgt cagaagaatt aaaaaattat 720 gctgaacagc agcaaatccc tgtggcacac acccttttgg ggctcggagg cttcccggct 780 gaccatccgc ttttcctagg gatggcggga atgcacggta cttatacagc caatatggcc 840 cttcatgaat gtgatctatt aatcagtatc ggcgcccgtt ttgatgaccg tgtcacagga 900 aacctgaaac actttgccag aaacgcaaag atagcccaca tcgatattga tccagctgaa 960 atcggaaaaa tcatgaaaac acagatteet gtagteggag acagcaaaat tgteetgeag 1020 gagetgatea aacaagaegg caaacaaage gatteaageg aatggaaaaa acagetegea 1080 gaatggaaag aagagtatcc gctctggtat gtagataatg aagaagaagg ttttaaacct 1140 cagaaattga ttgaatatat tcatcaattt acaaaaggag aggccattgt cgcaacggat 1200 gtaggccagc atcaaatgtg gtcagcgcaa ttttatccgt tccaaaaagc agataaatgg 1260 gtcacgtcag gcggacttgg aacgatggga ttcggtcttc cggcggcgat cggcgcacag 1320 ctggccgaaa aagatgctac tgttgtcgcg gttgtcggag acggcggatt ccaaatgacg 1380 cttcaagaac tcgatgttat tcgcgaatta aatcttccgg tcaaggtagt gattttaaat 1440 aacgettgte teggaatggt eagacagtgg eaggaaattt tetatgaaga aegttattea 1500 gaatctaaat tegettetea geetgaette gteaaattgt eegaageata eggeattaaa 1560 ggcatcagaa tttcatcaga agcggaagca aaggaaaagc tggaagaggc attaacatca 1620 agagaacctg ttgtcattga cgtgcgggtt gccagcgaag aaaaagtatt cccgatggtg 1680 gctccgggga aagggctgca tgaaatggtg ggggtgaaac cttgaaaaga attatcacat 1740 tgactgtggt gaaccgctcc ggggtgttaa accggatcac cggtctattc acaaaaaggc 1800 attacaacat tgaaagcatt acagttggac acacagaaac agccggcgtt tccagaatca 1860 ccttcgtcgt tcatgttgaa ggtgaaaatg atgttgaaca gttaacgaaa cagctcaaca 1920 aacagattga tgtgctgaaa gtcacagaca tcacaaatca atcgattgtc cagagggagc 1980 tggccttaat caaggttgtc tccgcacctt caacaagaac agagattaat ggaatcatag 2040 aaccgtttag agcctctgtc gttgatgtca gcagagacag catcgttgtt caggtgacag 2100 gtgaatctaa caaaattgaa gcgcttattg agttattaaa accttatggc attaaagaaa 2160 tcgcgagaac aggtacaacg gcttttgcga ggggaaccag caaaaggcgt catccaataa 2220 aacaatatct attgtataaa acataacaag ggagagattg aaatggtaaa agtatattat 2280 aacggtgata tcaaagagaa cgtattggct ggaaaaacag tagcggttat cgggtacggt 2340 tegeaaggee acgeacatge eetgaacett aaagaaageg gagtagaegt gategteggt 2400 gttagacaag gaaaatcttt cactcaagcc caagaagacg gacataaagt attttcagta 2460 aaagaagegg cageecaage egaaateate atggttetge tteeggatga geageageaa 2520 aaagtatacg aagctgaaat caaagatgaa ttgacagcag gaaaatcatt agtattcgct 2580 catggattta acgtgcattt ccatcaaatt gttcctccgg cggatgtaga tgtattctta 2640 gtggccccta aaggcccggg acacttggta agaagaacat atgagcaagg agctggcgta 2700 cctgcattgt tcgcaatcta tcaagatgtg actggagaag caagagacaa agccctcgct 2760 tatgctaaag gaatcggcgg cgcaagagcg ggcgtattag aaacgacatt taaagaagaa 2820 acagaaacag atttgttcgg tgagcaagca gttctttgcg gcggattaag cgcgcttgtc 2880 aaagccggat ttgaaacctt aactgaagca ggttatcagc ctgaacttgc atacttcgag 2940 tgtcttcatg agctgaaatt aatcgtagac cttatgtacg aagaaggact tgcaggaatg 3000 agatattcaa tctctgacac agcacagtgg ggagatttcg tatcaggccc tcgcgttgtg 3060 gacgccaaag taaaagaatc tatgaaagaa gtattaaaag atatccaaaa cggtacattc 3120 gcaaaagagt ggatcgtcga aaaccaagta aaccgtcctc gtttcaacgc tatcaatgca 3180 agcgagaacg aacatcaaat cgaagtagtg ggaagaaagc ttcgtgaaat gatgccgttt 3240 gtgaaacaag gcaagaagaa ggaagcggtg gtctccgttg cgcaaaatta a 3291

<210> 59

<211> 2363

<212> DNA

<213> Bacillus subtilis

<220>

<221> CDS

<222> (242)..(1072)

<220>

<221> CDS

<222> (1077)..(1934)

<221> CDS <222> (1939)..(2319)

<400> 59 ttggtacaag cccgttgatt ttggtatact tccattgggc agtatcgcct gcgaactgca 60 cctattatta aaatagatag acattgcagc agtctgcctt gatccaaaaa aggactggga 120 cagagggatg aaactcgccg aactttagaa agtgaagaat ccttctcgtt gtaacggaag 180 gttttttggc ttgcagaaga aaacggcaga tcatctcctc taaacatgag gaggagaaaa 240 c atg aaa aca aaa ctg gat ttt cta aaa atg aag gag tct gaa gaa ccg 289 Met Lys Thr Lys Leu Asp Phe Leu Lys Met Lys Glu Ser Glu Glu Pro att gtc atg ctg acc gct tat gat tat ccg gca gct aaa ctt gct gaa 337 Ile Val Met Leu Thr Ala Tyr Asp Tyr Pro Ala Ala Lys Leu Ala Glu caa gcg gga gtt gac atg att tta gtc ggt gat tca ctt gga atg gtc 385 Gln Ala Gly Val Asp Met Ile Leu Val Gly Asp Ser Leu Gly Met Val 35 gtc ctc ggc ctt gat tca act gtc ggt gtg aca gtt gcg gac atg atc 433 Val Leu Gly Leu Asp Ser Thr Val Gly Val Thr Val Ala Asp Met Ile 50 cat cat aca aaa gcc gtt aaa agg ggt gcg ccg aat acc ttt att gtg 481 His His Thr Lys Ala Val Lys Arg Gly Ala Pro Asn Thr Phe Ile Val aca gat atg ccg ttt atg tct tat cac ctg tct aag gaa gat acg ctg 529 Thr Asp Met Pro Phe Met Ser Tyr His Leu Ser Lys Glu Asp Thr Leu aaa aat gca gcg gct atc gtt cag gaa agc gga gct gac gca ctg aag 577 Lys Asn Ala Ala Ala Ile Val Gln Glu Ser Gly Ala Asp Ala Leu Lys ctt gag ggc gga gaa ggc gtg ttt gaa tcc att cgc gca ttg acg ctt 625 Leu Glu Gly Gly Glu Gly Val Phe Glu Ser Ile Arg Ala Leu Thr Leu 115 gga ggc att cca gta gtc agt cac tta ggt ttg aca ccg cag tca gtc 673 Gly Gly Ile Pro Val Val Ser His Leu Gly Leu Thr Pro Gln Ser Val 130 135 ggc gta ctg ggc ggc tat aaa gta cag ggc aaa gac gaa caa agc gcc Gly Val Leu Gly Gly Tyr Lys Val Gln Gly Lys Asp Glu Gln Ser Ala 145 150 aaa aaa tta ata gaa gac agt ata aaa tgc gaa gaa gca gga gct atg Lys Lys Leu Ile Glu Asp Ser Ile Lys Cys Glu Glu Ala Gly Ala Met 165 170 atg ctt gtg ctg gaa tgt gtg ccg gca gaa ctc aca gcc aaa att gcc 817 Met Leu Val Leu Glu Cys Val Pro Ala Glu Leu Thr Ala Lys Ile Ala 180 1.85 190

gag acg cta agc ata ccg gtc att gga atc ggg gct ggt gtg aaa gcg

865

Glu Thr Leu Ser Ile Pro Val Ile Gly Ile Gly Ala Gly Val Lys Ala 195 200 205	
gac gga caa gtt ctc gtt tat cat gat att atc ggc cac ggt gtt gag Asp Gly Gln Val Leu Val Tyr His Asp Ile Ile Gly His Gly Val Glu 210 215 220	913
aga aca cct aaa ttt gta aag caa tat acg cgc att gat gaa acc atc Arg Thr Pro Lys Phe Val Lys Gln Tyr Thr Arg Ile Asp Glu Thr Ile 225 230 235 240	
gaa aca gca atc agc gga tat gtt cag gat gta aga cat cgt gct ttc Glu Thr Ala Ile Ser Gly Tyr Val Gln Asp Val Arg His Arg Ala Phe 245 250 255	1009
cct gaa caa aag cat tcc ttt caa atg aac cag aca gtg ctt gac ggc Pro Glu Gln Lys His Ser Phe Gln Met Asn Gln Thr Val Leu Asp Gly 260 265 270	1057
ttg tac ggg gga aaa taag atg aga cag att act gat att tca cag ct Leu Tyr Gly Gly Lys Met Arg Gln Ile Thr Asp Ile Ser Gln Le 275 280 285	g 1106 eu
aaa gaa gcc ata aaa caa tac cat tca gag ggc aag tca atc gga ttt Lys Glu Ala Ile Lys Gln Tyr His Ser Glu Gly Lys Ser Ile Gly Pho 290 295 300	: 1154 e
gtt ccg acg atg ggg ttt ctg cat gag ggg cat tta acc tta gca gag Val Pro Thr Met Gly Phe Leu His Glu Gly His Leu Thr Leu Ala Asg 305 310 315	1202
aaa gca aga caa gaa aac gac gcc gtt att atg agt att ttt gtg aat Lys Ala Arg Gln Glu Asn Asp Ala Val Ile Met Ser Ile Phe Val Ass 320 325 330	വ
cct gca caa ttc ggc cct aat gaa gat ttt gaa gca tat ccg cgc ga Pro Ala Gln Phe Gly Pro Asn Glu Asp Phe Glu Ala Tyr Pro Arg As 340 345 350	t 1298 p
att gag cgg gat gca gct ctt gca gaa aac gcc gga gtc gat att ct Ile Glu Arg Asp Ala Ala Leu Ala Glu Asn Ala Gly Val Asp Ile Le 355 360 365	t 1346 u
ttt acg cca gat gct cat gat atg tat ccc ggt gaa aag aat gtc ac Phe Thr Pro Asp Ala His Asp Met Tyr Pro Gly Glu Lys Asn Val Th 370 375 380	g 1394 r
att cat gta gaa aga cgc aca gac gtg tta tgc ggg cgc tca aga ga Ile His Val Glu Arg Arg Thr Asp Val Leu Cys Gly Arg Ser Arg Gl 385 390 395	a 1442 u
gga cat ttt gac ggg gtc gcg atc gta ctg acg aag ctt ttc aat ct Gly His Phe Asp Gly Val Ala Ile Val Leu Thr Lys Leu Phe Asn Le 400 405 410	u
gtc aag ccg act cgt gcc tat ttc ggt tta aaa gat gcg cag cag gt Val Lys Pro Thr Arg Ala Tyr Phe Gly Leu Lys Asp Ala Gln Gln Va 420 425 430	a 1538 .1
gct gtt gtt gat ggg tta atc agc gac ttc ttc atg gat att gaa tt Ala Val Val Asp Gly Leu Ile Ser Asp Phe Phe Met Asp Ile Glu Le	g 1586 eu

445 440 435 gtt cct gtc gat acg gtc aga gag gaa gac ggc tta gcc aaa agc tct Val Pro Val Asp Thr Val Arg Glu Glu Asp Gly Leu Ala Lys Ser Ser 455 450 cgc aat gta tac tta aca gct gag gaa aga aaa gaa gcg cct aag ctg 1682 Arg Asn Val Tyr Leu Thr Ala Glu Glu Arg Lys Glu Ala Pro Lys Leu 465 tat cgg gcc ctt caa aca agt gcg gaa ctt gtc caa gcc ggt gaa aga 1730 Tyr Arg Ala Leu Gln Thr Ser Ala Glu Leu Val Gln Ala Gly Glu Arg 485 480 gat cct gaa gcg gtg ata aaa gct gca aaa gat atc att gaa acg act 1778 Asp Pro Glu Ala Val Ile Lys Ala Ala Lys Asp Ile Ile Glu Thr Thr 505 agc gga acc ata gac tat gta gag ctt tat tcc tat ccg gaa ctc gag 1826 Ser Gly Thr Ile Asp Tyr Val Glu Leu Tyr Ser Tyr Pro Glu Leu Glu 520 515 cct gtg aat gaa att gct gga aag atg att ctc gct gtt gca gtt gct Pro Val Asn Glu Ile Ala Gly Lys Met Ile Leu Ala Val Ala Val Ala 530 ttt tca aaa gcg cgt tta ata gat aat atc att att gat att cga gaa 1922 Phe Ser Lys Ala Arg Leu Ile Asp Asn Ile Ile Ile Asp Ile Arg Glu atg gag aga ata taat atg tat cga aca atg atg agc ggc aaa ctt cac Met Tyr Arg Thr Met Met Ser Gly Lys Leu His Met Glu Arg Ile 565 560 agg gca act gtt acg gaa gca aac ctg aac tat gtg gga agc att aca 2019 Arg Ala Thr Val Thr Glu Ala Asn Leu Asn Tyr Val Gly Ser Ile Thr 580 att gat gaa gat ctc att gat gct gtg gga atg ctt cct aat gaa aaa 2067 Ile Asp Glu Asp Leu Ile Asp Ala Val Gly Met Leu Pro Asn Glu Lys 595 gta caa att gtg aat aat aat gga gca cgt ctt gaa acg tat att 2115 Val Gln Ile Val Asn Asn Asn Gly Ala Arg Leu Glu Thr Tyr Ile 615 610 att cct ggt aaa cgg gga agc ggc gtc ata tgc tta aac ggt gca gcc 2163 Ile Pro Gly Lys Arg Gly Ser Gly Val Ile Cys Leu Asn Gly Ala Ala 630 gca cgc ctt gtg cag gaa gga gat aag gtc att att att tcc tac aaa 2211 Ala Arg Leu Val Gln Glu Gly Asp Lys Val Ile Ile Ile Ser Tyr Lys 645 atg atg tct gat caa gaa gcg gca agc cat gag ccg aaa gtg gct gtt 2259 Met Met Ser Asp Gln Glu Ala Ala Ser His Glu Pro Lys Val Ala Val 2307 ctg aat gat caa aac aaa att gaa caa atg ctg ggg aac gaa cca gcc Leu Asn Asp Gln Asn Lys Ile Glu Gln Met Leu Gly Asn Glu Pro Ala 680

cgt aca att ttg tagaagaaaa gcccccttta tcgggggttt tcttttaaga tttt 2363 Arg Thr Ile Leu 690

<210> 60

<211> 293

<212> PRT

<213> Bacillus subtilis

<400> 60

Met Ser Ile Ala Val Ser Glu Glu Glu Ala Lys Ala Val Glu Gly Leu

1 5 10 15

Asn Asp Tyr Leu Ser Val Glu Glu Val Glu Thr Ile Tyr Ile Pro Leu 20 25 30

Val Arg Leu His Leu His Val Lys Ser Ala Ala Glu Arg Asn Lys 35 40 45

His Val Asn Val Phe Leu Lys His Pro His Ser Ala Lys Ile Pro Phe 50 55 60

Ile Ile Gly Ile Ala Gly Ser Val Ala Val Gly Lys Ser Thr Thr Ala 65 70 75 80

Arg Ile Leu Gln Lys Leu Leu Ser Arg Leu Pro Asp Arg Pro Lys Val 85 90 95

Ser Leu Ile Thr Thr Asp Gly Phe Leu Phe Pro Thr Ala Glu Leu Lys
100 105 110

Lys Lys Asn Met Met Ser Arg Lys Gly Phe Pro Glu Ser Tyr Asp Val 115 120 125

Lys Ala Leu Leu Glu Phe Leu Asn Asp Leu Lys Ser Gly Lys Asp Ser 130 140

Val Lys Ala Pro Val Tyr Ser His Leu Thr Tyr Asp Arg Glu Gly 145 150 155 160

Val Phe Glu Val Val Glu Gln Ala Asp Ile Val Ile Ile Glu Gly Ile 165 170 175

Asn Val Leu Gln Ser Pro Thr Leu Glu Asp Asp Arg Glu Asn Pro Arg 180 185 190

Ile Phe Val Ser Asp Phe Phe Asp Phe Ser Ile Tyr Val Asp Ala Glu 195 200 205

Glu Ser Arg Ile Phe Thr Trp Tyr Leu Glu Arg Phe Arg Leu Leu Arg 210 215 220

Glu Thr Ala Phe Gln Asn Pro Asp Ser Tyr Phe His Lys Phe Lys Asp 225 230 235 240

Leu Ser Asp Glu Glu Ala Asp Glu Met Ala Ala Ser Ile Trp Glu Ser 245 250 255

Val Asn Arg Pro Asn Leu Tyr Glu Asn Ile Leu Pro Thr Lys Phe Arg

260 265 270

Ser Asp Leu Ile Leu Arg Lys Gly Asp Gly His Lys Val Glu Glu Val 275 280 285

Leu Val Arg Arg Val 290

<210> 61

<211> 281

<212> PRT

<213> Bacillus subtilis

<400> 61

Met Glu Gly Leu Asn Asp Tyr Leu Ser Val Glu Glu Val Glu Thr Ile 1 5 10 15

Tyr Ile Pro Leu Val Arg Leu Leu His Leu His Val Lys Ser Ala Ala 20 25 30

Glu Arg Asn Lys His Val Asn Val Phe Leu Lys His Pro His Ser Ala 35 40 45

Lys Ile Pro Phe Ile Ile Gly Ile Ala Gly Ser Val Ala Val Gly Lys 50 55 60

Ser Thr Thr Ala Arg Ile Leu Gln Lys Leu Leu Ser Arg Leu Pro Asp 65 70 75 80

Arg Pro Lys Val Ser Leu Ile Thr Thr Asp Gly Phe Leu Phe Pro Thr 85 90 95

Ala Glu Leu Lys Lys Lys Asn Met Met Ser Arg Lys Gly Phe Pro Glu 100 105 110

Ser Tyr Asp Val Lys Ala Leu Leu Glu Phe Leu Asn Asp Leu Lys Ser 115 120 125

Gly Lys Asp Ser Val Lys Ala Pro Val Tyr Ser His Leu Thr Tyr Asp 130 135 140

Ile Glu Gly Ile Asn Val Leu Gln Ser Pro Thr Leu Glu Asp Asp Arg 165 170 175

Glu Asn Pro Arg Ile Phe Val Ser Asp Phe Phe Asp Phe Ser Ile Tyr 180 185 190

Val Asp Ala Glu Glu Ser Arg Ile Phe Thr Trp Tyr Leu Glu Arg Phe 195 200 205

Arg Leu Leu Arg Glu Thr Ala Phe Gln Asn Pro Asp Ser Tyr Phe His 210 215 220

Lys Phe Lys Asp Leu Ser Asp Gln Glu Ala Asp Glu Met Ala Ala Ser 225 230 235 240

Ile Trp Glu Ser Val Asn Arg Pro Asn Leu Tyr Glu Asn Ile Leu Pro

245 250 255

Thr Lys Phe Arg Ser Asp Leu Ile Leu Arg Lys Gly Asp Gly His Lys 260 265 270

Val Glu Glu Val Leu Val Arg Arg Val 275 280

<210> 62

<211> 1092

<212> DNA

<213> Bacillus subtilis

<220>

<221> CDS

<222> (1)..(1089)

<400> 62

atg act aaa caa aca att cgc gtt gaa ttg aca tca aca aaa aaa ccg 48
Met Thr Lys Gln Thr Ile Arg Val Glu Leu Thr Ser Thr Lys Lys Pro

1 5 10 15

aaa cca gac cca aat cag ctt tcg ttc gga aga gtg ttt aca gac cac 96 Lys Pro Asp Pro Asn Gln Leu Ser Phe Gly Arg Val Phe Thr Asp His

atg ttt gta atg gac tat gcc gca gat aaa ggt tgg tac gat cca aga 144 Met Phe Val Met Asp Tyr Ala Ala Asp Lys Gly Trp Tyr Asp Pro Arg

atc att cct tat caa ccc tta tca atg gat cca act gca atg gtc tat 192
Ile Ile Pro Tyr Gln Pro Leu Ser Met Asp Pro Thr Ala Met Val Tyr
50 60

cac tac ggc caa acc gtg ttt gaa ggg tta aag gct tac gtg tca gag 240 His Tyr Gly Gln Thr Val Phe Glu Gly Leu Lys Ala Tyr Val Ser Glu 65 70 75 80

gat gac cat gtt ctg ctt ttc aga ccg gaa aaa aat atg gaa cgc ctg 288 Asp Asp His Val Leu Leu Phe Arg Pro Glu Lys Asn Met Glu Arg Leu 85 90 95

aat caa tca aac gac cgc ctc tgc atc ccg caa att gat gaa gaa cag 336 Asn Gln Ser Asn Asp Arg Leu Cys Ile Pro Gln Ile Asp Glu Glu Gln 100 105 110

gtt ctt gaa ggc tta aag cag ctt gtc gca att gat aaa gac tgg att 384 Val Leu Glu Gly Leu Lys Gln Leu Val Ala Ile Asp Lys Asp Trp Ile 115 120 125

acc gag cct ttc ctt ggt gtt gcg gca tct cat acg tat aag ctc ttg 480 Thr Glu Pro Phe Leu Gly Val Ala Ala Ser His Thr Tyr Lys Leu Leu 145 150 155 160

atc att ctt tct ccg gtc ggc tct tat tac aaa gaa ggc att aag ccg 528 Ile Ile Leu Ser Pro Val Gly Ser Tyr Tyr Lys Glu Gly Ile Lys Pro

BGI-141CP - 59 -

165	170	175
gtc aaa atc gct gtt gaa a Val Lys Ile Ala Val Glu 9 180	-	
aca gga aat gcc aaa acc g Thr Gly Asn Ala Lys Thr 2 195		
cag cag gta gcc gaa gag a Gln Gln Val Ala Glu Glu I 210	— -	
ggc att gag aag aaa tac a Gly Ile Glu Lys Lys Tyr 3 225 230		
ttc aaa atc aac ggt gaa a Phe Lys Ile Asn Gly Glu 1 245		
ctg gaa ggc att acg cgc a Leu Glu Gly Ile Thr Arg 2 260		9 99
ggc ctt caa gtt tca gaa o Gly Leu Gln Val Ser Glu 2 275		
gcc cat aaa gac ggc atc o Ala His Lys Asp Gly Ile 1 290		
gct gtt att tcc cca gtc g Ala Val Ile Ser Pro Val 0 305 310		
tcg atc aac aac ggt gaa a Ser Ile Asn Asn Gly Glu ' 325		-
acg att aca ggc att caa a Thr Ile Thr Gly Ile Gln i 340		
acg acc gaa gtc gca gcg Thr Thr Glu Val Ala Ala : 355		taa 1092
<210> 63 <211> 363 <212> PRT <213> Bacillus subtilis		
<400> 63 Met Thr Lys Gln Thr Ile 2 1 5	Arg Val Glu Leu Thr 10	Ser Thr Lys Lys Pro 15

Lys Pro Asp Pro Asn Gln Leu Ser Phe Gly Arg Val Phe Thr Asp His

20 25 30 Met Phe Val Met Asp Tyr Ala Ala Asp Lys Gly Trp Tyr Asp Pro Arg Ile Ile Pro Tyr Gln Pro Leu Ser Met Asp Pro Thr Ala Met Val Tyr His Tyr Gly Gln Thr Val Phe Glu Gly Leu Lys Ala Tyr Val Ser Glu Asp Asp His Val Leu Leu Phe Arg Pro Glu Lys Asn Met Glu Arg Leu Asn Gln Ser Asn Asp Arg Leu Cys Ile Pro Gln Ile Asp Glu Glu Gln Val Leu Glu Gly Leu Lys Gln Leu Val Ala Ile Asp Lys Asp Trp Ile Pro Asn Ala Glu Gly Thr Ser Leu Tyr Ile Arg Pro Phe Ile Ile Ala Thr Glu Pro Phe Leu Gly Val Ala Ala Ser His Thr Tyr Lys Leu Leu Ile Ile Leu Ser Pro Val Gly Ser Tyr Tyr Lys Glu Gly Ile Lys Pro Val Lys Ile Ala Val Glu Ser Glu Phe Val Arg Ala Val Lys Gly Gly Thr Gly Asn Ala Lys Thr Ala Gly Asn Tyr Ala Ser Ser Leu Lys Ala Gln Gln Val Ala Glu Glu Lys Gly Phe Ser Gln Val Leu Trp Leu Asp 215 220 Gly Ile Glu Lys Lys Tyr Ile Glu Glu Val Gly Ser Met Asn Ile Phe 230 Phe Lys Ile Asn Gly Glu Ile Val Thr Pro Met Leu Asn Gly Ser Ile 245 250 Leu Glu Gly Ile Thr Arg Asn Ser Val Ile Ala Leu Leu Lys His Trp 265 Gly Leu Gln Val Ser Glu Arg Lys Ile Ala Ile Asp Glu Val Ile Gln 280 Ala His Lys Asp Gly Ile Leu Glu Glu Ala Phe Gly Thr Gly Thr Ala Ala Val Ile Ser Pro Val Gly Glu Leu Ile Trp Gln Asp Glu Thr Leu 305 310 315 Ser Ile Asn Asn Gly Glu Thr Gly Glu Ile Ala Lys Lys Leu Tyr Asp 325 330 Thr Ile Thr Gly Ile Gln Lys Gly Ala Val Ala Asp Glu Phe Gly Trp Thr Thr Glu Val Ala Ala Leu Thr Glu Ser Lys 355 360

<210> 64 <211> 1071 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(1068) <400> 64 ttg aat aag ctt att gaa cga gaa aaa act gta tat tat aag gaa aag 48 Met Asn Lys Leu Ile Glu Arg Glu Lys Thr Val Tyr Tyr Lys Glu Lys ccc gac ccg tct tcc ttg ggg ttt gga caa tat ttt aca gat tat atg 96 Pro Asp Pro Ser Ser Leu Gly Phe Gly Gln Tyr Phe Thr Asp Tyr Met ttt gtg atg gac tac gaa gag ggg att gga tgg cat cat ccg aga att 144 Phe Val Met Asp Tyr Glu Glu Gly Ile Gly Trp His His Pro Arg Ile gcg ccg tac gca ccg ctt acg ctt gat ccg tct tca tct gtt ttt cat 192 Ala Pro Tyr Ala Pro Leu Thr Leu Asp Pro Ser Ser Val Phe His tac ggc cag gct gtt ttt gaa gga tta aaa gca tac aga aca gac gac 240 Tyr Gly Gln Ala Val Phe Glu Gly Leu Lys Ala Tyr Arg Thr Asp Asp ggc agg gtg ctg ctg ttc cgt ccg gat caa aat atc aaa cgg ctg aac 288 Gly Arg Val Leu Leu Phe Arg Pro Asp Gln Asn Ile Lys Arg Leu Asn 85 90 aga tcg tgt gag cgc atg agc atg ccc cct tta gac gaa gag ctg gtg 336 Arg Ser Cys Glu Arg Met Ser Met Pro Pro Leu Asp Glu Glu Leu Val 100 105 ctt gag gca ttg acg caa tta gtt gag ctg gag aaa gat tgg gtt cca 384 Leu Glu Ala Leu Thr Gln Leu Val Glu Leu Glu Lys Asp Trp Val Pro 115 120 aag gaa aaa gga acg tca ctg tat att cgt cct ttt gtc att gcc aca 432 Lys Glu Lys Gly Thr Ser Leu Tyr Ile Arg Pro Phe Val Ile Ala Thr 130 135 gaa ccg agt ctc ggt gtg aag gca tcc agg agc tat aca ttt atg atc 480 Glu Pro Ser Leu Gly Val Lys Ala Ser Arg Ser Tyr Thr Phe Met Ile 150 155 gtg ctt tcg cct gtc ggc tcc tat tat ggc gac gat cag ctg aag ccg 528 Val Leu Ser Pro Val Gly Ser Tyr Tyr Gly Asp Asp Gln Leu Lys Pro 165 170 175 gtt aga atc tat gtc gaa gat gag tat gtg agg gcg gtc aac gga gga 576

Val Arg Ile Tyr Val Glu Asp Glu Tyr Val Arg Ala Val Asn Gly Gly

BGI-141CP - 62 -

180		185	190	
		aac tat gcc gcc ag Asn Tyr Ala Ala Se 20	r Leu Gln Ala	524
		tat gac cag gta ct Tyr Asp Gln Val Le 220	3 33 3 3	572
		gaa gta ggg agc at Glu Val Gly Ser Me 235		720
Phe Val Ile Asn		aca cct gct tta ag Thr Pro Ala Leu Se 250	- 33	768
		gcg att gaa ttg at Ala Ile Glu Leu Il 265		316
		ata tcg att gat ga Ile Ser Ile Asp Gl 28	u Val Tyr Ala	364
		gag gtc ttt ggc ac Glu Val Phe Gly Th 300		912
		ctc aac atc cat gg Leu Asn Ile His Gl 315	2 3 3	960
		gac ctc tcg aaa aa Asp Leu Ser Lys Ly 330		1008
		aag gta aaa ggc co Lys Val Lys Gly Pr 345		1056
aca gtg gaa gtg Thr Val Glu Val 355	tga		1	L071
<210> 65 <211> 356 <212> PRT <213> Bacillus s	subtilis			
<400> 65 Met Asn Lys Leu 1	Ile Glu Arg Glu 5	Lys Thr Val Tyr Ty 10	r Lys Glu Lys 15	
Pro Asp Pro Ser 20	Ser Leu Gly Phe	Gly Gln Tyr Phe Th	r Asp Tyr Met 30	
Phe Val Met Asp 35	Tyr Glu Glu Gly 40	Ile Gly Trp His Hi	s Pro Arg Ile 5	

BGI-141CP

Ala Pro Tyr Ala Pro Leu Thr Leu Asp Pro Ser Ser Ser Val Phe His

Tyr Gly Gln Ala Val Phe Glu Gly Leu Lys Ala Tyr Arg Thr Asp Asp

Gly Arg Val Leu Leu Phe Arg Pro Asp Gln Asn Ile Lys Arg Leu Asn

Arg Ser Cys Glu Arg Met Ser Met Pro Pro Leu Asp Glu Glu Leu Val

Leu Glu Ala Leu Thr Gln Leu Val Glu Leu Glu Lys Asp Trp Val Pro 120

Lys Glu Lys Gly Thr Ser Leu Tyr Ile Arg Pro Phe Val Ile Ala Thr 135

Glu Pro Ser Leu Gly Val Lys Ala Ser Arg Ser Tyr Thr Phe Met Ile

Val Leu Ser Pro Val Gly Ser Tyr Tyr Gly Asp Asp Gln Leu Lys Pro 170

Val Arg Ile Tyr Val Glu Asp Glu Tyr Val Arg Ala Val Asn Gly Gly

Val Gly Phe Ala Lys Thr Ala Gly Asn Tyr Ala Ala Ser Leu Gln Ala

Gln Arg Lys Ala Asn Glu Leu Gly Tyr Asp Gln Val Leu Trp Leu Asp

Ala Ile Glu Lys Lys Tyr Val Glu Glu Val Gly Ser Met Asn Ile Phe

Phe Val Ile Asn Gly Glu Ala Val Thr Pro Ala Leu Ser Gly Ser Ile

Leu Ser Gly Val Thr Arg Ala Ser Ala Ile Glu Leu Ile Arg Ser Trp

Gly Ile Pro Val Arg Glu Glu Arg Ile Ser Ile Asp Glu Val Tyr Ala

Ala Ser Ala Arg Gly Glu Leu Thr Glu Val Phe Gly Thr Gly Thr Ala

Ala Val Val Thr Pro Val Gly Glu Leu Asn Ile His Gly Lys Thr Val 305

Ile Val Gly Asp Gly Gln Ile Gly Asp Leu Ser Lys Leu Tyr Glu 325

Thr Ile Thr Asp Ile Gln Leu Gly Lys Val Lys Gly Pro Phe Asn Trp 340 345

Thr Val Glu Val

355

<210> 66 <211> 1428 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(1425) <400> 66 atg tta aac ggc caa aaa gaa tat cgc gtg gaa aaa gac ttc ctt ggg 48 Met Leu Asn Gly Gln Lys Glu Tyr Arg Val Glu Lys Asp Phe Leu Gly gaa aaa caa att gaa gca gat gtt tat tac gga att cag acg ctc cgt 96 Glu Lys Gln Ile Glu Ala Asp Val Tyr Tyr Gly Ile Gln Thr Leu Arg 144 gct tct gaa aat ttt ccg atc aca gga tac aaa atc cat gag gaa atg Ala Ser Glu Asn Phe Pro Ile Thr Gly Tyr Lys Ile His Glu Glu Met att aac gca ctg gcg att gtg aaa aaa gct gcg gct ctt gcc aac atg 192 Ile Asn Ala Leu Ala Ile Val Lys Lys Ala Ala Leu Ala Asn Met 55 gac gtg aaa cgg ctg tat gaa gga att ggc caa gct atc gta caa gcc 240 Asp Val Lys Arg Leu Tyr Glu Gly Ile Gly Gln Ala Ile Val Gln Ala 65 gct gac gag att ctg gaa ggc aag tgg cac gat cag ttt atc gtc gat 288 Ala Asp Glu Ile Leu Glu Gly Lys Trp His Asp Gln Phe Ile Val Asp ccg att cag ggc ggt gcc gga act tct atg aac atg aac gcg aat gag 336 Pro Ile Gln Gly Gly Ala Gly Thr Ser Met Asn Met Asn Ala Asn Glu 100 384 gtt atc gga aac cgg gcg ctt gaa atc atg gga cat aaa aag gga gat Val Ile Gly Asn Arg Ala Leu Glu Ile Met Gly His Lys Lys Gly Asp 120 tat atc cat tta agt cca aac aca cat gtg aac atg tca cag tct cag 432 Tyr Ile His Leu Ser Pro Asn Thr His Val Asn Met Ser Gln Ser Gln 135 140 aac gat gtg ttc ccg act gct atc cat att tcc aca ttg aag ctc tta 480 Asn Asp Val Phe Pro Thr Ala Ile His Ile Ser Thr Leu Lys Leu Leu 150 155 gaa aaa ctg ctg aaa aca atg gaa gat atg cat agt gtg ttt aaa caa 528 Glu Lys Leu Leu Lys Thr Met Glu Asp Met His Ser Val Phe Lys Gln 165 170 aaa gca cag gag ttt cac tct gtt att aaa atg ggc cgg aca cac ctt 576 Lys Ala Gln Glu Phe His Ser Val Ile Lys Met Gly Arg Thr His Leu 180 185 190 caa gat gcg gtt ccg atc cgt ctt ggc cag gaa ttc gaa gct tac agc 624

Gln Asp Ala Val Pro Ile Arg Leu Gly Gln Glu Phe Glu Ala Tyr Ser

BGI-141CP - 65 -

		195					200					205				
-	_		gag Glu	_	_			-		-		-	-	_		672
			gtc Val													720
_	~		gaa Glu				_	_	-	_			-	-		768
_			cct Pro 260		-		_	-			_	_				816
		-	gcc Ala				_		_				-	-	_	864
_		_	tcg Ser	_				_	_	-		_		_		912
_	-	_	gga Gly			-			_		-	_	_	-		960
			atg Met	_			_		_		_			_		1008
			gcg Ala 340		_	_				_				_		1056
_		_	gcc Ala		_			_		_	_					1104
			ttg Leu													1152
~			gac Asp		-					_	_		_	-	_	1200
-	_		tac Tyr	_	-		_	_					-	_		1248
			ggg Gly 420													1296
			caa Gln		-		_		_	_	_		-		-	1344

1392

1428

act gaa gaa ttg gat att att tta aac cca tat gag atg acc aaa Thr Glu Glu Leu Asp Ile Ile Leu Asn Pro Tyr Glu Met Thr Lys 455 460 cca ggt atc gca ggg aaa gaa cta tta gaa aaa taa Pro Gly Ile Ala Gly Lys Glu Leu Leu Glu Lys 470 <210> 67 <211> 475 <212> PRT <213> Bacillus subtilis <400> 67 Met Leu Asn Gly Gln Lys Glu Tyr Arg Val Glu Lys Asp Phe Leu Gly Glu Lys Gln Ile Glu Ala Asp Val Tyr Tyr Gly Ile Gln Thr Leu Arg Ala Ser Glu Asn Phe Pro Ile Thr Gly Tyr Lys Ile His Glu Met Ile Asn Ala Leu Ala Ile Val Lys Lys Ala Ala Leu Ala Asn Met Asp Val Lys Arg Leu Tyr Glu Gly Ile Gly Gln Ala Ile Val Gln Ala Ala Asp Glu Ile Leu Glu Gly Lys Trp His Asp Gln Phe Ile Val Asp Pro Ile Gln Gly Gly Ala Gly Thr Ser Met Asn Met Asn Ala Asn Glu Val Ile Gly Asn Arg Ala Leu Glu Ile Met Gly His Lys Lys Gly Asp 120 Tyr Ile His Leu Ser Pro Asn Thr His Val Asn Met Ser Gln Ser Gln Asn Asp Val Phe Pro Thr Ala Ile His Ile Ser Thr Leu Lys Leu Leu 155 Glu Lys Leu Leu Lys Thr Met Glu Asp Met His Ser Val Phe Lys Gln Lys Ala Gln Glu Phe His Ser Val Ile Lys Met Gly Arg Thr His Leu Gln Asp Ala Val Pro Ile Arg Leu Gly Gln Glu Phe Glu Ala Tyr Ser Arg Val Leu Glu Arg Asp Ile Lys Arg Ile Lys Gln Ser Arg Gln His Leu Tyr Glu Val Asn Met Gly Ala Thr Ala Val Gly Thr Gly Leu Asn 225 230 235

Ala Asp Pro Glu Tyr Ile Lys Gln Val Val Lys His Leu Ala Asp Ile
245 250 255

Ser Gly Leu Pro Leu Val Gly Ala Asp His Leu Val Asp Ala Thr Gln 260 265 270

Asn Thr Asp Ala Tyr Thr Glu Val Ser Ala Ser Leu Lys Val Cys Met 275 280 285

Met Asn Met Ser Lys Ile Ala Asn Asp Leu Arg Leu Met Ala Ser Gly 290 295 300

Pro Arg Ala Gly Leu Ala Glu Ile Ser Leu Pro Ala Arg Gln Pro Gly 305 310 315 320

Ser Ser Ile Met Pro Gly Lys Val Asn Pro Val Met Ala Glu Leu Ile 325 330 335

Asn Gln Ile Ala Phe Gln Val Ile Gly Asn Asp Asn Thr Ile Cys Leu 340 345 350

Ala Ser Glu Ala Gly Gln Leu Glu Leu Asn Val Met Glu Pro Val Leu 355 360 365

Val Phe Asn Leu Gln Ser Ile Ser Ile Met Asn Asn Gly Phe Arg 370 375 380

Ser Phe Thr Asp Asn Cys Leu Lys Gly Ile Glu Ala Asn Glu Lys Arg 385 390 395 400

Met Lys Gln Tyr Val Glu Lys Ser Ala Gly Val Ile Thr Ala Val Asn \$405\$

Pro His Leu Gly Tyr Glu Ala Ala Ala Arg Ile Ala Arg Glu Ala Ile 420 425 430

Met Thr Gly Gln Ser Val Arg Asp Leu Cys Leu Gln His Asp Val Leu 435 440 445

Thr Glu Glu Glu Leu Asp Ile Ile Leu Asn Pro Tyr Glu Met Thr Lys 450 455 460

Pro Gly Ile Ala Gly Lys Glu Leu Leu Glu Lys 465 470

<210> 68

<211> 768

<212> DNA

<213> Bacillus subtilis

<220>

<221> CDS

<222> (1)..(765)

<400> 68

atg aaa cga gaa agc aac att caa gtg ctc agc cgt ggt caa aaa gat Met Lys Arg Glu Ser Asn Ile Gln Val Leu Ser Arg Gly Gln Lys Asp 1 5 10 15

cag cct gtg agc cag att tat caa gta tca aca atg act tct cta tta 96

Gln	Pro	Val	Ser 20	Gln	Ile	Tyr	Gln	Val 25	Ser	Thr	Met	Thr	Ser 30	Leu	Leu	
_											gag Glu					144
											gac Asp 60				_	192
		_		_			_		_		gac Asp					240
											tca Ser					288
	_	_	_	_				_	-		atg Met		-			336
	_					_	-	_		_	aga Arg					384
											cag Gln 140					432
											gaa Glu					480
_	_					_			_		acg Thr		_			528
											tct Ser					576
			-	_		_					cac His					624
			_	_	_	-	_				aaa Lys 220	_		_		672
	_		_				_				gcg Ala		_	_		720
	~				-		_			-	gga Gly	_		_	taa	768

<211> 255

<212> PRT

<213> Bacillus subtilis

<400> 69

Met Lys Arg Glu Ser Asn Ile Gln Val Leu Ser Arg Gly Gln Lys Asp 1 5 10 15

Gln Pro Val Ser Gln Ile Tyr Gln Val Ser Thr Met Thr Ser Leu Leu 20 25 30

Asp Gly Val Tyr Asp Gly Asp Phe Glu Leu Ser Glu Ile Pro Lys Tyr 35 40 45

Gly Asp Phe Gly Ile Gly Thr Phe Asn Lys Leu Asp Gly Glu Leu Ile 50 55 60

Gly Phe Asp Gly Glu Phe Tyr Arg Leu Arg Ser Asp Gly Thr Ala Thr 65 70 75 80

Pro Val Gln Asn Gly Asp Arg Ser Pro Phe Cys Ser Phe Thr Phe Phe
85 90 95

Thr Pro Asp Met Thr His Lys Ile Asp Ala Lys Met Thr Arg Glu Asp 100 105 110

Phe Glu Lys Glu Ile Asn Ser Met Leu Pro Ser Arg Asn Leu Phe Tyr 115 120 125

Ala Ile Arg Ile Asp Gly Leu Phe Lys Lys Val Gln Thr Arg Thr Val 130 135 140

Glu Leu Gln Glu Lys Pro Tyr Val Pro Met Val Glu Ala Val Lys Thr 145 150 155 160

Gln Pro Ile Phe Asn Phe Asp Asn Val Arg Gly Thr Ile Val Gly Phe 165 170 175

Leu Thr Pro Ala Tyr Ala Asn Gly Ile Ala Val Ser Gly Tyr His Leu 180 185 190

His Phe Ile Asp Glu Gly Arg Asn Ser Gly Gly His Val Phe Asp Tyr 195 200 205

Val Leu Glu Asp Cys Thr Val Thr Ile Ser Gln Lys Met Asn Met Asn 210 215 220

Leu Arg Leu Pro Asn Thr Ala Asp Phe Phe Asn Ala Asn Leu Asp Asn 225 230 235 240

Pro Asp Phe Ala Lys Asp Ile Glu Thr Thr Glu Gly Ser Pro Glu 245 250 255

<210> 70

<211> 1254

<212> DNA

<213> Escherichia coli

<220>

<221> CDS

BGI-141CP - 70 -

<222> (1)..(1251)

<400> 70 atg aca ttc tcc ctt ttt ggt gac aaa ttt acc cgc cac tcc ggc att 48 Met Thr Phe Ser Leu Phe Gly Asp Lys Phe Thr Arg His Ser Gly Ile acg ctg ttg atg gaa gat ctg aac gac ggt tta cgc acg cct ggc gcg Thr Leu Leu Met Glu Asp Leu Asn Asp Gly Leu Arg Thr Pro Gly Ala 25 att atg ctc ggc ggt ggt aat ccg gcg cag atc ccg gaa atg cag gac 144 Ile Met Leu Gly Gly Asn Pro Ala Gln Ile Pro Glu Met Gln Asp 40 tac ttc cag acg cta ctg acc gac atg ctg gaa agt ggc aaa gcg act 192 Tyr Phe Gln Thr Leu Leu Thr Asp Met Leu Glu Ser Gly Lys Ala Thr 55 gat gca ctg tgt aac tac gac ggt cca cag ggg aaa acg gag cta ctc 240 Asp Ala Leu Cys Asn Tyr Asp Gly Pro Gln Gly Lys Thr Glu Leu Leu aca ctg ctt gcc gga atg ctg cgc gag aag ttg ggt tgg gat atc gaa 288 Thr Leu Leu Ala Gly Met Leu Arg Glu Lys Leu Gly Trp Asp Ile Glu cca cag aat att gca cta aca aac ggc agc cag agc gcg ttt ttc tac 336 Pro Gln Asn Ile Ala Leu Thr Asn Gly Ser Gln Ser Ala Phe Phe Tyr 105 tta ttt aac ctg ttt gcc gga cgc cgt gcc gat ggt cgg gtc aaa aaa 384 Leu Phe Asn Leu Phe Ala Gly Arg Arg Ala Asp Gly Arg Val Lys Lys 115 120 gtg ctg ttc ccg ctt gca ccg gaa tac att ggc tat gct gac gca gga 432 Val Leu Phe Pro Leu Ala Pro Glu Tyr Ile Gly Tyr Ala Asp Ala Gly 130 135 140 ctg gaa gaa gat ctg ttt gtc tct gcg cgt ccg aat att gaa ctg ctg 480 Leu Glu Glu Asp Leu Phe Val Ser Ala Arg Pro Asn Ile Glu Leu Leu 150 155 ccg gaa ggc cag ttt aaa tac cac gtc gat ttt gag cat ctg cat att 528 Pro Glu Gly Gln Phe Lys Tyr His Val Asp Phe Glu His Leu His Ile 165 170 ggc gaa gaa acc ggg atg att tgc gtc tcc cgg ccg acg aat cca aca 576 Gly Glu Glu Thr Gly Met Ile Cys Val Ser Arg Pro Thr Asn Pro Thr ggc aat gtg att act gac gaa gag ttg ctg aag ctt gac gcg ctg ggc 624

aat caa cac ggc att ccg ctg gtg att gat aac gct tat ggc gtc ccg 672

Asn Gln His Gly Ile Pro Leu Val Ile Asp Asn Ala Tyr Gly Val Pro
210 215 220

ttc ccg ggt atc atc ttc agt gaa gcg cgc ccg cta tgg aat ccg aat 720

Phe Pro Gly Ile Ile Phe Ser Glu Ala Arg Pro Leu Trp Asn Pro Asn

Gly Asn Val Ile Thr Asp Glu Glu Leu Leu Lys Leu Asp Ala Leu Gly

BGI-141CP - 71 -

225	230	235	240
	Ser Leu Ser Lys Leu	ggt cta cct ggc tcc Gly Leu Pro Gly Ser 255	_
	_	atc acc gcc atc acc Ile Thr Ala Ile Thr 270	
		ggt att ggt ccg gcg Gly Ile Gly Pro Ala 285	-
		ctg cgc ctg tct gaa Leu Arg Leu Ser Glu 300	
_		cag gaa act atc gcc Gln Glu Thr Ile Ala 315	
	Pro Glu Asn Arg Cys	ctg att cat aaa ccg Leu Ile His Lys Pro 335	-
		gat ttg ccc att acg Asp Leu Pro Ile Thr 350	
		ggc gtg ctg atg gtg Gly Val Leu Met Val 365	_
		ccg tgg ccg cat acg Pro Trp Pro His Thr 380	
· · · · · · · · · · · · · · · · · · ·		ccg gag aaa att gag Pro Glu Lys Ile Glu 395	
	ı Ala Glu Glu Ile Glu	aga gcc tgg gct gaa Arg Ala Trp Ala Glu 415	
cac taa His			1254
<210> 71 <211> 417 <212> PRT <213> Escherichia d	coli		
<400> 71 Met Thr Phe Ser Let 1		Thr Arg His Ser Gly	Ile

Thr Leu Leu Met Glu Asp Leu Asn Asp Gly Leu Arg Thr Pro Gly Ala 20 25 30

Ile Met Leu Gly Gly Gly Asn Pro Ala Gln Ile Pro Glu Met Gln Asp 35 40 45

Tyr Phe Gln Thr Leu Leu Thr Asp Met Leu Glu Ser Gly Lys Ala Thr 50 55 60

Asp Ala Leu Cys Asn Tyr Asp Gly Pro Gln Gly Lys Thr Glu Leu Leu 65 70 75 80

Thr Leu Leu Ala Gly Met Leu Arg Glu Lys Leu Gly Trp Asp Ile Glu 85 90 95

Pro Gln Asn Ile Ala Leu Thr Asn Gly Ser Gln Ser Ala Phe Phe Tyr 100 105 110

Leu Phe Asn Leu Phe Ala Gly Arg Arg Ala Asp Gly Arg Val Lys Lys 115 120 125

Val Leu Phe Pro Leu Ala Pro Glu Tyr Ile Gly Tyr Ala Asp Ala Gly 130 135 140

Leu Glu Glu Asp Leu Phe Val Ser Ala Arg Pro Asn Ile Glu Leu Leu 145 5 55 160

Pro Glu Gly Gln Phe Lys Tyr His Val Asp Phe Glu His Leu His Ile 165 170 175

Gly Glu Glu Thr Gly Met Ile Cys Val Ser Arg Pro Thr Asn Pro Thr 180 185 190

Gly Asn Val Ile Thr Asp Glu Glu Leu Leu Lys Leu Asp Ala Leu Gly 195 200 205

Asn Gln His Gly Ile Pro Leu Val Ile Asp Asn Ala Tyr Gly Val Pro 210 215 220

Phe Pro Gly Ile Ile Phe Ser Glu Ala Arg Pro Leu Trp Asn Pro Asn 225 230 235 235

Ile Val Leu Cys Met Ser Leu Ser Lys Leu Gly Leu Pro Gly Ser Arg 245 250 255

Cys Gly Ile Ile Ile Ala Asn Glu Lys Ile Ile Thr Ala Ile Thr Asn 260 265 270

Met Asn Gly Ile Ile Ser Leu Ala Pro Gly Gly Ile Gly Pro Ala Met 275 280 285

Met Cys Glu Met Ile Lys Arg Asn Asp Leu Leu Arg Leu Ser Glu Thr 290 295 300

Val Ile Lys Pro Phe Tyr Tyr Gln Arg Val Gln Glu Thr Ile Ala Ile 305 310 315 320

Ile Arg Arg Tyr Leu Pro Glu Asn Arg Cys Leu Ile His Lys Pro Glu 325 330 335

Gly Ala Ile Phe Leu Trp Leu Trp Phe Lys Asp Leu Pro Ile Thr Thr 340 345 350

Lys Gln Leu Tyr Gln Arg Leu Lys Ala Arg Gly Val Leu Met Val Pro 355 360 365

Gly His Asn Phe Phe Pro Gly Leu Asp Lys Pro Trp Pro His Thr His 370 375 380

Gln Cys Met Arg Met Asn Tyr Val Pro Glu Pro Glu Lys Ile Glu Ala 385 390 395 400

Gly Val Lys Ile Leu Ala Glu Glu Ile Glu Arg Ala Trp Ala Glu Ser $405 \hspace{1.5cm} 410 \hspace{1.5cm} 415 \hspace{1.5cm}$

His

<210> 72

<211> 8803

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN294 plasmid

tgcgccgcta cagggcgcgt ccattcgcca ttcaggctgc gcaactgttg ggaagggcga 60 teggtgeggg cetetteget attacgceag etggegaaag ggggatgtge tgcaaggega 120 ttaagttggg taacgccagg gttttcccag tcacgacgtt gtaaaacgac ggccagtgaa 180 ttgtaatacg actcactata gggcgaattg ggcccgacgt cgcatgctgg atgaaaagcc 240 gatgaccgct tttcaggtct gtcagcagct ttttcctgct gtatatgaaa aggaattgtt 300 tttaacgatg tcagaaacgg caggtcacct tgatgtgttg gaggctgaag aagccatcac 360 gtcatattgg gaaggaaata ccgtatactt taaaacaatg aagaggtgaa atgggtgaaa 420 catatagcgg gaaaaaggat ttggataacc ggcgcttcag gagggcttgg agaaagaatc 480 gcatacttat gcgcggctga aggagcccat gtcctgctgt cggctagacg cgaggatcgt 540 ttgatagaaa tcaaaaggaa aataaccgag gaatggagcg gacagtgtga gatttttcct 600 ctggatgtcg gccgcctaga ggatatcgcc cgggtccgcg atcagatcgg ctcgattgat 660 gtactgatta acaatgcagg cttcggtata tttgaaacgg ttttagactc tacattggat 720 gacatgaaag cgatgtttga tgtgaatgtc ttcggcctga tcgcctgtac aaaagcggtg 780 cttccgcaaa tgcttgagca aaaaaaggga catatcatca atatcgcctc tcaagcgggg 840 aaaatcgcca caccgaagtc tagcctgtat tccgcgacca aacatgccgt gttaggttac 900 tcaaacgctt tgcggatgga gctttcggga accggcattt atgtgacaac agtcaacccg 960 ggcccgattc agacggactt tttttccatt gctgataaag gcggggacta cgccaaaaat 1020 gtcggccgct ggatgcttga tcctgatgac gtggcagctc aaattacagc tgcaattttt 1080 acgaaaaagc gggagatcaa tcttccgcgt ttaatgaatg ccggcactaa gctgtatcag 1140 ctgtttccag ctcttgtaga aaagctggca ggacgcgcgc tcatgaaaaa ataatgatag 1200 aactgcctgt ggtggagtgg cttgtttctc acggggcagt ttttgatagt ggaagggaga 1260 gattgttgaa tgtcagttca ttcagaagtc cttcatgctc tgcttaaaga tccgtttatt 1320 cagaaactga ttgatgcaga gcctgtattc tgggcaaatt caggcaagaa agaggggcca 1380 ttaccccgtg cagatgagtg ggcaaccgag atagcggaag cggaaaaaag aatgcagcgg 1440 tttgcacctt acattgccga ggtgtttcct gagacgaaag gcgctaaagg aatcatcgag 1500 teteegettt ttgaggtgca geatatgaag ggaaagetgg aageggeata teageageea 1560 tttcccggaa gatggctttt aaagtgcgac catgagcttc cgatttcagg atcgattaaa 1620 gcgaggggcg ggatttatga agtgttaaag tatgctgaaa atctcgcgct tcaagaagga 1680 atgcttcagg aaaccgatga ttaccgcatc ttacaggaag agcggtttac cgggttttc 1740 tecegetatt egattgetgt eggttegaca ggaaatetag gtttaageat eggeateate 1800 ggcgcggcac tcgggtttcg cgtgacagtg catatgtccg ccgatgctaa gcagtggaaa 1860 aaggatetee teegeeaaaa gggagteaet gttatggagt acgaaacaga ttacagtgaa 1920 gcggtgaacg aagggagacg gcaggcggaa caagatccat tctgttattt tattgatgat 1980 gaacattctc gtcagctgtt cttaggatat gctgttgctg caagccgatt aaaaacacag 2040 cttgactgta tgaatataaa gccaagtctt gagacgccct tgtttgtgta tctgccgtgc 2100 ggagtcggcg gaggaccggg cggtgtagca tttgggctga agcttttata cggagatgat 2160 gttcatgtgt ttttcgcaga accaactcat tcaccttgta tgctgttagg gctttattca 2220 qqacttcacq agaaqatctc cqtccaqqat atcqqcctgq ataatcagac ggctgctgac 2280 ggacttgccg tagggaggcc gtcaggattt gtcggcaagc tgattgaacc gcttctgagc 2340 ggctgttata cggtagagga caatacgctt tatactttgc ttcatatgct ggctgtatct 2400 gaagataaat atttagagcc ctctgctctt gctggcatgt tcgggccggt tcagcttttt 2460 tegacagaag agggaaggeg etatgeteag aaatataaga tggaacatge egtacatgte 2520 gtctggggaa cgggaggaag catggttcca aaagatgaaa tggctgcgta taaccgaatc 2580 ggtgctgatt tgctaaaaaa acgaaatgga aaataagcag acagtgaaaa ggttttccgt 2640 tacaatcttt gtaagggttt taacctacag agagtcaggt gtaaacagtg aaaaataaag 2700 aacttaacct acatacttta tatacacagc acaatcggga gtcttggtct ggttttgggg 2760 ggcatttgtc gattgctgta tctgaagaag aggcaaaagc tgtggaagga ttgaatgatt 2820 atctatctgt tgaagaagtg gagacgatct atattccgct tgttcgcttg cttcatttac 2880

BGI-141CP - 75 -

atgtcaagtc tgcggctgaa cgcaataagc atgtcaatgt ttttttgaag cacccacatt 2940 cagccaaaat tccgtttatt atcggcattg ccggcagtgt cgcagtcgga aaaagcacga 3000 eggegeggat ettgeagaag etgetttege gtttgeetga eegteeaaaa gtgageetta 3060 tcacgacaga tggtttttta tttcctactg ccgagctgaa aaagaaaaat atgatgtcaa 3120 gaaaaggatt teetgaaage tatgatgtaa aggegetget egaatttttg aatgaettaa 3180 aatcaggaaa ggacagcgta aaggccccgg tgtattccca tctaacctat gaccgcgagg 3240 aaggtgtgtt cgaggttgta gaacaggcgg atattgtgat tattgaaggc attaatgttc 3300 ttcagtcgcc caccttggag gatgaccggg aaaacccgcg tatttttgtt tccgatttct 3360 ttgatttttc gatttatgtg gatgcggagg aaagccggat tttcacttgg tatttagagc 3420 gttttcgcct gcttcgggaa acagcttttc aaaatcctga ttcatatttt cataaattta 3480 aagacttgtc cgatcaggag gctgacgaga tggcagcctc gatttgggag agtgtcaacc 3540 ggccgaattt atatgaaaat attttgccaa ctaaattcag gtcagatctc attttgcgta 3600 agggagacgg gcataaggtc gaggaagtgt tggtaaggag ggtatgaaat gtgctgcagc 3660 tegageaata gttaceetta ttateaagat aagaaagaaa aggattttte getaegetea 3720 aatcctttaa aaaaacacaa aagaccacat tttttaatgt ggtctttatt cttcaactaa 3780 ttatgttaca gtaatattga cttttaaaaa aggattgatt ctaatgaaga aagcagacaa 3900 qtaaqcctcc taaattcact ttagataaaa atttaggagg catatcaaat gaactttaat 3960 aaaattgatt tagacaattg gaagagaaaa gagatattta atcattattt gaaccaacaa 4020 acgactttta gtataaccac agaaattgat attagtgttt tataccgaaa cataaaacaa 4080 gaaggatata aattttaccc tgcatttatt ttcttagtga caagggtgat aaactcaaat 4140 acagetttta gaactggtta caatagegae ggagagttag gttattggga taagttagag 4200 ccactttata caatttttga tggtgtatct aaaacattct ctggtatttg gactcctgta 4260 aagaatgact tcaaagagtt ttatgattta tacctttctg atgtagagaa atataatggt 4320 tcggggaaat tgtttcccaa aacacctata cctgaaaatg ctttttctct ttctattatt 4380 ccatggactt catttactgg gtttaactta aatatcaata ataatagtaa ttaccttcta 4440 cccattatta cagcaggaaa attcattaat aaaggtaatt caatatattt accgctatct 4500 ttacaggtac atcattctgt ttgtgatggt tatcatgcag gattgtttat gaactctatt 4560 caggaattgt cagataggcc taatgactgg cttttataat atgagataat gccgactgta 4620 ctttttacag tcggttttct aatgtcacta acctgccccg ttagttgaag aaggttttta 4680 tattacaget gtegactegt gatettegga caggetgtte agetttttet caatgegate 4740 cagctgcqct tttcqqtttt tcqcatactt gaagcctgta acagccgcaa agacgacagc 4800 ctcatgtttg cgggagagat tcattctctt ccgtttttta tttaaagcgg cttttccaga 4920 egggaaeggt gttttgtggt etecatttte atttgeegat aggegaaege taaaaatgge 4980 aggeegagea gggtaatgee geteaggaea gaaaaaatat aaateggeeg geeagegeea 5040 aacaggteta tacatatece eeegaeeeaa gggeegatga egttteegag etgtggaaaa 5100 ccgattgccc cgaaataagt gccttttaat cctggttttg caatctggtc tacatacaaa 5160 tccatcatag agaataaaag cacttcgccg attgtaaatg tgatgacaat catcacaatt 5220 gatggaacac cgtgtgatac ggtgaaaatg gccatgctga tgctaaccat cacattaccg 5280 agcatcagag aacaaagcgg cgaaaaccgt tttgcaaaat ggacaatggg aaattgcgtc 5340 gccaacaca cgattgcgtt taatgtcagc atcagcccat acagcttcgt tccattgccg 5400 atcaaggggt totgcgccat atactgaggg aatgtggaac tgaattgtga gtagccgaag 5460 qtqcataqcq taatgccgac caaagcaatg gtaaaaagat aatccttttg cgtgaccata 5520 aacgcttccc qcacgctcat atttcgggac tgggctggtg ctgataagga tggatgtttt 5580 ttaaattgga gggcaagcac aattccgtat agtccgtaaa tgactgcagg caccaaaaag 5640 ggcgtagtcg attgcgatga gccgaaatat aggccaagca caggtccgaa gacaacgccg 5700 atattaatag ccgcatagcg taaattaaaa actagcagtc tcgttttttc ttctgtcata 5760 tcagacaaca aggcctttga agcgggctca aacagtgatt tgcaaagacc gtttaatgcg 5820 tttactacaa aaaacaccca gagattagat gctgccgcaa agcctgcaaa taccagcatc 5880 catccgaaaa tcgatacaag catcatgttt tttctgccga atttatctga gatatatccg 5940 ccgtaaaagc ttgcgaggat gccgactgat gagctcgcgg cgatgaccag ccctgcatag 6000 gaagetgatg egeettggae ggetgteaaa taaategeta aaaaaggaat geteategat 6060 gttgccattc tgccgaaaat ggttccgatt ataattgtac gcgttggatg catagcttga 6120 gtattctata gtgtcaccta aatagcttgg cgtaatcatg gtcatagctg tttcctgtgt 6180 gaaattgtta teegeteaca atteeacaca acataegage eggaageata aagtgtaaag 6240 cctggggtgc ctaatgagtg agctaactca cattaattgc gttgcgctca ctgcccgctt 6300 tccagtcggg aaacctgtcg tgccagctgc attaatgaat cggccaacgc gcggggagag 6360 geggtttgeg tattgggege tetteegett cetegeteae tgaetegetg egeteggteg 6420 tteggetgeg gegageggta teageteaet caaaggeggt aataeggtta teeacagaat 6480 caggggataa cgcaggaaag aacatgtgag caaaaggcca gcaaaaggcc aggaaccgta 6540 aaaaggccgc gttgctggcg tttttcgata ggctccgccc ccctgacgag catcacaaaa 6600 atcgacgete aagteagagg tggegaaaee egacaggaet ataaagatae caggegttte 6660 cccctggaag ctccctcgtg cgctctcctg ttccgaccct gccgcttacc ggatacctgt 6720 ccgcctttct cccttcggga agcgtggcgc tttctcatag ctcacgctgt aggtatctca 6780 gttcggtgta ggtcgttcgc tccaagctgg gctgtgtgca cgaacccccc gttcagcccg 6840 accgetgege ettateeggt aactategte ttgagteeaa eeeggtaaga caegaettat 6900 cgccactggc agcagccact ggtaacagga ttagcagagc gaggtatgta ggcggtgcta 6960 cagaqttctt gaaqtqqtqq cctaactacq qctacactaq aaqqacaqta tttgqtatct 7020 qcqctctqct qaaqccagtt accttcggaa aaagagttgg tagctcttga tccggcaaac 7080 aaaccaccgc tggtagcggt ggtttttttg tttgcaagca gcagattacg cgcagaaaaa 7140 aaggatetea agaagateet ttgatetttt etaeggggte tgaegeteag tggaaegaaa 7200 actcacgtta agggattttg gtcatgagat tatcaaaaag gatcttcacc tagatccttt 7260 taaattaaaa atgaagtttt aaatcaatct aaagtatata tgagtaaact tggtctgaca 7320 gttaccaatq cttaatcaqt gaggcaccta tctcagcgat ctgtctattt cgttcatcca 7380 tagttgcctg actccccgtc gtgtagataa ctacgatacg ggagggctta ccatctggcc 7440 ccagtgctgc aatgataccg cgagacccac gctcaccggc tccagattta tcagcaataa 7500 accagecage eggaagggee gagegeagaa gtggteetge aactttatee geeteeatee 7560 agtctattaa ttgttgccgg gaagctagag taagtagttc gccagttaat agtttgcgca 7620 acgttgttgg cattgctaca ggcatcgtgg tgtcacgctc gtcgtttggt atggcttcat 7680 teageteegg tteecaaega teaaggegag ttacatgate eeccatgttg tgeaaaaaag 7740 eggttagete etteggteet eegategttg teagaagtaa gttggeegea gtgttateae 7800 tcatggttat ggcagcactg cataattctc ttactgtcat gccatccgta agatgctttt 7860 ctgtgactgg tgagtactca accaagtcat tctgagaata ccgcgcccgg cgaccgagtt 7920 gctcttgccc ggcgtcaata cgggataata gtgtatgaca tagcagaact ttaaaagtgc 7980 tcatcattgg aaaacgttct teggggegaa aactetcaag gatettaceg etgttgagat 8040 ccagttcgat gtaacccact cgtgcaccca actgatcttc agcatctttt actttcacca 8100 gcgtttctgg gtgagcaaaa acaggaaggc aaaatgccgc aaaaaaggga ataagggcga 8160 cacggaaatg ttgaatactc atactcttcc tttttcaata ttattgaagc atttatcagg 8220 gttattgtct catgagcgga tacatatttg aatgtattta gaaaaataaa caaatagggg 8280 ttccgcgcac atttccccga aaagtgccac ctgtatgcgg tgtgaaatac cgcacagatg 8340 cgtaaggaga aaataccgca tcaggcgaaa ttgtaaacgt taatattttg ttaaaattcg 8400 cgttaaatat ttgttaaatc agetcattt ttaaccaata ggccgaaatc ggcaaaatcc 8460 cttataaatc aaaagaatag accgagatag ggttgagtgt tgttccagtt tggaacaaga 8520 gtccactatt aaagaacgtg gactccaacg tcaaagggcg aaaaaccgtc tatcagggcg 8580 atggcccact acgtgaacca tcacccaaat caagttttt gcggtcgagg tgccgtaaag 8640 ctctaaatcg gaaccctaaa gggagcccc gatttagagc ttgacggga aagccggcga 8700 acgtggcgag aaaggaaggg aagaaagcga aaggagcggg cgctagggcg ctggcaagtg 8760 tagcggtcac gctgcgcgta accaccacac ccgccgcgct taa 8803

<210> 73

<211> 8320

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant
pAN296 plasmid

tgcgccgcta cagggcgcgt ccattcgcca ttcaggctgc gcaactgttg ggaagggcga 60 teggtgeggg cetetteget attacgeeag etggegaaag ggggatgtge tgeaaggega 120 ttaagttggg taacgccagg gttttcccag tcacgacgtt gtaaaacgac ggccagtgaa 180 ttgtaatacg actcactata gggcgaattg ggcccgacgt cgcatgctgg atgaaaagcc 240 gatgaccgct tttcaggtct gtcagcagct ttttcctgct gtatatgaaa aggaattgtt 300 tttaacgatg tcagaaacgg caggtcacct tgatgtgttg gaggctgaag aagccatcac 360 gtcatattgg gaaggaaata ccgtatactt taaaacaatg aagaggtgaa atgggtgaaa 420 catatagcgg gaaaaaggat ttggataacc ggcgcttcag gagggcttgg agaaagaatc 480 gcatacttat gcgcggctga aggagcccat gtcctgctgt cggctagacg cgaggatcgt 540 ttgatagaaa tcaaaaggaa aataaccgag gaatggagcg gacagtgtga gatttttcct 600 ctggatgtcg gccgcctaga ggatatcgcc cgggtccgcg atcagatcgg ctcgattgat 660 gtactgatta acaatgcagg cttcggtata tttgaaacgg ttttagactc tacattggat 720 gacatgaaag cgatgtttga tgtgaatgtc ttcggcctga tcgcctgtac aaaagcggtg 780 cttccgcaaa tgcttgagca aaaaaaggga catatcatca atatcgcctc tcaagcgggg 840 aaaatcgcca caccgaagtc tagcctgtat tccgcgacca aacatgccgt gttaggttac 900 teaaacgett tgeggatgga gettteggga aceggeattt atgtgacaac agteaaceeg 960 ggcccgattc agacggactt tttttccatt gctgataaag gcggggacta cgccaaaaat 1020 gteggeeget ggatgettga teetgatgae gtggeagete aaattacage tgeaattttt 1080 acgaaaaagc gggagatcaa tcttccgcgt ttaatgaatg ccggcactaa gctgtatcag 1140 ctgtttccag ctcttgtaga aaagctggca ggacgcgcgc tcatgaaaaa ataatgatag 1200 aactgcctgt ggtggagtgg cttgtttctc acggggcagt ttttgatagt ggaagggaga 1260 gattgttgaa tgtcagttca ttcagaagtc cttcatgctc tgcttaaaga tccgtttatt 1320 cagaaactga ttgatgcaga gcctgtattc tgggcaaatt caggcaagaa agaggggcca 1380 ttaccccgtg cagatgagtg ggcaaccgag atagcggaag cggaaaaaag aatgcagcgg 1440 tttgcacctt acattgccga ggtgtttcct gagacgaaag gcgctaaagg aatcatcgag 1500 teteegettt ttgaggtgea geatatgaag ggaaagetgg aageggeata teageageea 1560 tttcccggaa gatggctttt aaagtgcgac catgagcttc cgatttcagg atcgattaaa 1620 gcgaggggcg ggatttatga agtgttaaag tatgctgaaa atctcgcgct tcaagaagga 1680 atgcttcagg aaaccgatga ttaccgcatc ttacaggaag agcggtttac cgggttttc 1740 tecegetatt egattgetgt eggttegaea ggaaatetag gtttaageat eggeateate 1800 ggcgcggcac tcgggtttcg cgtgacagtg catatgtccg ccgatgctaa gcagtggaaa 1860 aaggatetee teegeeaaaa qqqaqteact qttatqqaqt acqaaacaqa ttacaqtqaa 1920 gcggtgaacg aagggagacg gcaggcggaa caagatccat tctgttattt tattgatgat 1980 gaacattctc gtcagctgtt cttaggatat gctgttgctg caagccgatt aaaaacacag 2040 cttgactgta tgaatataaa gccaagtctt gagacgccct tgtttgtgta tctgccgtgc 2100 ggagtcggcg gaggaccggg cggtgtagca tttgggctga agcttttata cggagatgat 2160 gttcatgtgt ttttcgcaga accaactcat tcaccttgta tgctgttagg gctttattca 2220 ggacttcacg agaagatctc cgtccaggat atcggcctgg ataatcagac ggctgctgac 2280 ggacttgccg tagggaggcc gtcaggattt gtcggcaagc tgattgaacc gcttctgagc 2340 ggctgttata cggtagagga caatacgctt tatactttgc ttcatatgct ggctgtatct 2400 gaagataaat atttagagcc ctctgctctt gctggcatgt tcgggccggt tcagcttttt 2460 tcgacagaag agggaaggcg ctatgctcag aaatataaga tggaacatgc cgtacatgtc 2520 gtctggggaa cgggaggaag catggttcca aaagatgaaa tggctgcgta taaccgaatc 2580 ggtgctgatt tgctaaaaaa acgaaatgga aaataagcag acagtgaaaa ggttttccgt 2640 tacaatcttt gtaagggttt taacctacag agagtcaggt gtaaacagtg aaaaataaag 2700 aacttaacct acatacttta tatacacagc acaatcggga gtcttctgca gctcgagcaa 2760 tagttaccct tattatcaag ataagaaaga aaaggatttt tcgctacgct caaatccttt 2820 aaaaaaacac aaaagaccac attttttaat gtggtcttta ttcttcaact aaagcaccca 2880 ttagttcaac aaacgaaaat tggataaagt gggatatttt taaaaatatat atttatgtta 2940 caqtaatatt gactittaaa aaaggattga tictaatgaa gaaagcagac aagtaagcct 3000 cctaaattca ctttagataa aaatttagga ggcatatcaa atgaacttta ataaaattga 3060 tttagacaat tggaagagaa aagagatatt taatcattat ttgaaccaac aaacgacttt 3120 tagtataacc acagaaattg atattagtgt tttataccga aacataaaac aagaaggata 3180 taaattttac cctgcattta ttttcttagt gacaagggtg ataaactcaa atacagcttt 3240 tagaactggt tacaatagcg acggagagtt aggttattgg gataagttag agccacttta 3300 tacaattttt gatggtgtat ctaaaacatt ctctggtatt tggactcctg taaagaatga 3360 cttcaaagag ttttatgatt tatacctttc tgatgtagag aaatataatg gttcggggaa 3420 attgtttccc aaaacaccta tacctgaaaa tgctttttct ctttctatta ttccatggac 3480 ttcatttact gggtttaact taaatatcaa taataatagt aattaccttc tacccattat 3540 tacagcagga aaattcatta ataaaggtaa ttcaatatat ttaccgctat ctttacaggt 3600 acatcattet gtttgtgatg gttatcatge aggattgttt atgaacteta ttcaggaatt 3660 gtcagatagg cctaatgact ggcttttata atatgagata atgccgactg tactttttac 3720 agteggtttt ctaatgtcac taacetgeec egttagttga agaaggtttt tatattacag 3780 ctgtcgacta aggtcgagga agtgttggta aggagggtat gaaatgtgca tcatattgaa 3840 ctgtatgtct ctgatttgga ggcgtctagg cggtttttggg gctggttctt aaaagaactt 3900 ggttataaag agtatcaaaa atggagctca ggcatcagct ggaagaaaga tcgtttttac 3960 ctagtgattg tgcaggcgaa agagccattt ctagagccgg aataccatag atgccgagtc 4020 ggtctgaacc atctcgcatt tcatgctgaa tccaagcttc aagtcgatca gatgactgaa 4080 aaattgacgg caaaaggcta tcgtgtgttg taccgagaca ggcatccttt tgccggagga 4140 gacgggcatt atgcagtctt ttgtgaggat ccagaccgga ttaaggtaga gctcgttgcc 4200 ccaagctgtt aatcgtgatc ttcggacagg ctgttcagct ttttctcaat gcgatccagc 4260 tgcgcttttc ggtttttcgc atacttgaag cctgtaacag ccgcaaagac gacagcggca 4320 aatataataa atacaaacag ctgaaacatc acatcaccta tattcatgtt cttcacctca 4380 tgtttgcggg agagattcat tctcttccgt tttttattta aagcggcttt tccagacggg 4440 aacggtgttt tgtggtctcc attttcattt gccgataggc gaacgctaaa aatggcaggc 4500 cgagcagggt aatgccgctc aggacagaaa aaatataaat cggccggcca gcgccaaaca 4560 ggtctataca tatccccccg acccaagggc cgatgacgtt tccgagctgt ggaaaaccga 4620 ttgccccgaa ataagtgcct tttaatcctg gttttgcaat ctggtctaca tacaaatcca 4680 tcatagagaa taaaagcact tegeegattg taaatgtgat gacaatcate acaattgatg 4740 gaacaccgtg tgatacggtg aaaatggcca tgctgatgct aaccatcaca ttaccgagca 4800 tcagagaaca aagcggcgaa aaccgttttg caaaatggac aatgggaaat tgcgtcgcca 4860 acacaacgat tgcgtttaat gtcagcatca gcccatacag cttcgttcca ttgccgatca 4920 aggggttctg cgccatatac tgagggaatg tggaactgaa ttgtgagtag ccgaaggtgc 4980 atagcgtaat gccgaccaaa gcaatggtaa aaagataatc cttttgcgtg accataaacg 5040 cttcccgcac gctcatattt cgggactggg ctggtgctga taaggatgga tgttttttaa 5100 attggagggc aagcacaatt ccgtatagtc cgtaaatgac tgcaggcacc aaaaagggcg 5160 tagtcgattg cgatgagccg aaatataggc caagcacagg tccgaagaca acgccgatat 5220 taatagccgc atagcgtaaa ttaaaaacta gcagtctcgt tttttcttct gtcatatcag 5280 acaacaaggc ctttgaagcg ggctcaaaca gtgatttgca aagaccgttt aatgcgttta 5340 ctacaaaaaa cacccagaga ttagatgctg ccgcaaagcc tgcaaatacc agcatccatc 5400 cgaaaatcga tacaagcatc atgtttttc tgccgaattt atctgagata tatccgccgt 5460 aaaagettge gaggatgeeg actgatgage tegeggegat gaccageeet geataggaag 5520 ctgatgcgcc ttggacggct gtcaaataaa tcgctaaaaa aggaatgctc atcgatgttg 5580 ccattctgcc gaaaatggtt ccgattataa ttgtaacgcg ttggatgcat agcttgagta 5640 ttctatagtg tcacctaaat agcttggcgt aatcatggtc atagctgttt cctgtgtgaa 5700 attgttatcc gctcacaatt ccacacaaca tacgagccgg aagcataaag tgtaaagcct 5760 ggggtgccta atgagtgagc taactcacat taattgcgtt gcgctcactg cccgctttcc 5820 agtcgggaaa cctgtcgtgc cagctgcatt aatgaatcgg ccaacgcgcg gggagaggcg 5880 gtttgcgtat tgggcgctct tccgcttcct cgctcactga ctcgctgcgc tcggtcgttc 5940 ggctgcggcg agcggtatca gctcactcaa aggcggtaat acggttatcc acagaatcag 6000 gggataacgc aggaaagaac atgtgagcaa aaggccagca aaaggccagg aaccgtaaaa 6060 aggccgcgtt gctggcgttt ttcgataggc tccgccccc tgacgagcat cacaaaaatc 6120 gacgeteaag teagaggtgg egaaaceega eaggaetata aagataceag gegttteeee 6180 ctggaagete eetegtgege teteetgtte egaceetgee gettaeegga taeetgteeg 6240 cetttetece ttegggaage gtggegettt eteatagete aegetgtagg tateteagtt 6300 cggtgtaggt cgttcgctcc aagctgggct gtgtgcacga accccccgtt cagcccgacc 6360 gctgcgcctt atccggtaac tatcgtcttg agtccaaccc ggtaagacac gacttatcgc 6420 cactggcagc agccactggt aacaggatta gcagagcgag gtatgtaggc ggtgctacag 6480 agttettgaa gtggtggeet aactaegget acaetagaag gaeagtattt ggtatetgeg 6540 ctctgctgaa gccagttacc ttcggaaaaa gagttggtag ctcttgatcc ggcaaacaaa 6600 ccaccgctgg tagcggtggt ttttttgttt gcaagcagca gattacgcgc agaaaaaaa 6660 gatctcaaga agatcctttg atcttttcta cggggtctga cgctcagtgg aacgaaact 6720 cacgttaagg gattttggtc atgagattat caaaaaggat cttcacctag atccttttaa 6780 attaaaaatg aagttttaaa tcaatctaaa gtatatatga gtaaacttgg tctgacagtt 6840 accaatgett aatcagtgag geacetatet cagegatetg tetatttegt teatecatag 6900 ttgcctgact ccccgtcgtg tagataacta cgatacggga gggcttacca tctggcccca 6960 gtgctgcaat gataccgcga gacccacgct caccggctcc agatttatca gcaataaacc 7020 agccagccgg aagggccgag cgcagaagtg gtcctgcaac tttatccgcc tccatccagt 7080 ctattaattg ttgccgggaa gctagagtaa gtagttcgcc agttaatagt ttgcgcaacg 7140 ttgttggcat tgctacaggc atcgtggtgt cacgctcgtc gtttggtatg gcttcattca 7200 geteeggtte ecaacgatea aggegagtta catgateece catgttgtge aaaaaagegg 7260 ttagctcctt cggtcctccg atcgttgtca gaagtaagtt ggccgcagtg ttatcactca 7320 tggttatggc agcactgcat aattctctta ctgtcatgcc atccgtaaga tgcttttctg 7380 tgactggtga gtactcaacc aagtcattct gagaataccg cgcccggcga ccgagttgct 7440 cttgcccggc gtcaatacgg gataatagtg tatgacatag cagaacttta aaagtgctca 7500 tcattggaaa acgttcttcg gggcgaaaac tctcaaggat cttaccgctg ttgagatcca 7560 gttegatgta acceaetegt geaeceaaet gatetteage atetttaet tteaecageg 7620 tttctgggtg agcaaaaaca ggaaggcaaa atgccgcaaa aaagggaata agggcgacac 7680 ggaaatgttg aatactcata ctcttccttt ttcaatatta ttgaagcatt tatcagggtt 7740 attgtctcat gagcggatac atatttgaat gtatttagaa aaataaacaa ataggggttc 7800 cgcgcacatt tccccgaaaa gtgccacctg tatgcggtgt gaaataccgc acagatgcgt 7860 aaggagaaaa taccgcatca ggcgaaattg taaacgttaa tattttgtta aaattcgcgt 7920 taaatatttg ttaaatcagc tcattttta accaataggc cgaaatcggc aaaatccctt 7980 ataaatcaaa agaatagacc gagatagggt tgagtgttgt tccagtttgg aacaagagtc 8040 cactattaaa gaacgtggac tccaacgtca aagggcgaaa aaccgtctat cagggcgatg 8100 geceactacg tgaaccatca eccaaatcaa gttttttgeg gtegaggtge egtaaagete 8160 taaatcggaa ccctaaaggg agcccccgat ttagagcttg acggggaaag ccggcgaacg 8220 tggcgagaaa ggaagggaag aaagcgaaag gagcgggcgc tagggcgctg gcaagtgtag 8280 8320 cggtcacgct gcgcgtaacc accacacccg ccgcgcttaa

<210> 74

<211> 250

<212> PRT

<213> Clostridium acetobutylicum

<400> 74

Asn Lys Arg Ala Ala Phe Met Leu Leu Leu Phe Leu Arg Ser Val Leu 1 5 10 15

Lys Val Ile Leu Val Leu Asp Val Gly Asn Thr Asn Ile Val Leu Gly 20 25 30

Ile Tyr Asn Asp Thr Lys Leu Thr Ala Glu Trp Arg Leu Ser Thr Asp 35 40 45

Val Leu Arg Ser Ala Asp Glu Tyr Gly Ile Gln Val Met Asn Leu Phe 50 55 60

Gln Gln Asp Lys Leu Asp Pro Thr Leu Val Glu Gly Val Ile Ile Ser 65 70 75 80

Ser Val Val Pro Asn Ile Met Tyr Ser Leu Glu His Met Ile Arg Lys 85 90 95

Tyr Phe Lys Ile Asn Pro Leu Val Val Gly Pro Gly Ile Lys Thr Gly 100 105 110

Ile Asn Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile 115 120 125

Val Asn Ala Val Ala Ala His Glu Ile Tyr Lys Arg Ser Leu Ile Ile 130 135 140

Ile Asp Phe Gly Thr Ala Thr Thr Phe Cys Ala Val Arg Glu Asn Gly 145 150 155 160

Asp Tyr Leu Gly Gly Ala Ile Cys Pro Gly Ile Lys Val Ser Ser Glu 165 170 175

Ala Leu Phe Glu Lys Ala Ala Lys Leu Pro Arg Val Glu Leu Ile Lys 180 185 190

Pro Ala Tyr Ala Ile Cys Lys Asn Thr Ile Ser Ser Ile Gln Ser Gly 195 200 205

Ile Val Tyr Arg Tyr Leu Arg Gln Val Lys Tyr Leu Phe Glu Lys Leu 210 215 220

Lys Glu Asn Leu Pro Asp Gly Arg Arg Thr Arg Thr Ser Leu Val Leu 225 230 235 240

Ala Thr Gly Gly Leu Ala Lys Leu Ile Asn 245 250

<210> 75

<211> 258

<212> PRT

<213> Rhodobacter capsulatus

<400> 75

Met Leu Cys Ile Asp Cys Gly Asn Thr Asn Thr Val Phe Ser Val
1 5 10 15

Trp Asp Gly Thr Asp Phe Ala Ala Thr Trp Arg Ile Ala Thr Asp His 20 25 30

Arg Arg Thr Ala Asp Glu Tyr Phe Val Trp Leu Asn Thr Leu Met Gln 35 40 45

Leu Lys Gly Leu Gln Gly Arg Ile Ser Glu Ala Ile Ile Ser Ser Thr 50 60

Ala Pro Arg Val Val Phe Asn Leu Arg Val Leu Cys Asn Arg Tyr Phe 65 70 75 80

Asp Cys Arg Pro Tyr Val Val Gly Lys Pro Gly Cys Glu Leu Pro Val 85 90 95

Ala Pro Arg Val Asp Pro Gly Thr Thr Val Gly Pro Asp Arg Leu Val 100 105 110

Asn Thr Val Ala Gly Tyr Asp Arg His Gly Gly Asp Leu Ile Val Val 115 120 125

Asp Phe Gly Thr Ala Thr Thr Phe Asp Val Val Ala Pro Asp Gly Ala 130 135 140

Tyr Ile Gly Gly Val Ile Ala Pro Gly Val Asn Leu Ser Leu Glu Ala 145 150 155 160

Leu His Met Ala Ala Ala Ala Leu Pro His Val Asp Val Thr Lys Pro 165 170 175

Gln Gly Val Ile Gly Thr Asn Thr Val Ala Cys Ile Gln Ser Gly Val 180 185 190

Tyr Trp Gly Tyr Ile Gly Leu Val Glu Gly Ile Val Arg Gln Ile Arg 195 200 205

Met Glu Arg Asp Arg Pro Met Lys Val Ile Ala Thr Gly Gly Leu Ala 210 215 220

Ser Leu Phe Asp Leu Gly Phe Asp Leu Phe Asp Lys Val Glu Asp Asp 225 230 235 240

Leu Thr Met His Gly Leu Arg Leu Ile Phe Asp Tyr Asn Lys Gly Leu 245 250 255

Gly Ala

<210> 76

<211> 10801

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant
pAN240 plasmid

<400> 76 gaattttgcg gccgcttcga aagctgtaat ataaaaacct tcttcaacta acggggcagg 60 ttagtgacat tagaaaaccg actgtaaaaa gtacagtcgg cattatctca tattataaaa 120 gccagtcatt aggcctatct gacaattcct gaatagagtt cataaacaat cctgcatgat 180 aaccatcaca aacagaatga tgtacctgta aagatagcgg taaatatatt gaattacctt 240 tattaatgaa ttttcctgct gtaataatgg gtagaaggta attactatta ttattgatat 300 ttaagttaaa cccagtaaat gaagtccatg gaataataga aagagaaaaa gcattttcag 360 gtataggtgt tttgggaaac aatttccccg aaccattata tttctctaca tcagaaaggt 420 ataaatcata aaactctttg aagtcattct ttacaggagt ccaaatacca gagaatgttt 480 tagatacacc atcaaaaatt gtataaagtg gctctaactt atcccaataa cctaactctc 540 cgtcgctatt gtaaccagtt ctaaaagctg tatttgagtt tatcaccctt gtcactaaga 600 aaataaatgc agggtaaaat ttatatcctt cttgttttat gtttcggtat aaaacactaa 660 tatcaatttc tgtggttata ctaaaagtcg tttgttggtt caaataatga ttaaatatct 720 cttttctctt ccaattgtct aaatcaattt tattaaagtt catttgatat gcctcctaaa 780 tttttatcta aagtgaattt aggaggetta ettgtetget ttetteatta gaatcaatee 840 ttttttaaaa gtcaatatta ctgtaacata aatatatatt ttaaaaatat cccactttat 900 ccaattttcg tttgttgaac taatgggtgc tttagttgaa gaataaagac cacattaaaa 960 aatgtggtct tttgtgtttt tttaaaggat ttgagcgtag cgaaaaatcc ttttctttct 1020 tatcttgata ataagggtaa ctattgaatt cggtaccaag agtttgtaga aacgcaaaaa 1080 ggccatccgt caggatggcc ttctgcttaa tttgatgcct ggcagtttat ggcgggcgtc 1140 ctgcccgcca ccctccgggc cgttgcttcg caacgttcaa atccgctccc ggcggatttg 1200 tectacteag gagagegtte acegacaaac aacagataaa acgaaaggee cagtettteg 1260 actgagcett tegttttatt tgatgeetgg cagtteecta etetegeatg gggagaeece 1320 acactaccat cggcgctacg gcgtttcact tctgagttcg gcatggggtc aggtgggacc 1380 accgcgctac tgccgccagg caaattctgt tttatcagac cgcttctgcg ttctgattta 1440 atctgtatca ggctgaaaat cttctctcat ccgccaaaac aggatcctac ggaaatggag 1500 cggcaaaacc gttttactct caaaatctta aaagaaaacc cccgataaag ggggcttttc 1560 ttctacaaaa ttgtacgggc tggttcgttc cccagcattt gttcaatttt gttttgatca 1620 ttcagaacag ccactttcgg ctcatggctt gccgcttctt gatcagacat cattttgtag 1680 gaaataataa tgaccttatc tccttcctgc acaaggcgtg cggctgcacc gtttaagcat 1740 atgacgccgc ttccccgttt accaggaata atatacgttt caagacgtgc tccattatta 1800 ttattcacaa tttgtacttt ttcattagga agcattccca cagcatcaat gagatcttca 1860 tcaattgtaa tgcttcccac atagttcagg tttgcttccg taacagttgc cctgtgaagt 1920 ttgccgctca tcattgttcg atacatatta tattctctcc atttctcgaa tatcaataat 1980 gatattatct attaaacgcg cttttgaaaa agcaactgca acagcgagaa tcatctttcc 2040 agcaatttca ttcacaggct cgagttccgg ataggaataa agctctacat agtctatggt 2100 teegetagte gttteaatga tatettttge agettttate acegetteag gatetettte 2160 accggcttgg acaagttccg cacttgtttg aagggcccga tacagcttag gcgcttcttt 2220 tettteetea getgttaagt atacattgeg agagettttg getaageegt etteetetet 2280 gaccgtatcg acaggaacca attcaatatc catgaagaag tcgctgatta acccatcaac 2340 aacagctacc tgctgcgcat cttttaaacc gaaataggca cgagtcggct tgactagatt 2400 gaaaagcttc gtcagtacga tcgcgacccc gtcaaaatgt ccttctcttg agcgcccgca 2460 taacacgtct gtgcgtcttt ctacatgaat cgtgacattc ttttcaccgg gatacatatc 2520 atgagcatet ggegtaaaaa gaatategae teeggegttt tetgeaagag etgeateeeg 2580 ctcaatatcg cgcggatatg cttcaaaatc ttcattaggg ccgaattgtg caggattcac 2640 aaaaatactc ataataacgg cgtcgttttc ttgtcttgct ttgtctgcta aggttaaatg 2700 cccctcatgc agaaacccca tcgtcggaac aaatccgatt gacttgccct ctgaatggta 2760 ttgttttatg gcttctttca gctgtgaaat atcagtaatc tgtctcatct tattttcccc 2820 cgtacaagcc gtcaagcact gtctggttca tttgaaagga atgcttttgt tcagggaaag 2880 cacgatgtct tacatcctga acatatccgc tgattgctgt ttcgatggtt tcatcaatgc 2940 gcgtatattg ctttacaaat ttaggtgttc tctcaacacc gtggccgata atatcatgat 3000 aaacgagaac ttgtccgtcc gctttcacac cagccccgat tccaatgacc ggtatgctta 3060 gcgtctcggc aattttggct gtgagttctg ccggcacaca ttccagcaca agcatcatag 3120 ctcctgcttc ttcgcatttt atactgtctt ctattaattt tttggcgctt tgttcgtctt 3180 tgccctgtac tttatagccg cccagtacgc cgactgactg cggtgtcaaa cctaagtgac 3240 tgactactgg aatgcctcca agcgtcaatg cgcgaatgga ttcaaacacg ccttctccgc 3300 cctcaagctt cagtgcgtca gctccgcttt cctgaacgat agccgctgca tttttcagcg 3360 tatcttcctt agacaggtga taagacataa acggcatatc tgtcacaata aaggtattcg 3420 gegeacecet tttaaegget tttgtatgat ggateatgte egeaactgte acaeeggaeag 3480 ttgaatcaag gccgaggacg accattccaa gtgaatcacc gactaaaatc atgtcaactc 3540 ccgcttgttc agcaagttta gctgccggat aatcataagc ggtcagcatg acaatcggtt 3600 gagcgtcctg ctgttgttaa gattattata ccacaccttg tagataaagt caacaacttt 3720 ttgcaaaatt tttcaggaat tttagcagag gttgttctgg atgtagaaca aaacatcttt 3780 cegetettgt getgttagga tatetttett ggaagetagg taggeetega gttatggeag 3840 ttggttaaaa ggaaacaaaa agaccgtttt cacacaaaac ggtcttttc gatttctttt 3900 tacagtcaca gccacttttg caaaaaccgg acagettcat gccttataac tgctgtttcg 3960 gtcgacgatg atctgccgtt ttcttctgca agccaaaaaa ccttccgtta caacgagaag 4020 gattetteae tttetaaagt teggegagtt teateeetet gteeeagtee ttttttggat 4080 caaggcagac tgctgcaatg tctatctatt ttaataatag gtgcagttcg caggcgatac 4140 tgcccaatgg aagtatacca aaatcaacgg gcttgtacca acacattagc ccaattcgat 4200 atcggcagaa tagattttt taatgccttc gttcgtttct aaaagcagaa cgccttcatc 4260 atctatacct aacgccttac cgtaaaaggt tccgtttaac gttctggctc tcatattagt 4320 gccaataccg agcgcatagc tttcccataa aagcttaatc ggcgtaaatc cgtgcgtcat 4380 ataatcccgg taccgtttct caaagcatag taaaatatgc tggatgacgc cggcccgatc 4440 aattttttcc ccagcagctt ggctgaggct tgtcgcgatg tccttcaatt catctggaaa 4500 atcattaggc tgctggttaa cgttaatgcc gatcccaatg atcactgaac gtacgcggtc 4560 ttcttcagcc tgcatttccg ttaggatacc gactgttttt tttccgttaa tcaaaatatc 4620 atttggccat ttaatatccg tttggatgcc tgctgcctct tctattccct gcacaacagc 4680 tactgcagca agcagagtca gctgcggtgt tttttggagc ggaatgtcag gccgcaaaat 4740 caggeteate caaacacegt tteettettg agaatgeeat accetagaca tteggeeeet 4800 tccggctgtt tgtttgtcag ccaccacaag ggtgccttcc ggtgcgttat tattcgcgag 4860 ctcatgagcc gttttttgcg tgcttgaaag aacgtcatgg taaataagat gctggcccat 4920 cacttccgtt tttaatccaa aacgaatttc gctttcactg agttttccgg gttttttgat 4980 gageegatat cettttette taaeggette taetteataa eeetetttee gaagetette 5040 aatatgette cacacageag ttettgaaca geegagagea teaetgattt tttggeegga 5100 aataaattca ttgccggcct gagaaaataa ttcaataagg tcttttctta atgttgaccg 5160 catgtcttca gccactcctc tatgtgtttc ttttgattgg agagcttccc tgtcacaaca 5220 gcctgctcga tccactgtaa ttcttctgac acccattttc cggccggccg gtttcgaagc 5280 gcaagcaagt ccttacccgt gatatcaaga tccttaaggc ttttgatcgg caggttttga 5340 taagcgtact gaatgtcctt cagtttcttt tcatccagtt tttcgttttg ccgaagctgc 5400 gatattttgg ccgctgagag cagtgctttt ttcccagctc tgtacattgt cattgcgtca 5460 aggetetgge caaacgtate ggeaatgtga atggetteet tgateaettt teeegggage 5520 ttccaggett tcaggaaaag gggcgcgtct ttcaaaacta tgccaaggtt aattaaaaga 5580 gcagcccaaa gctcctcacg ggatgttaaa gagaagaatg gaaactcact cgttgaaatc 5640 aggttttctc gtttatgata aaaaccagga agctcttcat acaatctcgt ttgaatgagt 5700 gtttgaagcg cctggcgaga agctcttccc tgcagcaatt tctcaaactc tatagttttt 5760 cgttcgactg aaacatggga gaggagtgat ttttctttcg caatggcttc ttctgtttcc 5820 ggtgaaagcg taaagccaag ctggctcata aagcgtacgg ctctcagcat acgaagcgca 5880 tectettgaa atetateete aggettteea aeggttegaa teaetttetg ateaatatet 5940 ttettgeege caaaataate aageacette eegteegetg teatggeeat egeattgate 6000 gttaaatctc tgcgttttag atcctcttct aatgatgaga taaattgcac ttctgacggt 6060 cttctgaaat caacataatc agattcagtc cggaatgtcg tgacttcata ggtttcatcc 6120 teccagagea caataatggt ecegtgetet ttgeetacat caacagteeg etgaaacage 6180 cgttctactt gatcaggtgc cgcatctgtc gcgatatcga catctccgat cgttcgtttc 6240 atatagctgt cacgaactgc gcccccgaca aaataagcct gatggcccgc ttcgattaag 6300 atgcggagca cgggaagtgc tttgataaaa actttttcca tgtgatcact ccggttctgc 6360 taaatcggca taaatctgtt catactggct gacaattttt ttagaagaaa attcattttc 6420 aagcatetet attgeegeet ttgtaaaaeg attgettage tgtteatett etaaaatget 6480 catcgcgcgg gctgttgcgg ccgtaacatc accgacatcc accaaaaatc cgctcacatt 6540 gttttttata acctcaggga taccgccaat gtttgttcca atacaaggca ctccgcaagc 6600 categettea ageaggacaa ggeeaaaget ttettttea gatageagea getteaaate 6660 gctaatagaa taaagatett caacaeggte ttgattteea ageattaaga ettggtette 6720 caagccatat tttctgataa gctcgcaggc tgtcgatttc tccggaccgt ctccgactaa 6780 aagcagcttc gctttcgttt tgccagcgat attgcggaac acacggatga catcctgcac 6840 gcgtgcagcc actggtaaca ggattagcag agcgaggtat gtaggcggtg ctacagagtt 6900 cttgaagtgg tggcctaact acggctacac tagaaggaca gtatttggta tctgcgctct 6960 gctgaagcca gttaccttcg gaaaaagagt tggtagctct tgatccggca aacaaaccac 7020 cgctggtagc ggtggttttt ttgtttgcaa gcagcagatt acgcgcagaa aaaaaggatc 7080 tcaagaagat cetttgatet tttetaeggg gtetgaeget eagtggaaeg aaaaeteaeg 7140 ttaagggatt ttggtcatga gattatcaaa aaggatcttc acctagatcc ttttaaatta 7200 aaaatgaagt tttaaatcaa tctaaagtat atatgagtaa acttggtctg acagttacca 7260 atgcttaatc agtgaggcac ctatctcagc gatctgtcta tttcgttcat ccatagttgc 7320 ctgactcccc gtcgtgtaga taactacgat acgggagggc ttaccatctg gccccagtgc 7380 tgcaatgata ccgcgagacc cacgctcacc ggctccagat ttatcagcaa taaaccagcc 7440 agccggaagg gccgagcgca gaagtggtcc tgcaacttta tccgcctcca tccagtctat 7500 taattgttgc cgggaagcta gagtaagtag ttcgccagtt aatagtttgc gcaacgttgt 7560 tgccattgct acaggcatcg tggtgtcacg ctcgtcgttt ggtatggctt cattcagctc 7620 cqqttcccaa cgatcaaggc gagttacatg atcccccatg ttgtgcaaaa aagcggttag 7680 ctccttcggt cctccgatcg ttgtcagaag taagttggcc gcagtgttat cactcatggt 7740 tatggcagca ctgcataatt ctcttactgt catgccatcc gtaagatgct tttctgtgac 7800 tggtgagtac tcaaccaagt cattctgaga atagtgtatg cggcgaccga gttgctcttg 7860 cccggcgtca atacgggata ataccgcgcc acatagcaga actttaaaaag tgctcatcat 7920 tggaaaacgt tcttcggggc gaaaactctc aaggatctta ccgctgttga gatccagttc 7980 gatgtaaccc actcgtgcac ccaactgatc ttcagcatct tttactttca ccagcgtttc 8040 tgggtgagca aaaacaggaa ggcaaaatgc cgcaaaaaag ggaataaggg cgacacggaa 8100 atgttgaata ctcatactct tcctttttca atattattga agcatttatc agggttattg 8160 tctcatgagc ggatacatat ttgaatgtat ttagaaaaat aaacaaatag gggttccgcg 8220 cacatttccc cgaaaagtgc cacctgacgt ctaagaaacc attattatca tgacattaac 8280 ctataaaaat aggcgtatca cgaggccctt tcgtctcgca tgcggatcag tgagggtttg 8340 caactgeggg teaaggatet ggatttegat caeggeacga teategtgeg ggagggeaag 8400 ggctccaagg atcgggcctt gatgttaccc gagagcttgg cacccagcct gcgcgagcag 8460 gggaattgat ccggtggatg accttttgaa tgacctttaa tagattatat tactaattaa 8520 ttggggaccc tagaggtccc cttttttatt ttaaaaaattt tttcacaaaa cggtttacaa 8580 gcataacggg ttttgctgcc cgcaaacggg ctgttctggt gttgctagtt tgttatcaga 8640 ategeagate eggetteagg tttgeegget gaaagegeta tttetteeag aattgeeatg 8700 attttttccc cacgggaggc gtcactggct cccgtgttgt cggcagcttt gattcgataa 8760 gcagcatege etgtttcagg etgtctatgt gtgactgttg agetgtaaca agttgtctca 8820 ggtgttcaat ttcatgttct agttgctttg ttttactggt ttcacctgtt ctattaggtg 8880 ttacatgctg ttcatctgtt acattgtcga tctgttcatg gtgaacagct ttaaatgcac 8940 caaaaactcg taaaagctct gatgtatcta tcttttttac accgttttca tctgtgcata 9000 tggacagttt tccctttgat atctaacggt gaacagttgt tctacttttg tttgttagtc 9060 ttgatgette actgatagat acaagageea taagaacete agateettee gtatttagee 9120 agtatgttet etagtgtggt tegttgtttt tgegtgagee atgagaaega accattgaga 9180 tcatgcttac tttgcatgtc actcaaaaat tttgcctcaa aactggtgag ctgaattttt 9240 gcagttaaag catcgtgtag tgtttttctt agtccgttac gtaggtagga atctgatgta 9300 atggttgttg gtattttgtc accattcatt tttatctggt tgttctcaag ttcggttacg 9360 agatecattt gtetatetag tteaacttgg aaaateaacg tateagtegg geggeetege 9420 ttatcaacca ccaatttcat attgctgtaa gtgtttaaat ctttacttat tggtttcaaa 9480 acccattggt taagcctttt aaactcatgg tagttatttt caagcattaa catgaactta 9540 aattcatcaa ggctaatctc tatatttgcc ttgtgagttt tcttttgtgt tagttctttt 9600 aataaccact cataaatcct cataqaqtat ttgttttcaa aagacttaac atgttccaga 9660 ttatatttta tgaatttttt taactggaaa agataaggca atatctcttc actaaaaact 9720 aattetaatt titegetiga gaactiggea tagiitgiee aetggaaaat eteaaageet 9780 ttaaccaaag gatteetgat tteeacagtt etegteatea getetetggt tgetttaget 9840 aatacaccat aagcattttc cctactgatg ttcatcatct gagcgtattg gttataagtg 9900 aacgataccg tccgttcttt ccttgtaggg ttttcaatcg tggggttgag tagtgccaca 9960 cagcataaaa ttagcttggt ttcatgctcc gttaagtcat agcgactaat cgctagttca 10020 tttgctttga aaacaactaa ttcagacata catctcaatt ggtctaggtg attttaatca 10080 ctataccaat tgagatgggc tagtcaatga taattactag tccttttcct ttgagttgtg 10140 ggtatctgta aattctgcta gacctttgct ggaaaacttg taaattctgc tagaccctct 10200 gtaaattccg ctagaccttt gtgtgttttt tttgtttata ttcaagtggt tataatttat 10260 agaataaaga aagaataaaa aaagataaaa agaatagatc ccagccctgt gtataactca 10320 ctactttagt cagttccgca gtattacaaa aggatgtcgc aaacgctgtt tgctcctcta 10380 caaaacagac cttaaaaccc taaaggctta agtagcaccc tcgcaagctc gggcaaatcg 10440 ctgaatattc cttttgtctc cgaccatcag gcacctgagt cgctgtcttt ttcgtgacat 10500 tcagttcgct gcgctcacgg ctctggcagt gaatgggggt aaatggcact acaggcgcct 10560 tttatggatt catgcaagga aactacccat aatacaagaa aagcccgtca cgggcttctc 10620 agggegtttt atggegggte tgetatgtgg tgetatetga etttttgetg tteageagtt 10680 cetgecetet gattttecag tetgaceaet teggattate eegtgacagg teatteagae 10740 tggctaatgc acccagtaag gcagcggtat catcaacagg cttacccgtc ttactgtcaa 10800 C 10801

<210> 77 <211> 8654

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN236 plasmid

<400> 77 ctcgaggcct acctagcttc caagaaagat atcctaacag cacaagagcg gaaagatgtt 60 ttgttctaca tccagaacaa cctctgctaa aattcctgaa aaattttgca aaaagttgtt 120 gactttatct acaaggtgtg gtataataat cttaacaaca gcaggacgct ctagaggagg 180 agacatcatg aaaattggaa ttatcggcgg aggctccgtt ggtcttttat gcgcctatta 240 tttgtcactt tatcacgacg tgactgttgt gacgaggcgg caagaacagg ctgcggccat 300 tcagtctgaa ggaatccggc tttataaagg cggggaggaa ttcagggctg attgcagtgc 360 ggacacgagt atcaattcgg actttgacct gcttgtcgtg acagtgaagc agcatcagct 420 tcaatctgtt ttttcgtcgc ttgaacgaat cgggaagacg aatatattat ttttgcaaaa 480 eggeatgggg catatecaeg acetaaaaga etggeaegtt ggeeatteca titatgitgg 540 aatcgttgag cacggagctg taagaaaatc ggatacagct gttgatcata caggcctagg 600 tgcgataaaa tggagcgcgt tcgacgatgc tgaaccagac cggctgaaca tcttgtttca 660 gcataaccat teggatttte egatttatta tgagaeggat tggtacegte tgetgaeggg 720 caagetgatt gtaaatgegt gtattaatee tttaaetgeg ttattgeaag tgaaaaatgg 780 agaactgctg acaacgccag cttatctggc ttttatgaag ctggtatttc aggaggcatg 840 ccgcatttta aaacttgaaa atgaagaaaa ggcttgggag cgggttcagg ccgtttgtgg 900 gcaaacgaaa gagaatcgtt catcaatgct ggttgacgtc attggaggcc ggcagacgga 960 agctgacgcc attatcggat acttattgaa ggaagcaagt cttcaaggtc ttgatgccgt 1020 ccacctagag tttttatatg gcagcatcaa agcattggag cgaaatacaa acaaagtctt 1080 ttgagctttt tcggtaacat gctatactca tttcggatca ctaactattt attggaggat 1140 cctgttttgg cggatgagag aagattttca gcctgataca gattaaatca gaacgcagaa 1200 gcggtctgat aaaacagaat ttgcctggcg gcagtagcgc ggtggtccca cctgacccca 1260 tgccgaactc agaagtgaaa cgccgtagcg ccgatggtag tgtggggtct ccccatgcga 1320 gagtagggaa ctgccaggca tcaaataaaa cgaaaggctc agtcgaaaga ctgggccttt 1380 cgttttatct gttgtttgtc ggtgaacgct ctcctgagta ggacaaatcc gccgggagcg 1440 gatttgaacg ttgcgaagca acggcccgga gggtggcggg caggacgccc gccataaact 1500 gccaggcatc aaattaagca gaaggccatc ctgacggatg gcctttttgc gtttctacaa 1560 actcttggta cccagaaaaa gcggcaaaag cggctgttaa aaaagcgaaa tcgaagaagc 1620 tgtctgccgc taagacggaa tatcaaaagc gttctgctgt tgtgtcatct ttaaaagtca 1680 cagccgatga atcccagcaa gatgtcctaa aatacttgaa cacccagaaa gataaaggaa 1740 atgcagacca aattcattct tattatgtgg tgaacgggat tgctgttcat gcctcaaaag 1800 aggttatgga aaaagtggtg cagtttcccg aagtggaaaa ggtgcttcct aatgagaaac 1860 ggcagctttt taagtcatcc tccccattta atatgaaaaa agcacagaaa gctattaaag 1920 caactgacgg tgtggaatgg aatgtagacc aaatcgatgc cccaaaagct tgggcacttg 1980 gatatgatgg aactggcacg gttgttgcgt ccattgatac cggggtggaa tggaatcatc 2040 cggcattaaa agagaaatat cgcggatata atccggaaaa tcctaatgag cctgaaaatg 2100 aaatgaactg gtatgatgcc gtagcaggcg aggcaagccc ttatgatgat ttggctcatg 2160 gaacccacgt gacaggcacg atggtgggct ctgaacctga tggaacaaat caaatcggtg 2220 tagcacctgg cgcaaaatgg attgctgtta aagcgttctc tgaagatggc ggcactgatg 2280 ctgacatttt ggaagctggt gaatgggttt tagcaccaaa ggacgcggaa ggaaatcccc 2340 acceggaaat ggctcctgat gttgtcaata actcatgggg agggggctct ggacttgatg 2400 aatggtacag agacatggtc aatgcctggc gttcggccga tattttccct gagttttcag 2460 cggggaatac ggatctcttt attcccggcg ggcctggttc tatcgcaaat ccggcaaact 2520 atccagaatc gtttgcaact ggagcgactg agaattccaa ttccccatgg agagaaaaga 2580 aaatcgctaa tgttgattac tttgaacttc tgcatattct tgaatttaaa aaggctgaaa 2640 gagtaaaaga ttgtgctgaa atattagagt ataaacaaaa tcgtgaaaca ggcgaaagaa 2700 agttgtatcg agtgtggttt tgtaaatcca ggctttgtcc aatgtgcaac tggaggagag 2760 caatgaaaca tggcattcag tcacaaaagg ttgttgctga agttattaaa caaaagccaa 2820 cagttcgttg gttgtttctc acattaacag ttaaaaatgt ttatgatggc gaagaattaa 2880 ataagagttt gtcagatatg gctcaaggat ttcgccgaat gatgcaatat aaaaaaatta 2940 ataaaaatct tgttggtttt atgcgtgcaa cggaagtgac aataaataat aaagataatt 3000 cttataatca gcacatgcat gtattggtat gtgtggaacc aacttatttt aagaatacag 3060 aaaactacgt gaatcaaaaa caatggattc aattttggaa aaaggcaatg aaattagact 3120 atgatccaaa tgtaaaagtt caaatgattc gaccgaaaaa taaatataaa tcggatatac 3180 aatcggcaat tgacgaaact gcaaaatatc ctgtaaagga tacggatttt atgaccgatg 3240 atgaagaaaa gaatttgaaa cgtttgtctg atttggagga aggtttacac cgtaaaaggt 3300 taatctccta tggtggtttg ttaaaagaaa tacataaaaa attaaacctt gatgacacag 3360 aagaaggcga tttgattcat acagatgatg acgaaaaagc cgatgaagat ggattttcta 3420 ttattgcaat gtggaattgg gaacggaaaa attattttat taaagagtag ttcaacaaac 3480 gggccatatt gttgtataag tgatgaaata ctgaatttaa aacttagttt atatgtggta 3540 caattaaaag agggaagcgt atcattaacc ctataaacta cgtctgccct cattattgga 3660 gggtgaaatg tgaatacatc ctattcacaa tcgaatttac gacacaacca aattttaatt 3720 tggctttgca ttttatcttt ttttagcgta ttaaatgaaa tggttttgaa cgtctcatta 3780 cetgatattg caaatgattt taataaacca cetgegagta caaactgggt gaacacagee 3840 tttatgttaa ccttttccat tggaacagct gtatatggaa agctatctga tcaattaggc 3900 atcaaaaggt tactcctatt tggaattata ataaattgtt tcgggtcggt aattgggttt 3960 gttggccatt ctttcttttc cttacttatt atggctcgtt ttattcaagg ggctggtgca 4020 gctgcatttc cagcactcgt aatggttgta gttgcgcgct atattccaaa ggaaaatagg 4080 ggtaaagcat ttggtcttat tggatcgata gtagccatgg gagaaggagt cggtccagcg 4140 attggtggaa tgatagccca ttatattcat tggtcctatc ttctactcat tcctatgata 4200 acaattatca ctgttccgtt tcttatgaaa ttattaaaga aagaagtaag gataaaaggt 4260 cattttgata tcaaaggaat tatactaatg tctgtaggca ttgtattttt tatgttgttt 4320 acaacatcat atagcatttc ttttcttatc gttagcgtgc tgtcattcct gatatttgta 4380 aaacatatca ggaaagtaac agatcctttt gttgatcccg gattagggaa aaatatacct 4440 tttatgattg gagttctttg tgggggaatt atatttggaa cagtagcagg gtttgtctct 4500 atggttcctt atatgatgaa agatgttcac cagctaagta ctgccgaaat cggaagtgta 4560 attattttcc ctggaacaat gagtgtcatt attttcggct acattggtgg gatacttgtt 4620 gatagaagag gtcctttata cgtgttaaac atcggagtta catttctttc tgttagcttt 4680 ttaactgctt cctttctttt agaaacaaca tcatggttca tgacaattat aatcgtattt 4740 gttttaggtg ggctttcgtt caccaaaaca gttatatcaa caattgtttc aagtagcttg 4800 aaacagcagg aagctggtgc tggaatgagt ttgcttaact ttaccagctt tttatcagag 4860 ggaacaggta ttgcaattgt aggtggttta ttatccatac ccttacttga tcaaaggttg 4920 ttacctatgg aagttgatca gtcaacttat ctgtatagta atttgttatt acttttttca 4980 ggaatcattg tcattagttg gctggttacc ttgaatgtat ataaacattc tcaaagggat 5040 ttctaaatcg ttaagggatc aactttggga gagagttcaa aattgatcct ttttttataa 5100 cagttcgaag cggccgcaat tcttgaagac gaaagggcct cgtgatacgc ctattttat 5160 aggttaatgt catgataata atggtttett agacgteagg tggcaetttt eggggaaatg 5220 tgcgcggaac ccctatttgt ttatttttct aaatacattc aaatatgtat ccgctcatga 5280 gacaataacc ctgataaatg cttcaataat attgaaaaag gaagagtatg agtattcaac 5340 atttccgtgt cgcccttatt cccttttttg cggcattttg ccttcctgtt tttgctcacc 5400 cagaaacgct ggtgaaagta aaagatgctg aagatcagtt gggtgcacga gtgggttaca 5460 tcgaactgga tctcaacagc ggtaagatcc ttgagagttt tcgccccgaa gaacgttttc 5520 caatgatgag cacttttaaa gttctgctat gtggcgcggt attatcccgt attgacgccg 5580 ggcaagagca actcggtcgc cgcatacact attctcagaa tgacttggtt gagtactcac 5640 cagtcacaga aaagcatctt acggatggca tgacagtaag agaattatgc agtgctgcca 5700 taaccatgag tgataacact gcggccaact tacttctgac aacgatcgga ggaccgaagg 5760 agctaaccgc ttttttgcac aacatggggg atcatgtaac tcgccttgat cgttgggaac 5820 cggagctgaa tgaagccata ccaaacgacg agcgtgacac cacgatgcct gcagcaatgg 5880 caacaacgtt gcgcaaacta ttaactggcg aactacttac tctagcttcc cggcaacaat 5940 taatagactg gatggaggcg gataaagttg caggaccact tctgcgctcg gcccttccgg 6000 ctggctggtt tattgctgat aaatctggag ccggtgagcg tgggtctcgc ggtatcattg 6060 cagcactggg gccagatggt aagccctccc gtatcgtagt tatctacacg acggggagtc 6120 aggcaactat ggatgaacga aatagacaga tcgctgagat aggtgcctca ctgattaagc 6180 attggtaact gtcagaccaa gtttactcat atatacttta gattgattta aaacttcatt 6240 tttaatttaa aaggatctag gtgaagatcc tttttgataa tctcatgacc aaaatccctt 6300 aacgtgagtt ttcgttccac tgagcgtcag accccgtaga aaagatcaaa ggatcttctt 6360 gagateettt ttttetgege gtaatetget gettgeaaac aaaaaaacca eegetaecag 6420 cggtggtttg tttgccggat caagagctac caactctttt tccgaaggta actggcttca 6480 gcagagegea gataceaaat actgteette tagtgtagee gtagttagge caccaettea 6540 agaactetgt agcacegeet acataceteg etetgetaat eetgttaeea gtggetgetg 6600 ccagtggcga taagtcgtgt cttaccgggt tggactcaag acgatagtta ccggataagg 6660 cgcagcggtc gggctgaacg gggggttcgt gcacacagcc cagcttggag cgaacgacct 6720 acaccgaact gagataccta cagcgtgagc tatgagaaag cgccacgctt cccgaaggga 6780 gaaaggegga caggtateeg gtaageggea gggteggaac aggagagege aegagggage 6840 ttccaggggg aaacgcctgg tatctttata gtcctgtcgg gtttcgccac ctctgacttg 6900 agcgtcgatt tttgtgatgc tcgtcagggg ggcggagcct atggaaaaac gccagcaacg 6960 eggeettttt aeggtteetg geettttget ggeettttge teacatgtte ttteetgegt 7020 tatcccctga ttctgtggat aaccgtatta ccgcctttga gtgagctgat accgctcgcc 7080 gcagccgaac gaccgagcgc agcgagtcag tgagcgagga agcggaagag cgcctgatgc 7140 ggtattttct ccttacgcat ctgtgcggta tttcacaccg catatggtgc actctcagta 7200 caatctgctc tgatgccgca tagttaagcc agtatacact ccgctatcgc tacgtgactg 7260 ggtcatggct gcgccccgac acccgccaac acccgctgac gcgccctgac gggcttgtct 7320 gctcccggca tccgcttaca gacaagctgt gaccgtctcc gggagctgca tgtgtcagag 7380 gttttcaccg tcatcaccga aacgcgcgag gcagctgcgg taaagctcat cagcgtggtc 7440 gtgaagcgat tcacagatgt ctgcctgttc atccgcgtcc agctcgttga gtttctccag 7500 aagcgttaat gtctggcttc tgataaagcg ggccatgtta agggcggttt tttcctgttt 7560 ggtcacttga tgcctccgtg taagggggaa tttctgttca tgggggtaat gataccgatg 7620 aaacgagaga ggatgctcac gatacgggtt actgatgatg aacatgcccg gttactggaa 7680 cgttgtgagg gtaaacaact ggcggtatgg atgcggcggg accagagaaa aatcactcag 7740 ggtcaatgcc agcgcttcgt taatacagat gtaggtgttc cacagggtag ccagcagcat 7800 cetgcgatgc agateeggaa cataatggtg cagggegetg actteegegt ttecagaett 7860 tacgaaacac ggaaaccgaa gaccattcat gttgttgctc aggtcgcaga cgttttgcag 7920 cagcagtege tteaegtteg etegegtate ggtgatteat tetgetaace agtaaggeaa 7980 ccccgccagc ctagccgggt cctcaacgac aggagcacga tcatgcgcac ccgtggccag 8040 gacccaacgc tgcccgagat gcgccgcgtg cggctgctgg agatggcgga cgcgatggat 8100 atgttctgcc aagggttggt ttgcgcattc acagttctcc gcaagaattg attggctcca 8160 attcttggag tggtgaatcc gttagcgagg tgccgccggc ttccattcag gtcgaggtgg 8220 cccggctcca tgcaccgcga cgcaacgcgg ggaggcagac aaggtatagg gcggcgccta 8280 caatccatgc caacccgttc catgtgctcg ccgaggcggc ataaatcgcc gtgacgatca 8340 geggtecagt gategaagtt aggetggtaa gageegegag egateettga agetgteeet 8400 gatggtcgtc atctacctgc ctggacagca tggcctgcaa cgcgggcatc ccgatgccgc 8460 cggaagcgag aagaatcata atggggaagg ccatccagcc tcgcgtcggc ggccgcttcg 8520 tcgaccgaaa cagcagttat aaggcatgaa gctgtccggt ttttgcaaaa gtggctgtga 8580 ctgtaaaaag aaatcgaaaa agaccgtttt gtgtgaaaac ggtctttttg tttcctttta 8640 8654 accaactgcc ataa

<210> 78

<211> 8093

<212> DNA

<213> Artificial Sequence

<223> Description of Artificial Sequence: Recombinant pAN423 plasmid

<400> 78 ggcggccgct tcgtcgaccg aaacagcagt tataaggcat gaagctgtcc ggtttttgca 60 aaagtggctg tgactgtaaa aagaaatcga aaaagaccgt tttgtgtgaa aacggtcttt 120 ttgtttcctt ttaaccaact gccataactc gaggcctacc tagcttccaa gaaagatatc 180 ctaacagcac aagageggaa agatgttttg ttetacatee agaacaacet etgetaaaat 240 teetgaaaaa ttttgeaaaa agttgttgae tttatetaea aggtgtggta taataatett 300 aacaacagca ggacgctcta gaaaaggagg aatttaaatg tatcgtacga tgatgagcgg 360 caaacttcac agggcaactg ttacggaagc aaacctgaac tatgtgggaa gcattacaat 420 tgatgaagat ctcattgatg ctgtgggaat gcttcctaat gaaaaagtac aaattgtgaa 480 taataataat ggagcacgtc ttgaaacgta tattattcct ggtaaacggg gaagcggcgt 540 catatgctta aacggtgcag ccgcacgcct tgtgcaggaa ggagataagg tcattattat 600 ttcctacaaa atgatgtctg atcaagaagc ggcaagccat gagccgaaag tggctgttct 660 gaatgatcaa aacaaaattg aacaaatgct ggggaacgaa ccagcccgta caattttgta 720 aaggateetg ttttggegga tgagagaaga tttteageet gatacagatt aaateagaac 780 gcagaagcgg tctgataaaa cagaatttgc ctggcggcag tagcgcggtg gtcccacctg 840 accccatgcc gaactcagaa gtgaaacgcc gtagcgccga tggtagtgtg gggtctcccc 900 atgcgagagt agggaactgc caggcatcaa ataaaacgaa aggctcagtc gaaagactgg 960 gcctttcgtt ttatctgttg tttgtcggtg aacgctctcc tgagtaggac aaatccgccg 1020 ggageggatt tgaaegttge gaageaaegg eeeggagggt ggegggeagg aegeeegeea 1080 taaactgcca ggcatcaaat taagcagaag gccatcctga cggatggcct ttttgcgttt 1140 ctacaaactc ttggtaccca gaaaaagcgg caaaagcggc tgttaaaaaa gcgaaatcga 1200 agaagctgtc tgccgctaag acggaatatc aaaagcgttc tgctgttgtg tcatctttaa 1260 aagtcacagc cgatgaatcc cagcaagatg tcctaaaata cttgaacacc cagaaagata 1320 aaggaaatgc agaccaaatt cattcttatt atgtggtgaa cgggattgct gttcatgcct 1380 caaaagaggt tatggaaaaa gtggtgcagt ttcccgaagt ggaaaaggtg cttcctaatg 1440 agaaacggca gctttttaag tcatcctccc catttaatat gaaaaaagca cagaaagcta 1500 ttaaagcaac tgacggtgtg gaatggaatg tagaccaaat cgatgcccca aaagcttggg 1560 cacttggata tgatggaact ggcacggttg ttgcgtccat tgataccggg gtggaatgga 1620 atcatccggc attaaaagag aaatatcgcg gatataatcc ggaaaatcct aatgagcctg 1680 aaaatgaaat gaactggtat gatgccgtag caggcgaggc aagcccttat gatgatttgg 1740 ctcatggaac ccacgtgaca ggcacgatgg tgggctctga acctgatgga acaaatcaaa 1800 teggtgtage acctggcgca aaatggattg ctgttaaage gttetetgaa gatggeggca 1860 ctgatgctga cattttggaa gctggtgaat gggttttagc accaaaggac gcggaaggaa 1920 atccccaccc ggaaatggct cctgatgttg tcaataactc atggggaggg ggctctggac 1980 ttgatgaatg gtacagagac atggtcaatg cctggcgttc ggccgatatt ttccctgagt 2040 tttcagcggg gaatacggat ctctttattc ccggcgggcc tggttctatc gcaaatccgg 2100 caaactatcc agaatcgttt gcaactggag cgactgagaa ttccaattcc ccatggagag 2160 aaaagaaaat cgctaatgtt gattactttg aacttctgca tattcttgaa tttaaaaagg 2220 ctgaaagagt aaaagattgt gctgaaatat tagagtataa acaaaatcgt gaaacaggcg 2280 aaagaaagtt gtatcgagtg tggttttgta aatccaggct ttgtccaatg tgcaactgga 2340 ggagagcaat gaaacatggc attcagtcac aaaaggttgt tgctgaagtt attaaacaaa 2400 agccaacagt tcgttggttg tttctcacat taacagttaa aaatgtttat gatggcgaag 2460 aattaaataa gagtttgtca gatatggctc aaggatttcg ccgaatgatg caatataaaa 2520 aaattaataa aaatcttgtt ggttttatgc gtgcaacgga agtgacaata aataataaag 2580 ataattetta taateageae atgeatgtat tggtatgtgt ggaaceaaet tattttaaga 2640 atacagaaaa ctacgtgaat caaaaacaat ggattcaatt ttggaaaaaag gcaatgaaat 2700 tagactatga tccaaatgta aaagttcaaa tgattcgacc gaaaaataaa tataaatcgg 2760 atatacaatc ggcaattgac gaaactgcaa aatatcctgt aaaggatacg gattttatga 2820 ccgatgatga agaaaagaat ttgaaacgtt tgtctgattt ggaggaaggt ttacaccgta 2880 aaaggttaat ctcctatggt ggtttgttaa aagaaataca taaaaaatta aaccttgatg 2940 acacagaaga aggcgatttg attcatacag atgatgacga aaaagccgat gaagatggat 3000 tttctattat tgcaatgtgg aattgggaac ggaaaaatta ttttattaaa gagtagttca 3060 acaaacgggc catattgttg tataagtgat gaaatactga atttaaaact tagtttatat 3120 gtggtaaaat gttttaatca agtttaggag gaattaatta tgaagtgtaa tgaatgtaac 3180 agggttcaat taaaagaggg aagcgtatca ttaaccctat aaactacgtc tgccctcatt 3240 attggagggt gaaatgtgaa tacatcctat tcacaatcga atttacgaca caaccaaatt 3300 ttaatttggc tttgcatttt atctttttt agcgtattaa atgaaatggt tttgaacgtc 3360 tcattacctg atattgcaaa tgattttaat aaaccacctg cgagtacaaa ctgggtgaac 3420 acagcettta tgttaacett tteeattgga acagetgtat atggaaaget atetgateaa 3480 ttaggcatca aaaggttact cctatttgga attataataa attgtttcgg gtcggtaatt 3540 gggtttgttg gccattcttt cttttcctta cttattatgg ctcgttttat tcaaggggct 3600 ggtgcagctg catttccagc actcgtaatg gttgtagttg cgcgctatat tccaaaggaa 3660 aataggggta aagcatttgg tcttattgga tcgatagtag ccatgggaga aggagtcggt 3720 ccagegattg gtggaatgat ageceattat atteattggt cetatettet acteatteet 3780 atgataacaa ttatcactgt tccgtttctt atgaaattat taaagaaaga agtaaggata 3840 aaaggtcatt ttgatatcaa aggaattata ctaatgtctg taggcattgt attttttatg 3900 ttgtttacaa catcatatag catttctttt cttatcgtta gcgtgctgtc attcctgata 3960 tttgtaaaac atatcaggaa agtaacagat ccttttgttg atcccggatt agggaaaaat 4020 atacctttta tgattggagt tctttgtggg ggaattatat ttggaacagt agcagggttt 4080 gtctctatgg ttccttatat gatgaaagat gttcaccagc taagtactgc cgaaatcgga 4140 agtgtaatta ttttccctgg aacaatgagt gtcattattt tcggctacat tggtgggata 4200 cttgttgata gaagaggtcc tttatacgtg ttaaacatcg gagttacatt tctttctgtt 4260 agetttttaa etgetteett tettttagaa acaacateat ggtteatgae aattataate 4320 gtatttgttt taggtgggct ttcgttcacc aaaacagtta tatcaacaat tgtttcaagt 4380 agettgaaac ageaggaage tggtgetgga atgagtttge ttaactttae cagettttta 4440 tcagagggaa caggtattgc aattgtaggt ggtttattat ccataccctt acttgatcaa 4500 aggttgttac ctatggaagt tgatcagtca acttatctgt atagtaattt gttattactt 4560 ttttcaggaa tcattgtcat tagttggctg gttaccttga atgtatataa acattctcaa 4620 agggatttet aaategttaa gggateaact ttgggagaga gtteaaaatt gateettttt 4680 ttataacagt tcgaagcggc cgcaattctt gaagacgaaa gggcctcgtg atacgcctat 4740 ttttataggt taatgtcatg ataataatgg tttcttagac gtcaggtggc acttttcggg 4800 gaaatgtgcg cggaacccct atttgtttat ttttctaaat acattcaaat atgtatccgc 4860 tcatgagaca ataaccctga taaatgcttc aataatattg aaaaaggaag agtatgagta 4920 ttcaacattt ccgtgtcgcc cttattccct tttttgcggc attttgcctt cctgtttttg 4980 ctcacccaga aacgctggtg aaagtaaaag atgctgaaga tcagttgggt gcacgagtgg 5040gttacatcga actggatctc aacagcggta agatccttga gagttttcgc cccgaagaac 5100 gttttccaat gatgagcact tttaaagttc tgctatgtgg cgcggtatta tcccgtattg 5160 acgccgggca agagcaactc ggtcgccgca tacactattc tcagaatgac ttggttgagt 5220 actcaccagt cacagaaaag catcttacgg atggcatgac agtaagagaa ttatgcagtg 5280 ctgccataac catgagtgat aacactgcgg ccaacttact tctgacaacg atcggaggac 5340 cgaaggagct aaccgctttt ttgcacaaca tgggggatca tgtaactcgc cttgatcgtt 5400 gggaaccgga gctgaatgaa gccataccaa acgacgagcg tgacaccacg atgcctgcag 5460 caatggcaac aacgttgcgc aaactattaa ctggcgaact acttactcta gcttcccggc 5520 aacaattaat agactggatg gaggcggata aagttgcagg accacttctg cgctcggccc 5580 ttccggctgg ctggtttatt gctgataaat ctggagccgg tgagcgtggg tctcgcggta 5640 tcattgcagc actggggcca gatggtaagc cctcccgtat cgtagttatc tacacgacgg 5700 ggagtcaggc aactatggat gaacgaaata gacagatcgc tgagataggt gcctcactga 5760 ttaagcattg gtaactgtca gaccaagttt actcatatat actttagatt gatttaaaac 5820 ttcattttta atttaaaagg atctaggtga agatcctttt tgataatctc atgaccaaaa 5880 tecettaaeg tgagtttteg ttecaetgag egteagaece egtagaaaag ateaaaggat 5940 cttcttgaga tcctttttt ctgcgcgtaa tctgctgctt gcaaacaaaa aaaccaccgc 6000 taccageggt ggtttgtttg ceggateaag agetaceaac tetttteeg aaggtaactg 6060 getteageag agegeagata ceaaatactg teettetagt gtageegtag ttaggeeace 6120 acttcaagaa ctctgtagca ccgcctacat acctcgctct gctaatcctg ttaccagtgg 6180 ctgctgccag tggcgataag tcgtgtctta ccgggttgga ctcaagacga tagttaccgg 6240 ataaggegea geggteggge tgaacggggg gttegtgeac acageceage ttggagegaa 6300 cgacctacac cgaactgaga tacctacagc gtgagctatg agaaagcgcc acgcttcccg 6360 aagggagaaa ggcggacagg tatccggtaa gcggcagggt cggaacagga gagcgcacga 6420 gggagettee agggggaaac geetggtate tttatagtee tgtegggttt egeeacetet 6480 gacttgagcg tcgatttttg tgatgctcgt caggggggcg gagcctatgg aaaaacgcca 6540 gcaacgcggc ctttttacgg ttcctggcct tttgctggcc ttttgctcac atgttctttc 6600 ctgcgttatc ccctgattct gtggataacc gtattaccgc ctttgagtga gctgataccg 6660 ctcgccgcag ccgaacgacc gagcgcagcg agtcagtgag cgaggaagcg gaagagcgcc 6720 tgatgcggta ttttctcctt acgcatctgt gcggtatttc acaccgcata tggtgcactc 6780 tcagtacaat ctgctctgat gccgcatagt taagccagta tacactccgc tatcgctacg 6840 tgactgggtc atggctgcgc cccgacaccc gccaacaccc gctgacgcgc cctgacgggc 6900 ttgtctgctc ccggcatccg cttacagaca agctgtgacc gtctccggga gctgcatgtg 6960 tcagaggttt tcaccgtcat caccgaaacg cgcgaggcag ctgcggtaaa gctcatcagc 7020 gtggtcgtga agcgattcac agatgtctgc ctgttcatcc gcgtccagct cgttgagttt 7080 ctccagaagc gttaatgtct ggcttctgat aaagcgggcc atgttaaggg cggttttttc 7140 ctgtttggtc acttgatgcc tccgtgtaag ggggaatttc tgttcatggg ggtaatgata 7200 ccgatgaaac gagagaggat gctcacgata cgggttactg atgatgaaca tgcccggtta 7260 ctggaacgtt gtgagggtaa acaactggcg gtatggatgc ggcgggacca gagaaaaatc 7320 actcagggtc aatgccagcg cttcgttaat acagatgtag gtgttccaca gggtagccag 7380 cagcatectg egatgeagat eeggaacata atggtgeagg gegetgaett eegegtttee 7440 agactttacg aaacacggaa accgaagacc attcatgttg ttgctcaggt cgcagacgtt 7500 ttgcagcagc agtcgcttca cgttcgctcg cgtatcggtg attcattctg ctaaccagta 7560 aggcaacccc gccagcctag ccgggtcctc aacgacagga gcacgatcat gcgcacccgt 7620 ggccaggacc caacgctgcc cgagatgcgc cgcgtgcggc tgctggagat ggcggacgcg 7680 atggatatgt tetgeeaagg gttggtttge geatteaeag tteteegeaa gaattgattg 7740 gctccaattc ttggagtggt gaatccgtta gcgaggtgcc gccggcttcc attcaggtcg 7800 aggtggcccg gctccatgca ccgcgacgca acgcggggag gcagacaagg tatagggcgg 7860 cgcctacaat ccatgccaac ccgttccatg tgctcgccga ggcggcataa atcgccgtga 7920 cgatcagcgg tccagtgatc gaagttaggc tggtaagagc cgcgagcgat ccttgaagct 7980 gtccctgatg gtcgtcatct acctgcctgg acagcatggc ctgcaacgcg ggcatcccga 8040 tgccgccgga agcgagaaga atcataatgg ggaaggccat ccagcctcgc gtc 8093

<220>

<223> Description of Artificial Sequence: Recombinant pAN429 plasmid

<400> 79
ggcggccgcttcgtcgaccgaaacagcagttataaggcatgaagetgteeggtttttgaa60aaagtggctgtgactgtaaaaagaaatcgaaaaagaccgttttgtgtgaaaacggtcttt120ttgtttccttttaaccaactgccataactcgaggcctacctagcttccaagaaagatatc180ctaacagcacaagagcggaaagatgtttgttctacatccagaacaacctctgctaaaat240tcctgaaaaattttgcaaaaagttgttgactttatctacaaggtgtgtataataatctt300aacaacagcaggacgctctagattagaaaggaggtttaattaatgtatcgtacgatgatg360agcggcaaacttcacagggcaactgttacggaagcaaacctgaactatgtgggaagcatt420acaattgatgaagatctcattgatgctgtgggaatgcttcctaatgaaaaagtacaaatt480gtgaataataataatggagcacgtcttgaaacgtatattattcctggtaaacggggaagc540ggcgtcatatgcttaaacggtgcagccgcacgccttgtgaggaaggagataaggtcatt600

<210> 79 <211> 8098 <212> DNA

<213> Artificial Sequence

attatttcct acaaaatgat qtctgatcaa gaagcggcaa gccatgagcc gaaagtggct 660 gttctgaatg atcaaaacaa aattgaacaa atgctgggga acgaaccagc ccgtacaatt 720 ttgtaaagga tcctgttttg gcggatgaga gaagattttc agcctgatac agattaaatc 780 agaacgcaga agcggtctga taaaacagaa tttgcctggc ggcagtagcg cggtggtccc 840 acctgacccc atgccgaact cagaagtgaa acgccgtagc gccgatggta gtgtggggtc 900 tececatgeg agagtaggga aetgeeagge ateaaataaa aegaaagget eagtegaaag 960 actgggcctt tcgttttatc tgttgtttgt cggtgaacgc tctcctgagt aggacaaatc 1020 cgccgggagc ggatttgaac gttgcgaagc aacggcccgg agggtggcgg gcaggacgcc 1080 cgccataaac tgccaggcat caaattaagc agaaggccat cctgacggat ggcctttttg 1140 cgtttctaca aactcttggt acccagaaaa agcggcaaaa gcggctgtta aaaaagcgaa 1200 atcgaagaag ctgtctgccg ctaagacgga atatcaaaag cgttctgctg ttgtgtcatc 1260 tttaaaagtc acagccgatg aatcccagca agatgtccta aaatacttga acacccagaa 1320 agataaagga aatgcagacc aaattcattc ttattatgtg gtgaacggga ttgctgttca 1380 tgcctcaaaa gaggttatgg aaaaagtggt gcagtttccc gaagtggaaa aggtgcttcc 1440 taatgagaaa cggcagcttt ttaagtcatc ctccccattt aatatgaaaa aagcacagaa 1500 agctattaaa gcaactgacg gtgtggaatg gaatgtagac caaatcgatg ccccaaaagc 1560 ttgggcactt ggatatgatg gaactggcac ggttgttgcg tccattgata ccggggtgga 1620 atggaatcat ccggcattaa aagagaaata tcgcggatat aatccggaaa atcctaatga 1680 gcctgaaaat gaaatgaact ggtatgatgc cgtagcaggc gaggcaagcc cttatgatga 1740 tttggctcat ggaacccacg tgacaggcac gatggtgggc tctgaacctg atggaacaaa 1800 tcaaatcggt gtagcacctg gcgcaaaatg gattgctgtt aaagcgttct ctgaagatgg 1860 cggcactgat gctgacattt tggaagctgg tgaatgggtt ttagcaccaa aggacgcgga 1920 aggaaatccc cacceggaaa tggctcctga tgttgtcaat aactcatggg gagggggctc 1980 tggacttgat gaatggtaca gagacatggt caatgcctgg cgttcggccg atattttccc 2040 tgagttttca geggggaata eggatetett tatteeegge gggeetggtt etategeaaa 2100 tccggcaaac tatccagaat cgtttgcaac tggagcgact gagaattcca attccccatg 2160 gagagaaaag aaaatcgcta atgttgatta ctttgaactt ctgcatattc ttgaatttaa 2220 aaaggctgaa agagtaaaag attgtgctga aatattagag tataaacaaa atcgtgaaac 2280 aggcgaaaga aagttgtatc gagtgtggtt ttgtaaatcc aggctttgtc caatgtgcaa 2340 ctggaggaga gcaatgaaac atggcattca gtcacaaaag gttgttgctg aagttattaa 2400 acaaaagcca acagttcgtt ggttgtttct cacattaaca gttaaaaatg tttatgatgg 2460

cgaagaatta aataagagtt tgtcagatat ggctcaagga tttcgccgaa tgatgcaata 2520 taaaaaaatt aataaaaatc ttgttggttt tatgcgtgca acggaagtga caataaataa 2580 taaagataat tettataate ageaeatgea tgtattggta tgtgtggaae caaettattt 2640 taagaataca gaaaactacg tgaatcaaaa acaatggatt caattttgga aaaaggcaat 2700 gaaattagac tatgatccaa atgtaaaagt tcaaatgatt cgaccgaaaa ataaatataa 2760 atcggatata caatcggcaa ttgacgaaac tgcaaaatat cctgtaaagg atacggattt 2820 tatgaccgat gatgaagaaa agaatttgaa acgtttgtct gatttggagg aaggtttaca 2880 ccgtaaaagg ttaatctcct atggtggttt gttaaaagaa atacataaaa aattaaacct 2940 tgatgacaca gaagaaggcg atttgattca tacagatgat gacgaaaaag ccgatgaaga 3000 tggattttct attattgcaa tgtggaattg ggaacggaaa aattattta ttaaagagta 3060 gttcaacaaa cgggccatat tgttgtataa gtgatgaaat actgaattta aaacttagtt 3120 tatatgtggt aaaatgtttt aatcaagttt aggaggaatt aattatgaag tgtaatgaat 3180 gtaacagggt tcaattaaaa gagggaageg tatcattaac cctataaact acgtctgccc 3240 tcattattgg agggtgaaat gtgaatacat cctattcaca atcgaattta cgacacaacc 3300 aaattttaat ttggctttgc attttatctt tttttagcgt attaaatgaa atggttttga 3360 acgteteatt acetgatatt geaaatgatt ttaataaace acetgegagt acaaactggg 3420 tgaacacage etttatgtta acetttteea ttggaacage tgtatatgga aagetatetg 3480 atcaattagg catcaaaagg ttactcctat ttggaattat aataaattgt ttcgggtcgg 3540 taattgggtt tgttggccat tctttctttt ccttacttat tatggctcqt tttattcaaq 3600 gggctggtgc agctgcattt ccagcactcg taatggttgt agttgcgcgc tatattccaa 3660 aggaaaatag gggtaaagca tttggtctta ttggatcgat agtagccatg ggagaaggag 3720 teggteeage gattggtgga atgatageee attatattea ttggteetat ettetaetea 3780 ttoctatgat aacaattato actgttoogt ttottatgaa attattaaag aaagaagtaa 3840 ggataaaagg tcattttgat atcaaaggaa ttatactaat gtctgtaggc attgtatttt 3900 ttatgttgtt tacaacatca tatagcattt cttttcttat cgttagcgtg ctgtcattcc 3960 tgatatttgt aaaacatatc aggaaagtaa cagatccttt tgttgatccc ggattaggga 4020 aaaatatacc ttttatgatt ggagttettt gtgggggaat tatatttgga acagtagcag 4080 ggtttgtctc tatggttcct tatatgatga aagatgttca ccagctaagt actgccgaaa 4140 teggaagtgt aattatttte eetggaacaa tgagtgteat tattttegge tacattggtg 4200 ggatacttgt tgatagaaga ggtcctttat acgtgttaaa catcggagtt acatttcttt 4260 ctgttagctt tttaactgct tcctttcttt tagaaacaac atcatggttc atgacaatta 4320 taatcgtatt tgttttaggt gggctttcgt tcaccaaaac agttatatca acaattgttt 4380 caagtagett gaaacagcag gaagetggtg etggaatgag tttgettaac tttaccaget 4440 ttttatcaga gggaacaggt attgcaattg taggtggttt attatccata cccttacttg 4500 atcaaaggtt gttacctatg gaagttgatc agtcaactta tctgtatagt aatttgttat 4560 tacttttttc aggaatcatt gtcattagtt ggctggttac cttgaatgta tataaacatt 4620 ctcaaaggga tttctaaatc gttaagggat caactttggg agagagttca aaattgatcc 4680 tttttttata acagttcgaa gcggccgcaa ttcttgaaga cgaaagggcc tcgtgatacg 4740 cctattttta taggttaatg tcatgataat aatggtttct tagacgtcag gtggcacttt 4800 tcggggaaat gtgcgcggaa cccctatttg tttatttttc taaatacatt caaatatgta 4860 tccgctcatg agacaataac cctgataaat gcttcaataa tattgaaaaa ggaagagtat 4920 gagtatteaa cattteegtg tegecettat teeetttttt geggeatttt geetteetgt 4980 ttttgctcac ccagaaacgc tggtgaaagt aaaagatgct gaagatcagt tgggtgcacg 5040 agtgggttac atcgaactgg atctcaacag cggtaagatc cttgagagtt ttcgccccga 5100 agaacgtttt ccaatgatga gcacttttaa agttctgcta tgtggcgcgg tattatcccg 5160 tattgacgcc gggcaagagc aactcggtcg ccgcatacac tattctcaga atgacttggt 5220 tgagtactca ccagtcacag aaaagcatct tacggatggc atgacagtaa gagaattatg 5280 cagtgctgcc ataaccatga gtgataacac tgcggccaac ttacttctga caacgatcgg 5340 aggaccgaag gagctaaccg cttttttgca caacatgggg gatcatgtaa ctcgccttga 5400 tcgttgggaa ccggagctga atgaagccat accaaacqac gagcqtgaca ccacqatqcc 5460 tgcagcaatg gcaacaacgt tgcgcaaact attaactggc gaactactta ctctagcttc 5520 ccggcaacaa ttaatagact ggatggaggc ggataaagtt gcaggaccac ttctgcgctc 5580 ggcccttccg gctggctggt ttattgctga taaatctgga gccggtgagc gtgggtctcg 5640 eggtateatt geageactgg ggeeagatgg taageeetee egtategtag ttatetaeae 5700 gacggggagt caggcaacta tggatgaacg aaatagacag atcgctgaga taggtgcctc 5760 actgattaag cattggtaac tgtcagacca agtttactca tatatacttt agattgattt 5820 aaaacttcat ttttaattta aaaggatcta ggtgaagatc ctttttgata atctcatgac 5880 caaaatccct taacgtgagt tttcgttcca ctgagcgtca gaccccgtag aaaagatcaa 5940 aggatettet tgagateett tttttetgeg egtaatetge tgettgeaaa eaaaaaaace 6000 acegetacea geggtggttt gtttgeegga teaagageta eeaactettt tteegaaggt 6060 aactggette ageagagege agataceaaa tactgteett etagtgtage egtagttagg 6120

ccaccacttc aagaactctg tagcaccgcc tacatacctc gctctgctaa tcctgttacc 6180 agtggctgct gccagtggcg ataagtcgtg tcttaccggg ttggactcaa gacgatagtt 6240 accggataag gegeageggt egggetgaac ggggggtteg tgeacacage ceagettgga 6300 gcgaacgacc tacaccgaac tgagatacct acagcgtgag ctatgagaaa gcgccacgct 6360 teeegaaqgg aqaaaggegg acaqgtatee ggtaagegge agggteggaa eaggagageg 6420 cacqaqqqaq cttccaqqqq qaaacqcctg gtatctttat agtcctgtcg ggtttcgcca 6480 cctctgactt gagcgtcgat ttttgtgatg ctcgtcaggg gggcggagcc tatggaaaaa 6540 cgccagcaac gcggcctttt tacggttcct ggccttttgc tggccttttg ctcacatgtt 6600 ctttcctgcg ttatcccctg attctgtgga taaccgtatt accgcctttg agtgagctga 6660 taccgctcgc cgcagccgaa cgaccgagcg cagcgagtca gtgagcgagg aagcggaaga 6720 gegeetgatg eggtatttte teettaegea tetgtgeggt attteacaee geatatggtg 6780 cacteteagt acaatetget etgatgeege atagttaage cagtatacae teegetateg 6840 ctacgtgact gggtcatggc tgcgccccga cacccgccaa cacccgctga cgcgccctga 6900 egggettgte tgeteeegge ateegettae agacaagetg tgaeegtete egggagetge 6960 atgtgtcaga ggttttcacc gtcatcaccg aaacgcgcga ggcagctgcg gtaaagctca 7020 teagegtggt egtgaagega tteacagatg tetgeetgtt cateegegte cagetegttg 7080 agtttctcca gaagcgttaa tgtctggctt ctgataaagc gggccatgtt aagggcggtt 7140 ttttcctgtt tggtcacttg atgcctccgt gtaaggggga atttctgttc atgggggtaa 7200 tgataccgat gaaacgagag aggatgctca cgatacgggt tactgatgat gaacatgccc 7260 ggttactgga acgttgtgag ggtaaacaac tggcggtatg gatgcggcgg gaccagagaa 7320 aaatcactca gggtcaatgc cagcgcttcg ttaatacaga tgtaggtgtt ccacagggta 7380 gccagcagca tcctgcgatg cagatccgga acataatggt gcagggcgct gacttccgcg 7440 tttccagact ttacgaaaca cggaaaccga agaccattca tgttgttgct caggtcgcag 7500 acgttttgca gcagcagtcg cttcacgttc gctcgcgtat cggtgattca ttctgctaac 7560 cagtaaggca accccgccag cctagccggg tcctcaacga caggagcacg atcatgcgca 7620 cccgtggcca ggacccaacg ctgcccgaga tgcgccgcgt gcggctgctg gagatggcgg 7680 acgcgatgga tatgttctgc caagggttgg tttgcgcatt cacagttctc cgcaagaatt 7740 gattggctcc aattcttgga gtggtgaatc cgttagcgag gtgccgccgg cttccattca 7800 ggtcgaggtg gcccggctcc atgcaccgcg acgcaacgcg gggaggcaga caaggtatag 7860 ggcggcgcct acaatccatg ccaacccgtt ccatgtgctc gccgaggcgg cataaatcgc 7920

cgtgacgatc agcggtccag tgatcgaagt taggctggta agagccgcaa gcgatccttg 7980 aagctgtccc tgatggtcgt catctacctg cctggacagc atggcctgca acgcgggcat 8040 cccgatgccg ccggaagcga gaagaatcat aatggggaag gccatccagc ctcgcgtc 8098

<210> 80

<211> 4450

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN443 plasmid

<400> 80 ccatcgaatg gccagatgat taattcctaa tttttgttga cactctatca ttgatagagt 60 tattttacca ctccctatca gtgatagaga aaagtgaaat gaatagttcg acaaaaatct 120 agattagaaa ggaggattta aatatgagac agattactga tatttcacag ctgaaagaag 180 ccataaaaca ataccattca gagggcaagt caatcggatt tgttccgacg atggggtttc 240 tgcatgaggg gcatttaacc ttagcagaca aagcaagaca agaaaacgac gccgttatta 300 tgagtatttt tgtgaatcct gcacaattcg gccctaatga agattttgaa gcatatccgc 360 gcgatattga gcgggatgca gctcttgcag aaaacgccgg agtcgatatt ctttttacgc 420 cagatgetea tgatatgtat eceggtgaaa agaatgteae gatteatgta gaaagaegea 480 cagacgtgtt atgcgggcgc tcaagagaag gacattttga cggggtcgcg atcgtactga 540 cgaagetttt caatetagte aageegaete gtgeetattt eggtttaaaa gatgegeage 600 aggtagctgt tgttgatggg ttaatcagcg acttcttcat ggatattgaa ttggttcctg 660 tegataeggt cagagaggaa gaeggettag ecaaaagete tegeaatgta taettaacag 720 ctgaggaaag aaaagaagcg cctaagctgt atcgggccct tcaaacaagt gcggaacttg 780 tccaagccgg tgaaagagat cctgaagcgg tgataaaagc tgcaaaagat atcattgaaa 840 cgactagcgg aaccatagac tatgtagagc tttattccta tccggaactc gagcctgtga 900 atgaaattgc tggaaagatg attctcgctg ttgcagttgc tttttcaaaa gcgcgtttaa 960 tagataatat cattattgat attcgtagaa aggaggtgaa ttaatatgta tcgtacgatg 1020 atgagcggca aacttcacag ggcaactgtt acggaagcaa acctgaacta tgtgggaagc 1080 attacaattg atgaagatct cattgatgct gtgggaatgc ttcctaatga aaaagtacaa 1140 attgtgaata ataataatgg agcacgtctt gaaacgtata ttattcctgg taaacgggga 1200 ageggegtea tatgettaaa eggtgeagee geaegeettg tgeaggaagg agataaggte 1260 attattattt cctacaaaat gatgtctgat caagaagcgg caagccatga gccgaaagtg 1320 gctgttctga atgatcaaaa caaaattgaa caaatgctgg ggaacgaacc agcccgtaca 1380 attttgtaaa ggatcccccg gggatccctc gaggtcgacc tgcaggggga ccatggtctc 1440 agcgcttgga gccacccgca gttcgaaaaa taataagctt gacctgtgaa gtgaaaaatg 1500 gcgcacattg tgcgacattt tttttgtctg ccgtttaccg ctactgcgtc acggatctcc 1560 acgcgccctg tagcggcgca ttaagcgcgg cgggtgtggt ggttacgcgc agcgtgaccg 1620 cgttcgccgg ctttccccgt caagctctaa atcgggggct ccctttaggg ttccgattta 1740 gtgctttacg gcacctcgac cccaaaaaac ttgattaggg tgatggttca cgtagtgggc 1800 catcgccctg atagacggtt tttcgccctt tgacgttgga gtccacgttc tttaatagtg 1860 gactettgtt ecaaactgga acaacaetca accetatete ggtetattet tttgatttat 1920 aagggatttt gccgatttcg gcctattggt taaaaaatga gctgatttaa caaaaattta 1980 acgcgaattt taacaaaata ttaacgctta caatttcagg tggcactttt cggggaaatg 2040 tgcgcggaac ccctatttgt ttattttct aaatacattc aaatatgtat ccgctcatga 2100 gacaataacc ctgataaatg cttcaataat attgaaaaag gaagagtatg agtattcaac 2160 atttccgtgt cgcccttatt cccttttttg cggcattttg ccttcctgtt tttgctcacc 2220 cagaaacgct ggtgaaagta aaagatgctg aagatcagtt gggtgcacga gtgggttaca 2280 tcgaactgga tctcaacagc ggtaagatcc ttgagagttt tcgccccgaa gaacgttttc 2340 caatgatgag cacttttaaa gttctgctat gtggcgcggt attatcccgt attgacgccg 2400 ggcaagagca acteggtege egeatacaet atteteagaa tgaettggtt gagtaeteae 2460 cagtcacaga aaagcatctt acggatggca tgacagtaag agaattatgc agtgctgcca 2520 taaccatgag tgataacact gcggccaact tacttctgac aacgatcgga ggaccgaagg 2580 agctaaccgc ttttttgcac aacatggggg atcatgtaac tcgccttgat cgttgggaac 2640 cggagctgaa tgaagccata ccaaacgacg agcgtgacac cacgatgcct gtagcaatgg 2700 caacaacgtt gcgcaaacta ttaactggcg aactacttac tctagcttcc cggcaacaat 2760 tgatagactg gatggaggcg gataaagttg caggaccact tctgcgctcg gcccttccgg 2820 ctggctggtt tattgctgat aaatctggag ccggtgagcg tggctctcgc ggtatcattg 2880 cagcactggg gccagatggt aagccctccc gtatcgtagt tatctacacg acggggagtc 2940 aggcaactat ggatgaacga aatagacaga tcgctgagat aggtgcctca ctgattaagc 3000 attggtagga attaatgatg tctcgtttag ataaaagtaa agtgattaac agcgcattag 3060 agctgcttaa tgaggtcgga atcgaaggtt taacaacccg taaactcgcc cagaagctag 3120 gtgtagagca gcctacattg tattggcatg taaaaaataa gcgggctttg ctcgacgcct 3180 tagccattga gatgttagat aggcaccata ctcacttttg ccctttagaa ggggaaagct 3240 ggcaagattt tttacgtaat aacgctaaaa gttttagatg tgctttacta agtcatcgcg 3300 atggagcaaa agtacattta ggtacacggc ctacagaaaa acagtatgaa actctcgaaa 3360 atcaattagc ctttttatgc caacaaggtt tttcactaga gaatgcatta tatgcactca 3420 gcgcagtggg gcattttact ttaggttgcg tattggaaga tcaagagcat caagtcgcta 3480 aagaagaaag ggaaacacct actactgata gtatgccgcc attattacga caagctatcg 3540 aattatttga tcaccaaggt gcagagccag ccttcttatt cggccttgaa ttgatcatat 3600 geggattaga aaaacaactt aaatgtgaaa gtgggtetta aaagcagcat aacettttte 3660 cgtgatggta acttcactag tttaaaagga tctaggtgaa gatccttttt gataatctca 3720 tgaccaaaat cccttaacgt gagttttcgt tccactgagc gtcagacccc gtagaaaaga 3780 tcaaaggatc ttcttgagat ccttttttc tgcgcgtaat ctgctgcttg caaacaaaa 3840 aaccaccgct accagcggtg gtttgtttgc cggatcaaga gctaccaact ctttttccga 3900 aggtaactgg cttcagcaga gcgcagatac caaatactgt ccttctagtg tagccgtagt 3960 taggccacca cttcaagaac tctgtagcac cgcctacata cctcgctctg ctaatcctgt 4020 taccagtggc tgctgccagt ggcgataagt cgtgtcttac cgggttggac tcaagacgat 4080 agttaccgga taaggcgcag cggtcgggct gaacgggggg ttcgtgcaca cagcccagct 4140 tggagcgaac gacctacacc gaactgagat acctacagcg tgagctatga gaaagcgcca 4200 cgcttcccga agggagaaag gcggacaggt atccggtaag cggcagggtc ggaacaggag 4260 agegeacgag ggagetteca gggggaaacg cetggtatet ttatagteet gtegggttte 4320 gccacctctg acttgagcgt cgatttttgt gatgctcgtc aggggggcgg agcctatgga 4380 aaaacgccag caacgcggcc tttttacggt tcctggcctt ttgctggcct tttgctcaca 4440 4450 tgacccgaca

gcggccgcta aaaagagctt gaggatttgc ggagtgaaaa tcagacattg cggaatcagc 60 tagagatgac agaagaggat tacaaggcac tgatcgatat catggatcgg gccagaaaaa 120 tggttgtttc gaaggaagac ggaagaatga aaaaagcggc tcaagaaacg taaagaaacg 180

<210> 81

<211> 10212

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN251 plasmid

cctgaaatga accggcccta tagtaagaat aggccggttg ttttgatttc tatgcagact 240 ctcccggtgt catttcgcga tccatatcag gatgccagat gagcgggtct tcccctttgt 300 cccgcgccat atcatactta acagttttaa agttcatttt gttccaaaat tccgctgatt 360 tcattctcgg atttgtccgg atcggcattt tgaatgattt tgcaaattca accaaggctc 420 tecegtatee cetgttetgg tageeeggaa gaaceteaag etteeacage tecaaataat 480 cctggcggtt gtcaaaatag ggattcgatt tgccgttaac ttgatacaga ctcattcgtg 540 ctacaagttt atcgccaaaa taaatcccgt aaaaaggcga ggtgctgtca ttttcaataa 600 tattatcctg aagttcttca agcattgaaa gctcctgaat gccgtattct ttgaatttct 660 tgaattette cagegtttta tagttgataa geagaegtte taeetttgte aaacaaatet 720 ccccctttgt tgtttctaca tatattgtaa acgctttatt taaaaaatcc aaatatttaa 780 actttaattt taagcacatg ggatctttga gaagtaattt cttcttactt ctgctatgat 840 aatacgtaaa tgcgtcgacc gaaacagcag ttataaggca tgaagctgtc cggtttttgc 900 aaaagtggct gtgactgtaa aaagaaatcg aaaaagaccg ttttgtgtga aaacggtctt 960 tttgtttcct tttaaccaac tgccataact cgaggcctac ctagcttcca agaaagatat 1020 cctaacagca caagagcgga aagatgtttt gttctacatc cagaacaacc tctgctaaaa 1080 ttcctgaaaa attttgcaaa aagttgttga ctttatctac aaggtgtggt ataataatct 1140 taacaacagc aggacgctct agaggaggag acatcatgaa aattggaatt atcggcggag 1200 gctccgttgg tcttttatgc gcctattatt tgtcacttta tcacgacgtg actgttgtga 1260 cgaggcggca agaacaggct gcggccattc agtctgaagg aatccggctt tataaaggcg 1320 gggaggaatt cagggctgat tgcagtgcgg acacgagtat caattcggac tttgacctgc 1380 ttgtcgtgac agtgaagcag catcagcttc aatctgtttt ttcgtcgctt gaacgaatcg 1440 ggaagacgaa tatattattt ttgcaaaacg gcatggggca tatccacgac ctaaaagact 1500 ggcacgttgg ccattccatt tatgttggaa tcgttgagca cggagctgta agaaaatcgg 1560 atacagctgt tgatcataca ggcctaggtg cgataaaatg gagcgcgttc gacgatgctg 1620 aaccagaccg gctgaacatc ttgtttcagc ataaccattc ggattttccg atttattatg 1680 agacggattg gtaccgtctg ctgacgggca agctgattgt aaatgcgtgt attaatcctt 1740 taactgcgtt attgcaagtg aaaaatggag aactgctgac aacgccagct tatctggctt 1800 ttatgaagct ggtatttcag gaggcatgcc gcattttaaa acttgaaaat gaagaaaagg 1860 cttgggagcg ggttcaggcc gtttgtgggc aaacgaaaga gaatcgttca tcaatgctgg 1920 ttgacgtcat tggaggccgg cagacggaag ctgacgccat tatcggatac ttattgaagg 1980 aagcaagtet teaaggtett gatgeegtee acetagagtt tttatatgge agcateaaag 2040 cattggagcg aaatacaaac aaagtctttt gagctttttc ggtaacatgc tatactcatt 2100 teggateact aactatttat tggagaaagg aagttetaga agatgeaget aactgaactt 2160 tccatcaaaa atcagaatgt gtttgtacag cactatatag atggcaaaga agaaatgtct 2220 tetttttttg attacagtat teatcataag gacatgtgge gegaaagaet ggaagaetta 2280 tetteceggt ttttegeaag agaggaattg geggegtaet taacetetta eeataataaa 2340 ttcggttcaa gtgcgatgca gtctgctatt gagaagctga aggacccgtc aagtgccgct 2400 gtagtcggcg gacagcaggc aggactttta acaggaccgc tttacaccat acataaaatc 2460 atttcaatca ttgttttagc aaagcaacaa gaaaaggaac tgcaagtgcc tgtcatacca 2520 atcttctggg tggctggaga agaccacgat ttggatgaga ttaattttgt tcacacatct 2580 gaagagaatg ggcctgtgaa aaaaaagctg cctcagtctt attggaagaa atcatcagca 2640 gcgagtacat cgcttgatca ggaaaagtgt gccgcgtgga tagatgatgt ttttgccgct 2700 tttgaagaaa cagaccatac gaatacactt ctcgacaatg tgaaacgatg tttaagggaa 2760 tetgttacgt ttactgactt etttgaactg etgategegg atttgtteea agaagaggge 2820 ttagttttat taaattctgg ggatcctgtt ttggcggatg agagaagatt ttcagcctga 2880 tacagattaa atcagaacgc agaagcggtc tgataaaaca gaatttgcct ggcggcagta 2940 gegeggtggt eccaectgae eccatgeega acteagaagt gaaaegeegt agegeegatg 3000 gtagtgtggg gtctccccat gcgagagtag ggaactgcca ggcatcaaat aaaacgaaag 3060 geteagtega aagaetggge etttegtttt atetgttgtt tgteggtgaa egeteteetg 3120 agtaggacaa atccgccggg agcggatttg aacgttgcga agcaacggcc cggagggtgg 3180 cgggcaggac gcccgccata aactgccagg catcaaatta agcagaaggc catcctgacg 3240 gatggccttt ttgcgtttct acaaactctt ggtacccaga aaaagcggca aaagcggctg 3300 ttaaaaaagc gaaatcgaag aagctgtctg ccgctaagac ggaatatcaa aagcgttctg 3360 ctgttgtgtc atctttaaaa gtcacagccg atgaatccca gcaagatgtc ctaaaatact 3420 tgaacaccca gaaagataaa ggaaatgcag accaaattca ttcttattat gtggtgaacg 3480 ggattgctgt tcatgcctca aaagaggtta tggaaaaagt ggtgcagttt cccgaagtgg 3540 aaaaggtgct tcctaatgag aaacggcagc tttttaagtc atcctcccca tttaatatga 3600 aaaaagcaca gaaagctatt aaagcaactg acggtgtgga atggaatgta gaccaaatcg 3660 atgccccaaa agcttgggca cttggatatg atggaactgg cacggttgtt gcgtccattg 3720 ataccggggt ggaatggaat catccggcat taaaagagaa atatcgcgga tataatccgg 3780 aaaatcctaa tgagcctgaa aatgaaatga actggtatga tgccgtagca ggcgaggcaa 3840 gecettatga tgatttgget catggaacce acgtgacagg cacgatggtg ggetetgaac 3900 ctgatggaac aaatcaaatc ggtgtagcac ctggcgcaaa atggattgct gttaaagcgt 3960 tctctgaaga tggcggcact gatgctgaca ttttggaagc tggtgaatgg gttttagcac 4020 caaaggacgc ggaaggaaat ccccacccgg aaatggctcc tgatgttgtc aataactcat 4080 ggggaggggg ctctggactt gatgaatggt acagagacat ggtcaatgcc tggcgttcgg 4140 ccgatatttt ccctgagttt tcagcgggga atacggatct ctttattccc ggcgggcctg 4200 gttctatcgc aaatccggca aactatccag aatcgtttgc aactggagcg actgagaatt 4260 ccaattcccc atggagagaa aagaaaatcg ctaatgttga ttactttgaa cttctgcata 4320 ttcttgaatt taaaaaggct gaaagagtaa aagattgtgc tgaaatatta gagtataaac 4380 aaaatcgtga aacaggcgaa agaaagttgt atcgagtgtg gttttgtaaa tccaggcttt 4440 gtccaatgtg caactggagg agagcaatga aacatggcat tcagtcacaa aaggttgttg 4500 ctgaagttat taaacaaaag ccaacagttc gttggttgtt tctcacatta acagttaaaa 4560 atgtttatga tggcgaagaa ttaaataaga gtttgtcaga tatggctcaa ggatttcgcc 4620 gaatgatgca atataaaaaa attaataaaa atcttgttgg ttttatgcgt gcaacggaag 4680 tgacaataaa taataaagat aattettata atcagcacat geatgtattg gtatgtgtgg 4740 aaccaactta ttttaagaat acagaaaact acgtgaatca aaaacaatgg attcaatttt 4800 ggaaaaaggc aatgaaatta gactatgatc caaatgtaaa agttcaaatg attcgaccga 4860 aaaataaata taaatcggat atacaatcgg caattgacga aactgcaaaa tatcctgtaa 4920 aggatacgga ttttatgacc gatgatgaag aaaagaattt gaaacgtttg tctgatttgg 4980 aggaaggttt acaccgtaaa aggttaatct cctatggtgg tttgttaaaa gaaatacata 5040 aaaaattaaa ccttgatgac acagaagaag gcgatttgat tcatacagat gatgacgaaa 5100 aagccgatga agatggattt tctattattg caatgtggaa ttgggaacgg aaaaattatt 5160 ttattaaaga gtagttcaac aaacgggcca tattgttgta taagtgatga aatactgaat 5220 ttaaaactta gtttatatgt ggtaaaatgt tttaatcaag tttaggagga attaattatg 5280 aagtgtaatg aatgtaacag ggttcaatta aaagagggaa gcgtatcatt aaccctataa 5340 actacgtctg ccctcattat tggagggtga aatgtgaata catcctattc acaatcgaat 5400 ttacgacaca accaaatttt aatttggctt tgcattttat ctttttttag cgtattaaat 5460 gaaatggttt tgaacgtctc attacctgat attgcaaatg attttaataa accacctgcg 5520 agtacaaact gggtgaacac agcctttatg ttaacctttt ccattggaac agctgtatat 5580 ggaaagctat ctgatcaatt aggcatcaaa aggttactcc tatttggaat tataataaat 5640 tgtttcgggt cggtaattgg gtttgttggc cattctttct tttccttact tattatggct 5700 cgttttattc aaggggctgg tgcagctgca tttccagcac tcgtaatggt tgtagttgcg 5760 cgctatattc caaaggaaaa taggggtaaa gcatttggtc ttattggatc gatagtagcc 5820 atgggagaag gagtcggtcc agcgattggt ggaatgatag cccattatat tcattggtcc 5880 tatcttctac tcattcctat gataacaatt atcactgttc cgtttcttat gaaattatta 5940 aagaaagaag taaggataaa aggtcatttt gatatcaaag gaattatact aatgtctgta 6000 gtgctgtcat tcctgatatt tgtaaaacat atcaggaaag taacagatcc ttttgttgat 6120 cccggattag ggaaaaatat accttttatg attggagttc tttgtggggg aattatattt 6180 ggaacagtag cagggtttgt ctctatggtt ccttatatga tgaaagatgt tcaccagcta 6240 agtactgccg aaatcggaag tgtaattatt ttccctggaa caatgagtgt cattattttc 6300 ggctacattg gtgggatact tgttgataga agaggtcctt tatacgtgtt aaacatcgga 6360 gttacatttc tttctgttag ctttttaact gcttcctttc ttttagaaac aacatcatgg 6420 ttcatgacaa ttataatcgt atttgtttta ggtgggcttt cgttcaccaa aacagttata 6480 tcaacaattg tttcaagtag cttgaaacag caggaagctg gtgctggaat gagtttgctt 6540 aactttacca gctttttatc agagggaaca ggtattgcaa ttgtaggtgg tttattatcc 6600 ataccettae ttgatcaaag gttgttaeet atggaagttg atcagtcaae ttatetgtat 6660 agtaatttgt tattactttt ttcaggaatc attgtcatta gttggctggt taccttgaat 6720 gtatataaac attctcaaag ggatttctaa atcgttaagg gatcaacttt gggagagagt 6780 tcaaaattga tcctttttt ataacagttc gaagcggccg caattcttga agacgaaagg 6840 geetegtgat aegeetattt ttataggtta atgteatgat aataatggtt tettagaegt 6900 caggtggcac ttttcgggga aatgtgcgcg gaacccctat ttgtttattt ttctaaatac 6960 attcaaatat gtatccgctc atgagacaat aaccctgata aatgcttcaa taatattgaa 7020 aaaggaagag tatgagtatt caacatttcc gtgtcgccct tattcccttt tttgcggcat 7080 tttgccttcc tgtttttgct cacccagaaa cgctggtgaa agtaaaagat gctgaagatc 7140 agttgggtgc acgagtgggt tacatcgaac tggatctcaa cagcggtaag atccttgaga 7200 gttttcgccc cgaagaacgt tttccaatga tgagcacttt taaagttctg ctatgtggcg 7260 cggtattatc ccgtattgac gccgggcaag agcaactcgg tcgccgcata cactattctc 7320 agaatgactt ggttgagtac tcaccagtca cagaaaagca tcttacggat ggcatgacag 7380 taagagaatt atgcagtgct gccataacca tgagtgataa cactgcggcc aacttacttc 7440 tgacaacgat cggaggaccg aaggagctaa ccgctttttt gcacaacatg ggggatcatg 7500 taactcgcct tgatcgttgg gaaccggagc tgaatgaagc cataccaaac gacgagcgtg 7560 acaccacgat gcctgcagca atggcaacaa cgttgcgcaa actattaact ggcgaactac 7620 ttactctagc ttcccggcaa caattaatag actggatgga ggcggataaa gttgcaggac 7680 cacttctgcg ctcggccctt ccggctggct ggtttattgc tgataaatct ggagccggtg 7740 agegtgggte tegeggtate attgcageae tggggeeaga tggtaageee teeegtateg 7800 tagttatcta cacgacgggg agtcaggcaa ctatggatga acgaaataga cagatcgctg 7860 agataggtgc ctcactgatt aagcattggt aactgtcaga ccaagtttac tcatatatac 7920 tttagattga tttaaaactt catttttaat ttaaaaggat ctaggtgaag atcctttttg 7980 ataatctcat gaccaaaatc ccttaacgtg agttttcgtt ccactgagcg tcagaccccg 8040 tagaaaagat caaaggatct tettgagate ettttttet gegegtaate tgetgettge 8100 aaacaaaaaa accaccgcta ccagcggtgg tttgtttgcc ggatcaagag ctaccaactc 8160 tttttccgaa ggtaactggc ttcagcagag cgcagatacc aaatactgtc cttctagtgt 8220 agccgtagtt aggccaccac ttcaagaact ctgtagcacc gcctacatac ctcgctctgc 8280 taatcctgtt accagtggct gctgccagtg gcgataagtc gtgtcttacc gggttggact 8340 caagacgata gttaccggat aaggcgcagc ggtcgggctg aacggggggt tcgtgcacac 8400 ageceagett ggagegaaeg acetaeaeeg aactgagata eetaeagegt gagetatgag 8460 aaagcgccac gcttcccgaa gggagaaagg cggacaggta tccggtaagc ggcagggtcg 8520 gaacaggaga gcgcacgagg gagcttccag ggggaaacgc ctggtatctt tatagtcctg 8580 tegggttteg ceacetetga ettgagegte gatttttgtg atgetegtea ggggggegga 8640 gcctatggaa aaacgccagc aacgcggcct ttttacggtt cctggccttt tgctggcctt 8700 ttgctcacat gttctttcct gcgttatccc ctgattctgt ggataaccgt attaccgcct 8760 ttgagtgagc tgataccgct cgccgcagcc gaacgaccga gcgcagcgag tcagtgagcg 8820 aggaagegga agagegeetg atgeggtatt tteteettae geatetgtge ggtattteae 8880 accgcatatg gtgcactctc agtacaatct gctctgatgc cgcatagtta agccagtata 8940 cactccgcta tcgctacgtg actgggtcat ggctgcgccc cgacacccgc caacacccgc 9000 tgacgcgccc tgacgggctt gtctgctccc ggcatccgct tacagacaag ctgtgaccgt 9060 ctccgggagc tgcatgtgtc agaggttttc accgtcatca ccgaaacgcg cgaggcagct 9120 geggtaaagc teateagegt ggtegtgaag egatteaeag atgtetgeet gtteateege 9180 gtccagctcg ttgagtttct ccagaagcgt taatgtctgg cttctgataa agcgggccat 9240 gttaagggcg gttttttcct gtttggtcac ttgatgcctc cgtgtaaggg ggaatttctg 9300

ttcatggggg taatgatacc gatgaaacga gagaggatgc tcacgatacg ggttactgat 9360 gatgaacatg cccggttact ggaacgttgt gagggtaaac aactggcggt atggatgcgg 9420 cgggaccaga gaaaaatcac tcagggtcaa tgccagcgct tcgttaatac agatgtaggt 9480 gttccacagg gtagccagca gcatcctgcg atgcagatcc ggaacataat ggtgcagggc 9540 gctgacttcc gcgtttccag actttacgaa acacggaaac cgaagaccat tcatgttgtt 9600 getcaggteg cagaegtttt geageageag tegetteaeg ttegetegeg tateggtgat 9660 tcattctgct aaccagtaag gcaaccccgc cagcctagcc gggtcctcaa cgacaggagc 9720 acgatcatgc gcacccgtgg ccaggaccca acgctgcccg agatgcgccg cgtgcggctg 9780 ctggagatgg cggacgcgat ggatatgttc tgccaagggt tggtttgcgc attcacagtt 9840 ctccgcaaga attgattggc tccaattctt ggagtggtga atccgttagc gaggtgccgc 9900 cggcttccat tcaggtcgag gtggcccggc tccatgcacc gcgacgcaac gcggggaggc 9960 agacaaggta tagggeggeg cetacaatee atgecaacee gttecatgtg etegeegagg 10020 cggcataaat cgccgtgacg atcagcggtc cagtgatcga agttaggctg gtaagagccg 10080 cgagcgatcc ttgaagctgt ccctgatggt cgtcatctac ctgcctggac agcatggcct 10140 gcaacgcggg catcccgatg ccgccggaag cgagaagaat cataatgggg aaggccatcc 10200 10212 agcctcgcgt cg

<400> 82
aacaaaattc tccagtcttc acatcggttt gaaaggagga agcggaagga tgaagtaaga 60

gggatttttg actccgaagt aagtcttcaa aaaatcaaat aaggagtgtc aagaatgttt 120

gcaaaacgat tcaaaacctc tttactgccg ttattcgctg gatttttatt gctgtttcat 180

ttggttctgg caggaccggc ggctgcgagt gctgaaacgg cgaacaaatc gaatgagctt 240

acagcaccgt cgatcaaaag cggaaccatt cttcatgcat ggaattggtc gttcaatacg 300

ttaaaacaca atatgaagga tattcatgat gcaggatata cagccattca gacatctccg 360

attaaccaag taaaggaagg gaatcaagga gataaaagca tgtcgaactg gtactggctg 420

tatcagccga catcgtatca aattggcaac cgttacttag gtactgaaca agaatttaaa 480

gaaatgtgtg cagccgctga agaatatggc ataaaggtca ttgttgacgc ggtcatcaat 540

<210> 82

<211> 10426

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN267 plasmid

BGI-141CP - 114 -

cataccacca gtgattatgc cgcgatttcc aatgaggtta agagtattcc aaactggaca 600 catggaaaca cacaaattaa aaactggtct gatcgaaata gtacataatg gatttcctta 660 cgcgaaatac gggcagacat ggcctgcccg gttattatta tttttgacac cagaccaact 720 ggtaatggta gcgaccggcg ctcaggatcg tctcggtacc aagagtttgt agaaacgcaa 780 aaaggccatc cgtcaggatg gccttctgct taatttgatg cctggcagtt tatggcgggc 840 gtcctgcccg ccaccctccg ggccgttgct tcgcaacgtt caaatccgct cccggcggat 900 ttgtcctact caggagagcg ttcaccgaca aacaacagat aaaacgaaag gcccagtctt 960 tegactgage etttegtttt atttgatgee tggeagttee etactetege atggggagae 1020 cccacactac catcggcgct acggcgtttc acttctgagt tcggcatggg gtcaggtggg 1080 accaccgcgc tactgccgcc aggcaaattc tgttttatca gaccgcttct gcgttctgat 1140 ttaatctgta tcaggctgaa aatcttctct catccgccaa aacaggatcc atcacgaagc 1200 gtcgtatcga aaaaattaat tttgcgcaac ggagaccacc gcttccttct tcttgccttg 1260 tttcacaaac ggcatcattt cacgaagett tetteecaet aettegattt gatgttegtt 1320 ctcgcttgca ttgatagcgt tgaaacgagg acggtttact tggttttcga cgatccactc 1380 ttttgcgaat gtaccgtttt ggatatcttt taatacttct ttcatagatt cttttacttt 1440 agcgtccaca acgcgagggc ctgatacgaa atctccccac tgtgctgtgt cagagattga 1500 atatctcatt cctgcaagtc cttcttcgta cataaggtct acgattaatt tcagctcatg 1560 aagacacteg aagtatgeaa gtteaggetg ataacetget teagttaagg ttteaaatee 1620 ggctttgaca agcgcgctta atccgccgca aagaactgct tgctcaccga acaaatctgt 1680 ttctgtttct tctttaaatg tcgtttctaa tacgcccgct cttgcgccgc cgattccttt 1740 agcataagcg agggetttgt etettgette tecagteaca tettgataga ttgcgaacaa 1800 tgcaggtacg ccagctcctt gctcatatgt tcttcttacc aagtgtcccg ggcctttagg 1860 ggccactaag aatacatcta catccgccgg aggaacaatt tgatggaaat gcacgttaaa 1920 tccatgagcg aatactaatg attttcctgc tgtcaattca tctttgattt cagcttcgta 1980 tactttttgc tgctgctcat ccggaagcag aaccatgatg atttcggctt gggctgccgc 2040 ttettttaet gaaaataett tatgteegte ttettggget tgagtgaaag atttteettg 2100 tctaacaccg acgatcacgt ctactccgct ttctttaagg ttcagggcat gtgcgtggcc 2160 ttgcgaaccg tacccgataa ccgctactgt ttttccagcc aatacgttct ctttgatatc 2220 acceptataa tatactttta ccatttcaat ctctcccttg ttatgtttta tacaatagat 2280 attgttttat tggatgacgc cttttgctgg ttcccctcgc aaaagccgtt gtacctgttc 2340 togogattto tttaatgoca taaggtttta ataactcaat aagogottoa attttgttag 2400 attcacctgt cacctgaaca acgatgctgt ctctgctgac atcaacgaca gaggctctaa 2460 acggttctat gattccatta atctctgttc ttgttgaagg tgcggagaca accttgatta 2520 aggccagete cetetggaca ategattgat ttgtgatgte tgtgaettte agcaeateaa 2580 tctgtttgtt gagctgtttc gttaactgtt caacatcatt ttcaccttca acatgaacga 2640 cgaaggtgat tetggaaacg eeggetgttt etgtgtgtee aactgtaatg ettteaatgt 2700 tgtaatgcct ttttgtgaat agaccggtga tccggtttaa caccccggag cggttcacca 2760 cagtcaatgt gataattett tteaaggttt caececeace attteatgea geeettteee 2820 cggagccacc atcgggaata ctttttcttc gctggcaacc cgcacgtcaa tgacaacagg 2880 ttetettgat gttaatgeet etteeagett tteetttget teegettetg atgaaattet 2940 gatgccttta atgccgtatg cttcggacaa tttgacgaag tcaggctgag aagcgaattt 3000 agattetgaa taaegttett eatagaaaat tteetgeeae tgtetgaeea tteegagaea 3060 agcgttattt aaaatcacta ccttgaccgg aagatttaat tcgcgaataa catcgagttc 3120 ttgaagcgtc atttggaatc cgccgtctcc gacaaccgcg acaacagtag catcttttc 3180 ggccagctgt gcgccgatcg ccgccggaag accgaatccc atcgttccaa gtccgcctga 3240 cgtgacccat ttatctgctt tttggaacgg ataaaattgc gctgaccaca tttgatgctg 3300 gcctacatcc gttgcgacaa tggcctctcc ttttgtaaat tgatgaatat attcaatcaa 3360 tttctgaggt ttaaaacctt cttcttcatt atctacatac cagageggat actcttcttt 3420 ccattctgcg agctgttttt tccattcgct tgaatcgctt tgtttgccgt cttgtttgat 3480 cageteetge aggacaattt tgetgtetee gactacagga atetgtgttt teatgatttt 3540 tccgatttca gctggatcaa tatcgatgtg ggctatcttt gcgtttctgg caaagtgttt 3600 caggitteet gigacacggi catcaaaacg ggcgccgata cigattaata gatcacattc 3660 atgaagggcc atattggctg tataagtacc gtgcattccc gccatcccta ggaaaagcgg 3720 atggtcagcc gggaagcctc cgagccccaa aagggtgtgt gccacaggga tttgctgctg 3780 ttcagcataa ttttttaatt cttctgacgc ttttccgtgc agtacgcccg cacccgccag 3840 gatcaccggt tttttcgcac tgctcacggc ttccacaagc ttgcggatct gcaaataatt 3900 cggctctgtt gtcggctggt atcccgggag attcatctca tgatcgtagc tgaattctcc 3960 ttcaattgtt gctacatctt tcggaatatc aatcaataca ggtccgggtc ttccagttgt 4020 tgcaatatgg aacgcttctt taatgatgcg cggcagatct tccggctggc gaacctggta 4080 gctgtgtttt gttactggca tcgtaatccc taaaatgtct gcttcctgaa atgcatcgct 4140 cccgattaca gaggttgcta cctgccctgt aaagacgact aacggcaatg aatcaatcat 4200 ggcatcagca aggcctgtaa caaggtttgt cgctcccggc ctgacgtggc aatgacgaca 4260 ceggtttccg gagaccettg cgtatccetc cgctgcatga attgctcctt gttcgtgacg 4320 ggaaggatat gtaccaacct gaatgtatag cttatcgtaa atcggaagca cagccccgcc 4380 cggataaccg aagatcattt ctactttctc tttttttaat gattcaatca gcattaatcg 4440 teegeteate gtetgtgtae atteggeaga tgetgaatee acetgtaeat tagteeceat 4500 tttatctcct cctctagagc gtcctgctgt tgttaagatt attataccac accttgtaga 4560 taaagtcaac aactttttgc aaaatttttc aggaatttta gcagaggttg ttctggatgt 4620 agaacaaaac atctttccgc tcttgtgctg ttaggatatc tttcttggaa gctaggtagg 4680 cctcgagtta tggcagttgg ttaaaaggaa acaaaaagac cgttttcaca caaaacggtc 4740 tttttcgatt tctttttaca gtcacagcca cttttgcaaa aaccggacag cttcatgcct 4800 tataactgct gtttcggtcg acgaagcggc cgccgtttaa acgaattcct gcagctggcg 4860 aatggcgatt ttcgttcgtg aatacatgtt ataataacta taactaataa cgtaacgtga 4920 ctggcaagag atatttttaa aacaatgaat aggtttacac ttactttagt tttatggaaa 4980 tgaaagatca tatcatatat aatctagaat aaaattaact aaaataatta ttatctagat 5040 aaaaaattta gaagccaatg aaatctataa ataaactaaa ttaagtttat ttaattaaca 5100 actatggata taaaataggt actaatcaaa atagtgagga ggatatattt gaatacatac 5160 gaacaaatta ataaagtgaa aaaaatactt cggaaacatt taaaaaataa ccttattggt 5220 acttacatgt ttggatcagg agttgagagt ggactaaaac caaatagtga tcttgacttt 5280 ttagtcgtcg tatctgaacc attgacagat caaagtaaag aaatacttat acaaaaaatt 5340 agacctattt caaaaaaaat aggagataaa agcaacttac gatatattga attaacaatt 5400 attattcagc aagaaatggt accgtggaat catcctccca aacaagaatt tatttatgga 5460 gaatggttac aagagcttta tgaacaagga tacattcctc agaaggaatt aaattcagat 5520 ttaaccataa tgctttacca agcaaaacga aaaaataaaa gaatatacgg aaattatgac 5580 ttagaggaat tactacctga tattccattt tctgatgtga gaagagccat tatggattcg 5640 tcagaggaat taatagataa ttatcaggat gatgaaacca actctatatt aactttatgc 5700 cgtatgattt taactatgga cacgggtaaa atcataccaa aagatattgc gggaaatgca 5760 gtggctgaat cttctccatt agaacatagg gagagaattt tgttagcagt tcgtagttat 5820 cttggagaga atattgaatg gactaatgaa aatgtaaatt taactataaa ctatttaaat 5880 aacagattaa aaaaattata aaaaaattga aaaaatggtg gaaacacttt tttcaatttt 5940 tttgttttat tatttaatat ttgggaaata ttcattctaa ttggtaatca gattttagaa 6000 aacaataaac ccttgcatag ggggatcgat atccgtttag gctgggcggt gatagcttct 6060 cgttcaggca gtacgcctct tttcttttcc agacctgagg gaggcggaaa tggtgtgagg 6120 ttcccgggga aaagccaaat aggcgatcgc gggagtgctt tatttgaaga tcaggctatc 6180 actgcggtca atagatttca caatgtgatg gctggacagc ctgaggaact ctcgaacccg 6240 aatggaaaca accagatatt tatgaatcag cgcggctcac atggcgttgt gctggcaaat 6300 gcaggttcat cctctgtctc tatcaatacg gcaacaaaat tgcctgatgg caggtatgac 6360 aataaagctg gagcgggttc atttcaagtg aacgatggta aactgacagg cacgatcaat 6420 gccaggtctg tagctgtgct ttatcctgat gatattgcaa aagcgcctca tgttttcctt 6480 gagaattaca aaacaggtgt aacacattct ttcaatgatc aactgacgat taccttgcgt 6540 gcagatgcga atacaacaaa agccgtttat caaatcaata atggaccaga cgacaggcgt 6600 ttaaggatgg agatcaattc acaatcggaa aaggagatcc aatttggcaa aacatacacc 6660 atcatgttaa aaggaacgaa cagtgatggt gtaacgagga ccgagaaata cagttttgtt 6720 aaaagagatc cagcgtcggc caaaaccatc ggctatcaaa atccgaatca ttggagccag 6780 gtaaatgett atatetataa acatgatggg ageegagtaa ttgaattgae eggatettgg 6840 cctggaaaac caatgactaa aaatgcagac ggaatttaca cgctgacgct gcctgcggac 6900 acggatacaa ccaacgcaaa agtgattttt aataatggca gcgcccaagt gcccggtcag 6960 aatcagcctg gctttgatta cgtgctaaat ggtttatata atgactcggg cttaagcggt 7020 tetetteece attgagggea aggetagaeg ggaettaeeg aaagaaacea teaatgatgg 7080 tttctttttt gttcataaat cagacaaaac ttttctcttg caaaagtttg tgaagtgttg 7140 cacaatataa atgtgaaata cttcacaaac aaaaagacat caaagagaaa cataccctgc 7200 aaggatgctg atattgtctg catttgcgcc ggagcaaacc aaaaacctgg tgagacacgc 7260 cttgaattag tagaaaagaa cttgaagatt ttcaaaggca tcgttagtga agtcatggcg 7320 ageggatttg aeggeatttt ettagteggg eggeaceteg etaaeggatt eaceaeteea 7380 agaattggag ccaatcaatt cttgcggaga actgtgaatg cgcaaaccaa cccttggcag 7440 aacatatcca tegegteege cateteeage ageegeacge ggegeatete gggeagegtt 7500 gggtcctggc cacgggtgcg catgatcgtg ctcctgtcgt tgaggacccg gctaggctgg 7560 cggggttgcc ttactggtta gcagaatgaa tcaccgatac gcgagcgaac gtgaagcgac 7620 tgctgctgca aaacgtctgc gacctgagca acaacatgaa tggtcttcgg tttccgtgtt 7680 tegtaaagte tggaaacgeg gaagteageg ceetgeacea ttatgtteeg gatetgeate 7740 gcaggatgct gctggctacc ctgtggaaca cctacatctg tattaacgaa gcgctggcat 7800 tgaccctgag tgatttttct ctggtcccgc cgcatccata ccgccagttg tttaccctca 7860 caacgttcca gtaaccgggc atgttcatca tcagtaaccc gtatcgtgag catcctctct 7920 cgtttcatcg gtatcattac ccccatgaac agaaatcccc cttacacgga ggcatcagtg 7980 accaaacagg aaaaaaccgc ccttaacatg gcccgcttta tcagaagcca gacattaacg 8040 cttctggaga aactcaacga gctggacgcg gatgaacagg cagacatctg tgaatcgctt 8100 cacgaccacg ctgatgagct ttaccgcagc tgcctcgcgc gtttcggtga tgacggtgaa 8160 aacetetgae acatgeaget eeeggagaeg gteacagett gtetgtaage ggatgeeggg 8220 agcagacaag cccgtcaggg cgcgtcagcg ggtgttggcg ggtgtcgggg cgcagccatg 8280 acccagtcac gtagcgatag cggagtgtat actggcttaa ctatgcggca tcagagcaga 8340 ttgtactgag agtgcaccat atgcggtgtg aaataccgca cagatgcgta aggagaaaat 8400 accgcatcag gcgctcttcc gcttcctcgc tcactgactc gctgcgctcg gtcgttcggc 8460 tgcggcgagc ggtatcagct cactcaaagg cggtaatacg gttatccaca gaatcagggg 8520 ataacgcagg aaagaacatg tgagcaaaag gccagcaaaa ggccaggaac cgtaaaaagg 8580 cegegttget ggegttttte cataggetee geeeceetga egageateae aaaaategae 8640 gctcaagtca gaggtggcga aacccgacag gactataaag ataccaggcg tttccccctg 8700 gaageteect egtgegetet eetgtteega eeetgeeget taeeggatae etgteegeet 8760 ttctcccttc gggaagcgtg gcgctttctc atagctcacg ctgtaggtat ctcagttcgg 8820 tgtaggtcgt tcgctccaag ctgggctgtg tgcacgaacc ccccgttcag cccgaccgct 8880 gegeettate eggtaactat egtettgagt eeaaceeggt aagacaegae ttategeeae 8940 tggcagcagc cactggtaac aggattagca gagcgaggta tgtaggcggt gctacagagt 9000 tettgaagtg gtggeetaac taeggetaca etagaaggae agtatttggt atetgegete 9060 tgctgaagcc agttaccttc ggaaaaagag ttggtagctc ttgatccggc aaacaaacca 9120 ccgctggtag cggtggtttt tttgtttgca agcagcagat tacgcgcaga aaaaaaggat 9180 ctcaagaaga teetttgate ttttetaegg ggtetgaege teagtggaae gaaaaeteae 9240 gttaagggat tttggtcatg agattatcaa aaaggatctt cacctagatc cttttaaatt 9300 aaaaatgaag ttttaaatca atctaaagta tatatgagta aacttggtct gacagttacc 9360 aatgcttaat cagtgaggca cctatctcag cgatctgtct atttcgttca tccatagttg 9420 cctgactccc cgtcgtgtag ataactacga tacgggaggg cttaccatct ggccccagtg 9480 ctgcaatgat accgcgagac ccacgctcac cggctccaga tttatcagca ataaaccagc 9540 cageeggaag ggeegagege agaagtggte etgeaacttt ateegeetee ateeagteta 9600 ttaattgttg ccgggaagct agagtaagta gttcgccagt taatagtttg cgcaacgttg 9660 ttgccattgc tgcaggcatc gtggtgtcac gctcgtcgtt tggtatggct tcattcagct 9720 ccggttcccaacgatcaaggcgagttacatgatccccatgttgtcaaaaaagcggtta9780gctccttcggtcctccgatcgttgtcagaagtaagttggccgcagtgttatcactcatgg9840ttatggcagcactgcataattctcttactgtcatgccatccgtaagatgcttttctgtga9900ctggtgagtactcaaccaagtcattctgagaatagtgtatgcggcgaccgagttgctctt9960gcccggcgtcaacacgggataataccgcgccacatagcagaactttaaaagtgctcatca10020ttggaaaacgttcttcggggcgaaaactctcaaggatctaccgctgttgagatccagtt10140ctgggtgagcaaaaacaggaaggcaaaatgccgcaaaaaagggaataagggcgacacgga10200aatgttgaatactcatactcttccttttcaatattattgaagcatttatcagggttatt10260gtctcatgagcggaaaagtgccacctgacgtctaagaaaataaacaaataggggttccgc10320gcacatttccccgaaaagtgccacctgacgtctaagaaaccattattatcatgacattaa10380cctataaaaataggcgtatcacgaggccctttcgtcttcaagaatt10426

<400> 83
ccaatacgca aaccgcctct ccccgcgcgt tggccgattc attaatgcag cccggatccg 60
aaagaagagg atatcccggt tttacagaag gcattggatg atccaaaggt gtccatcaga 120
agacaggctg ttgtgtactt aggaatgatt gaaacacctg atgttcttcc tctattgtat 180
aaagcacttg aggacaaagc tgtatcagtc agaagaacgg ccggagactg cctgtctgat 240
atcggcgatc ctcaagccat tcctgctatg atcaagtcat taagcgactc cagcaagctt 300
gttcgctggc gtgccgccat gttcctgtac gaagtcggcg atgaaagtgc aattgaagct 360
ttgcgcgctg ccgaagatga ccccgaattt gaggtcagcc ttcaagtcaa aatggcgctt 420
gaacgtattg agcatggaga agaagcaaaa ggttctgttt ggaaacaaat gacggaaagc 480
aggacaaaaag gcgaataaag ataaaaaagg tgcagatcat gcacctttt tatgtgaatt 540
ggtcgaccga aacagcagtt ataaggcatg aagctgtccg gtttttgcaa aagtggctg 600
gactgtaaaa agaaatcgaa aaagaccgtt ttggtgaaa acggtctttt tgtttccttt 660
taaccaactg ccataactcg aggcctacct agctccaag aaagatatcc taacagcaca 720
agagcggaaa gatgttttgt tctacatcca gaacaacctc tgctaaaat cctgaaaaat 780

<210> 83

<211> 4191

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN263 plasmid

tttgcaaaaa gttgttgact ttatctacaa ggtgtggtat aataatctta acaacagcag 840 gacgctctag aggaggagac accatggcag aattacgcag taatatgatc acacaaggaa 900 tegatagage teegeacege agtttgette gtgcageagg ggtaaaagaa gaggattteg 960 gcaagccgtt tattgcggtg tgtaattcat acattgatat cgttcccggt catgttcact 1020 tgcaggagtt tgggaaaatc gtaaaagaag caatcagaga agcagggggc gttccgtttg 1080 aatttaatac cattggggta gatgatggca tcgcaatggg gcatatcggt atgagatatt 1140 cgctgccaag ccgtgaaatt atcgcagact ctgtggaaac ggttgtatcc gcacactggt 1200 ttgacggaat ggtctgtatt ccgaactgcg acaaaatcac accgggaatg cttatggcgg 1260 caatgcgcat caacattccg acgatttttg tcagcggcgg accgatggcg gcaggaagaa 1320 caagttacgg gcgaaaaatc tccctttcct cagtattcga aggggtaggc gcctaccaag 1380 cagggaaaat caacgaaaac gagcttcaag aactagagca gttcggatgc ccaacgtgcg 1440 ggtcttgctc aggcatgttt acggcgaact caatgaactg tctgtcagaa gcacttggtc 1500 ttgctttgcc gggtaatgga accattctgg caacatctcc ggaacgcaaa gagtttgtga 1560 gaaaatcggc tgcgcaatta atggaaacga ttcgcaaaga tatcaaaccg cgtgatattg 1620 ttacagtaaa agcgattgat aacgcgtttg cactcgatat ggcgctcgga ggttctacaa 1680 ataccettct tcataccett geeettgeaa acgaageegg egttgaatae tetttagaae 1740 gcattaacga agtcgctgag cgcgtgccgc acttggctaa gctggcgcct gcatcggatg 1800 tgtttattga agatetteae gaagegggeg gegttteage ggetetgaat gagetttega 1860 agaaagaagg agcgcttcat ttagatgcgc tgactgttac aggaaaaact cttggagaaa 1920 ccattgccgg acatgaagta aaggattatg acgtcattca cccgctggat caaccattca 1980 ctgaaaaggg aggccttgct gttttattcg gtaatctagc tccggacggc gctatcatta 2040 aaacaggcgg cgtacagaat gggattacaa gacacgaagg gccggctgtc gtattcgatt 2100 ctcaggacga ggcgcttgac ggcattatca accgaaaagt aaaagaaggc gacgttgtca 2160 tcatcagata cgaagggcca aaaggcggac ctggcatgcc ggaaatgctg gcgccaacat 2220 cccaaatcgt tggaatggga ctcgggccaa aagtggcatt gattacggac ggacgttttt 2280 ccggagcctc ccgtggcctc tcaatcggcc acgtatcacc tgaggccgct gagggcgggc 2340 cgcttgcctt tgttgaaaac ggagaccata ttatcgttga tattgaaaaa cgcatcttgg 2400 atgtacaagt gccagaagaa gagtgggaaa aacgaaaagc gaactggaaa ggttttgaac 2460 cgaaagtgaa aaccggctac ctggcacgtt attctaaact tgtgacaagt gccaacaccg 2520 geggtattat gaaaatetag acceetggeg taatagegaa gaggeeegea eegategeee 2580 in .j

 BGI-141CP - 121 -

ttcccaacag ttgcgcagcc tgaatggcga atgagcttgc gccgtcccgt caagtcagcg 2640 taatgctctg ccagtgttac aaccaattaa ccaattctga ttagaaaaac tcatcgagca 2700 tcaaatgaaa ctgcaattta ttcatatcag gattatcaat accatatttt tgaaaaagcc 2760 gtttctgtaa tgaaggagaa aactcaccga ggcagttcca taggatggca agatcctggt 2820 atcggtctgc gattccgact cgtccaacat caatacaacc tattaatttc ccctcgtcaa 2880 aaataaggtt atcaagtgag aaatcaccat gagtgacgac tgaatccggt gagaatggca 2940 aaaggttatg catttctttc cagacttgtt caacaggcca gccattacgc tcgtcatcaa 3000 aatcactcgc atcaaccaaa ccgttattca ttcgtgattg cgcctgagcg agacgaaata 3060 cgcgatcgct gttaaaagga caattacaaa caggaatcga atgcaaccgg cgcaggaaca 3120 ctgccagcgc atcaacaata ttttcacctg aatcaggata ttcttctaat acctggaatg 3180 ctgttttccc agggatcgca gtggtgagta accatgcatc atcaggagta cggataaaat 3240 gettgatggt eggaagagge ataaatteeg teageeagtt tagtetgace ateteatetg 3300 taacatcatt ggcaacgcta cctttgccat gtttcagaaa caactctggc gcatcgggct 3360 teccatacaa teaatagatt gtegeaeetg attgeeegae attategega geeeatttat 3420 acccatataa atcagcatcc atgttggaat ttaatcgcgg cctcgacgag caagacgttt 3480 cccgttgaat atggctcata acaccccttg tattactgtt tatgtaagca gacagtttta 3540 ttgttcatga tgatatattt ttatcttgtg caatgtaaca tcagagattt tgagacactc 3600 gacaagatga tettettgag ategttttgg tetgegegta atetettget etgaaaaega 3660 aaaaaccgcc ttgcagggcg gtttttcgaa ggttctctga gctaccaact ctttgaaccg 3720 aggtaactgg cttggaggag cgcagtcacc aaaacttgtc ctttcagttt agccttaacc 3780 ggcgcatgac ttcaagacta actcctctaa atcaattacc agtggctgct gccagtggtg 3840 cttttgcatg tctttccggg ttggactcaa gacgatagtt accggataag gcgcagcggt 3900 cggactgaac ggggggttcg tgcatacagt ccagcttgga gcgaactgcc tacccggaac 3960 tgagtgtcag gcgtggaatg agacaaacgc ggccataaca gcggaatgac accggtaaac 4020 cgaaaggcag gaacaggaga gcgcacgagg gagccgccag gggaaacgcc tggtatcttt 4080 atagteetgt egggtttege eaceactgat ttgagegtea gatttegtga tgettgteag 4140 gggggcggag cctatggaaa aacggctttg ccgcggccct ctcacttccc t 4191

<210> 84

<211> 702

<212> DNA

<213> Bacillus subtilis

<221> CDS <222> (1)..(699)

<400> 84 ttg tta ctg gtt atc gat gtg ggg aac acc aat act gta ctt ggt gta Met Leu Leu Val Ile Asp Val Gly Asn Thr Asn Thr Val Leu Gly Val tat cat gat gga aaa tta gaa tat cac tgg cgt ata gaa aca agc agg 96 Tyr His Asp Gly Lys Leu Glu Tyr His Trp Arg Ile Glu Thr Ser Arg 2.0 cat aaa aca gaa gat gag ttt ggg atg att ttg cgc tcc tta ttt gat 144 His Lys Thr Glu Asp Glu Phe Gly Met Ile Leu Arg Ser Leu Phe Asp cac tcc ggg ctt atg ttt gaa cag ata gat ggc att att att tcg tca 192 His Ser Gly Leu Met Phe Glu Gln Ile Asp Gly Ile Ile Ile Ser Ser 50 gta gtg ccg cca atc atg ttt gcg tta gaa aga atg tgc aca aaa tac Val Val Pro Pro Ile Met Phe Ala Leu Glu Arg Met Cys Thr Lys Tyr 70 ttt cat atc gag cct caa att gtt ggt cca ggt atg aaa acc ggt tta Phe His Ile Glu Pro Gln Ile Val Gly Pro Gly Met Lys Thr Gly Leu 85 aat ata aaa tat gac aat ccg aaa gaa gta ggg gca gac aga atc gta Asn Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile Val 100 aat gct gtc gct gcg ata cac ttg tac ggc aat cca tta att gtt gtc Asn Ala Val Ala Ala Ile His Leu Tyr Gly Asn Pro Leu Ile Val Val 115 gat ttc gga acc gcc aca acg tac tgc tat att gat gaa aac aaa caa Asp Phe Gly Thr Ala Thr Thr Tyr Cys Tyr Ile Asp Glu Asn Lys Gln 130 tac atg ggc ggg gcg att gcc cct ggg att aca att tcg aca gag gcg Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala 155 ctt tac tcg cgt gca gca aag ctt cct cgt atc gaa atc acc cgg ccc Leu Tyr Ser Arg Ala Ala Lys Leu Pro Arg Ile Glu Ile Thr Arg Pro gac aat att atc gga aaa aac act gtt agc gcg atg caa tct gga att Asp Asn Ile Ile Gly Lys Asn Thr Val Ser Ala Met Gln Ser Gly Ile 180 tta ttt ggc tat gtc ggc caa gtg gaa gga atc gtt aag cga atg aaa Leu Phe Gly Tyr Val Gly Gln Val Glu Gly Ile Val Lys Arg Met Lys 200 tgg cag gca aaa cag gac cca agg tca ttg cga cag gag gcc tgg cgc Trp Gln Ala Lys Gln Asp Pro Arg Ser Leu Arg Gln Glu Ala Trp Arg 215

cgc tca ttg cga acg aat cag att gta tag

702

Arg Ser Leu Arg Thr Asn Gln Ile Val 225 230

<210> 85

<211> 233

<212> PRT

<213> Bacillus subtilis

<400> 85

Met Leu Leu Val Ile Asp Val Gly Asn Thr Asn Thr Val Leu Gly Val 1 5 10 15

Tyr His Asp Gly Lys Leu Glu Tyr His Trp Arg Ile Glu Thr Ser Arg 20 25 30

His Lys Thr Glu Asp Glu Phe Gly Met Ile Leu Arg Ser Leu Phe Asp 35 40 45

His Ser Gly Leu Met Phe Glu Gln Ile Asp Gly Ile Ile Ile Ser Ser 50 60

Val Val Pro Pro Ile Met Phe Ala Leu Glu Arg Met Cys Thr Lys Tyr 65 70 75 80

Phe His Ile Glu Pro Gln Ile Val Gly Pro Gly Met Lys Thr Gly Leu 85 90 95

Asn Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile Val

Asn Ala Val Ala Ala Ile His Leu Tyr Gly Asn Pro Leu Ile Val Val 115 120 125

Asp Phe Gly Thr Ala Thr Thr Tyr Cys Tyr Ile Asp Glu Asn Lys Gln 130 135 140

Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala 145 150 155 160

Leu Tyr Ser Arg Ala Ala Lys Leu Pro Arg Ile Glu Ile Thr Arg Pro 165 170 175

Asp Asn Ile Ile Gly Lys Asn Thr Val Ser Ala Met Gln Ser Gly Ile 180 185 190

Leu Phe Gly Tyr Val Gly Gln Val Glu Gly Ile Val Lys Arg Met Lys 195 200 205

Trp Gln Ala Lys Gln Asp Pro Arg Ser Leu Arg Gln Glu Ala Trp Arg 210 215 220

Arg Ser Leu Arg Thr Asn Gln Ile Val 225 230

<210> 86

<211> 1623

<212> DNA

<213> Bacillus subtilis

<220>

<221> CDS <222> (1)..(1620)

<400> 86 atg tat ttg gca ttc cag gtg caa aaa ttg atg cgg tat ttg acg ctt Met Tyr Leu Ala Phe Gln Val Gln Lys Leu Met Arg Tyr Leu Thr Leu tac aag ata aag gac ctg aaa tta tcg ttg ccc ggc acg aac aaa acg Tyr Lys Ile Lys Asp Leu Lys Leu Ser Leu Pro Gly Thr Asn Lys Thr 20 cag caa ttc atg gcc caa gca gtc ggc cgt tta act gga aaa ccg gga 144 Gln Gln Phe Met Ala Gln Ala Val Gly Arg Leu Thr Gly Lys Pro Gly 35 gtc gtg tta gtc aca tca gga ccg ggt gcc tct aac ttg gca aca ggc 192 Val Val Leu Val Thr Ser Gly Pro Gly Ala Ser Asn Leu Ala Thr Gly 55 50 ctg ctg aca gcg aac act gaa gga gac cct gtc gtt gcg ctt gct gga 240 Leu Leu Thr Ala Asn Thr Glu Gly Asp Pro Val Val Ala Leu Ala Gly 70 aac gtg atc cgt gca tat cgt tta aaa cgg aca cat caa tct ttg gat 288 Asn Val Ile Arg Ala Tyr Arg Leu Lys Arg Thr His Gln Ser Leu Asp aat gcg gcg cta ttc cag ccg att aca aaa tac agt gta gaa gtt caa Asn Ala Ala Leu Phe Gln Pro Ile Thr Lys Tyr Ser Val Glu Val Gln 100 105 gat gta aaa aat ata ccg gaa gct gtt aca aat gca ttt agg ata gcg Asp Val Lys Asn Ile Pro Glu Ala Val Thr Asn Ala Phe Arg Ile Ala 120 115 tca gca ggg cag gct ggg gcc gct ttt gtg agc ttt ccg caa gat gtt Ser Ala Gly Gln Ala Gly Ala Ala Phe Val Ser Phe Pro Gln Asp Val 130 gtg aat gaa gtc aca aat acg aaa aac gtg cgt gct gtt gca gcg cca Val Asn Glu Val Thr Asn Thr Lys Asn Val Arg Ala Val Ala Ala Pro aaa ctc ggt cct gca gca gat gat gca atc agt gcg gcc ata gca aaa Lys Leu Gly Pro Ala Ala Asp Asp Ala Ile Ser Ala Ala Ile Ala Lys 165 atc caa aca gca aaa ctt cct gtc gtt ttg gtc ggc atg aaa ggc gga Ile Gln Thr Ala Lys Leu Pro Val Val Leu Val Gly Met Lys Gly Gly 180 aga ccg gaa gca att aaa gcg gtt cgc aag ctt ttg aaa aag gtt cag 624 Arg Pro Glu Ala Ile Lys Ala Val Arg Lys Leu Leu Lys Lys Val Gln 200 ctt cca ttt gtt gaa aca tat caa gct gcc ggt acc ctt tct aga gat Leu Pro Phe Val Glu Thr Tyr Gln Ala Ala Gly Thr Leu Ser Arg Asp 215 tta gag gat caa tat ttt ggc cgt atc ggt ttg ttc cgc aac cag cct 720

Leu (225	Glu	Asp	Gln		Phe 230	Gly	Arg	Ile	Gly	Leu 235	Phe	Arg	Asn	Gln	Pro 240	
ggc (gat Asp	tta Leu	ctg Leu	cta Leu 245	gag Glu	cag Gln	gca Ala	Asp	gtt Val 250	gtt Val	ctg Leu	acg Thr	atc Ile	ggc Gly 255	tat Tyr	768
gac Asp	ccg Pro	att Ile	gaa Glu 260	tat Tyr	gat Asp	ccg Pro	aaa Lys	ttc Phe 265	tgg Trp	aat Asn	atc Ile	aat Asn	gga Gly 270	gac Asp	cgg Arg	816
aca Thr	att Ile	atc Ile 275	cat His	tta Leu	gac Asp	gag Glu	att Ile 280	atc Ile	gct Ala	gac Asp	att Ile	gat Asp 285	cat His	gct Ala	tac Tyr	864
cag Gln	cct Pro 290	gat Asp	ctt Leu	gaa Glu	ttg Leu	atc Ile 295	ggt Gly	gac Asp	att Ile	ccg Pro	tcc Ser 300	acg Thr	atc Ile	aat Asn	cat His	912
atc Ile 305	gaa Glu	cac His	gat Asp	gct Ala	gtg Val 310	aaa Lys	gtg Val	gaa Glu	ttt Phe	gca Ala 315	gag Glu	cgt Arg	gag Glu	cag Gln	aaa Lys 320	960
atc Ile	ctt Leu	tct Ser	gat Asp	tta Leu 325	aaa Lys	caa Gln	tat Tyr	atg Met	cat His 330	gaa Glu	ggt Gly	gag Glu	cag Gln	gtg Val 335	cct Pro	1008
gca Ala	gat Asp	tgg Trp	aaa Lys 340	tca Ser	gac Asp	aga Arg	gcg Ala	cac His 345	cct Pro	ctt Leu	gaa Glu	atc Ile	gtt Val 350	aaa Lys	gag Glu	1056
ttg Leu	cgt Arg	aat Asn 355	gca Ala	gtc Val	gat Asp	gat Asp	cat His 360	gtt Val	aca Thr	gta Val	act Thr	tgc Cys 365	gat Asp	atc Ile	ggt Gly	1104
tcg Ser	cac His 370	tcc Ser	att Ile	tgg Trp	atg Met	tca Ser 375	Arg	tat Tyr	ttc Phe	cgc Arg	ago Ser 380	tac Tyr	gag Glu	ccg Pro	tta Leu	1152
aca Thr 385	tta Leu	atg Met	ato	agt Ser	aac Asn 390	ggt Gly	atg Met	caa Gln	aca Thr	ctc Leu 395	. Gly	gtt Val	gcg Ala	ctt Leu	cct Pro 400	1200
tgg Trp	gca Ala	ato Ile	ggc Gly	gct Ala 405	Ser	ttg Leu	gtg Val	aaa Lys	ccg Pro 410	Gly	gaa Glu	a aaa 1 Lys	gtg Val	gtt Val 415	tct Ser	1248
gtc Val	tct Ser	ggt Gly	gac Asp 420	Gly	ggt Gly	ttc Phe	: tta : Leu	tto Phe 425	e Ser	gca Ala	atg Met	g gaa : Glu	tta Leu 430	. GIU	g aca 1 Thr	1296
gca Ala	gtt Val	cga Arg 435	, Lei	aaa Lys	gca Ala	cca Pro	a att 11e 440	val	cac His	att Ile	gta Val	a tgg l Trp 445) Asr	gac As <u>r</u>	e agc 9 Ser	1344
aca Thr	tat Tyr 450	Asp	ato Met	g gtg : Val	g cat His	tto Phe 455	e Glr	g caa n Glr	ı ttç ı Lei	g aaa 1 Lys	a aaa 5 Ly: 46	з Туг	aac Asr	: cgt n Arg	aca g Thr	1392
tct Ser	gcg Ala	g gto a Val	c gat L Asp	t tto p Phe	c gga e Gly	a aat 7 Asi	t ato	gat Asp	ato	c gtg e Val	g aa L Ly	a tat s Ty:	gcg Ala	g gaa a Glu	a agc ı Ser	1440

470 465 475 480 tte qqa qea act qeg ttg ege gta gaa tea eea gae eag etg gea gat 1488 Phe Gly Ala Thr Ala Leu Arg Val Glu Ser Pro Asp Gln Leu Ala Asp 485 490 gtt ctg cgt caa ggc atg aac gct gaa ggt cct gtc atc atc gat gtc 1536 Val Leu Arg Gln Gly Met Asn Ala Glu Gly Pro Val Ile Ile Asp Val 505 ccg gtt gac tac agt gat aac att aat tta gca agt gac aag ctt ccg 1584 Pro Val Asp Tyr Ser Asp Asn Ile Asn Leu Ala Ser Asp Lys Leu Pro 515 aaa gaa ttc ggg gaa ctc atg aaa acg aaa gct ctc tag 1623 Lys Glu Phe Gly Glu Leu Met Lys Thr Lys Ala Leu <210> 87 <211> 540 <212> PRT <213> Bacillus subtilis <400> 87 Met Tyr Leu Ala Phe Gln Val Gln Lys Leu Met Arg Tyr Leu Thr Leu Tyr Lys Ile Lys Asp Leu Lys Leu Ser Leu Pro Gly Thr Asn Lys Thr Gln Gln Phe Met Ala Gln Ala Val Gly Arg Leu Thr Gly Lys Pro Gly 45 Val Val Leu Val Thr Ser Gly Pro Gly Ala Ser Asn Leu Ala Thr Gly Leu Leu Thr Ala Asn Thr Glu Gly Asp Pro Val Val Ala Leu Ala Gly Asn Val Ile Arg Ala Tyr Arg Leu Lys Arg Thr His Gln Ser Leu Asp Asn Ala Ala Leu Phe Gln Pro Ile Thr Lys Tyr Ser Val Glu Val Gln 105 110 Asp Val Lys Asn Ile Pro Glu Ala Val Thr Asn Ala Phe Arg Ile Ala 120 Ser Ala Gly Gln Ala Gly Ala Ala Phe Val Ser Phe Pro Gln Asp Val 135 Val Asn Glu Val Thr Asn Thr Lys Asn Val Arg Ala Val Ala Ala Pro Lys Leu Gly Pro Ala Ala Asp Asp Ala Ile Ser Ala Ala Ile Ala Lys 165 170 Ile Gln Thr Ala Lys Leu Pro Val Val Leu Val Gly Met Lys Gly Gly

185

180

- Arg Pro Glu Ala Ile Lys Ala Val Arg Lys Leu Leu Lys Lys Val Gln 195 200 205
- Leu Pro Phe Val Glu Thr Tyr Gln Ala Ala Gly Thr Leu Ser Arg Asp 210 215 220
- Leu Glu Asp Gln Tyr Phe Gly Arg Ile Gly Leu Phe Arg Asn Gln Pro 225 230 235
- Gly Asp Leu Leu Glu Gln Ala Asp Val Val Leu Thr Ile Gly Tyr 245 250 255
- Asp Pro Ile Glu Tyr Asp Pro Lys Phe Trp Asn Ile Asn Gly Asp Arg 260 265 270
- Thr Ile Ile His Leu Asp Glu Ile Ile Ala Asp Ile Asp His Ala Tyr 275 280 285
- Gln Pro Asp Leu Glu Leu Ile Gly Asp Ile Pro Ser Thr Ile Asn His 290 295 300
- Ile Glu His Asp Ala Val Lys Val Glu Phe Ala Glu Arg Glu Gln Lys 305 310 315 320
- Ile Leu Ser Asp Leu Lys Gln Tyr Met His Glu Gly Glu Gln Val Pro 325 330 335
- Ala Asp Trp Lys Ser Asp Arg Ala His Pro Leu Glu Ile Val Lys Glu 340 345 350
- Leu Arg Asn Ala Val Asp Asp His Val Thr Val Thr Cys Asp Ile Gly 355 360 365
- Ser His Ser Ile Trp Met Ser Arg Tyr Phe Arg Ser Tyr Glu Pro Leu 370 375 380
- Thr Leu Met Ile Ser Asn Gly Met Gln Thr Leu Gly Val Ala Leu Pro 385 390 395 400
- Trp Ala Ile Gly Ala Ser Leu Val Lys Pro Gly Glu Lys Val Val Ser 405 410 415
- Val Ser Gly Asp Gly Gly Phe Leu Phe Ser Ala Met Glu Leu Glu Thr 420 425 430
- Ala Val Arg Leu Lys Ala Pro Ile Val His Ile Val Trp Asn Asp Ser 435 440 445
- Thr Tyr Asp Met Val His Phe Gln Gln Leu Lys Lys Tyr Asn Arg Thr 450 455 460
- Ser Ala Val Asp Phe Gly Asn Ile Asp Ile Val Lys Tyr Ala Glu Ser 465 470 475 480
- Phe Gly Ala Thr Ala Leu Arg Val Glu Ser Pro Asp Gln Leu Ala Asp 485 490 495
- Val Leu Arg Gln Gly Met Asn Ala Glu Gly Pro Val Ile Ile Asp Val 500 505 510
- Pro Val Asp Tyr Ser Asp Asn Ile Asn Leu Ala Ser Asp Lys Leu Pro

525 520 515 Lys Glu Phe Gly Glu Leu Met Lys Thr Lys Ala Leu 535 <210> 88 <211> 23 <212> DNA <213> Artificial Sequence <220> <223> Description of Artificial Sequence: ribosome binding site <220> <223> All occurrences of n indicate any nucleotide <400> 88 23 agaaaggagg tgannnnnn atg <210> 89 <211> 7 <212> PRT <213> Artificial Sequence <220> <223> Description of Artificial Sequence: PanC C terminus <400> 89 Ile Arg Glu Met Glu Arg Ile <210> 90 <211> 5 <212> PRT <213> Artificial Sequence <220> <223> Description of Artificial Sequence: PanC C terminus <400> 90 Ile Arg Glu Arg Arg <210> 91 <211> 7 <212> PRT <213> Artificial Sequence <223> Description of Artificial Sequence: PanC C terminus <400> 91

Ile Arg Arg Lys Glu Val Asn

1

<210> 92

<211> 6688

<212> DNA

<213> Artificial Sequence

5

<220>

<223> Description of Artificial Sequence: Recombinant pAN336 plasmid

<400> 92 tgcgccgcta cagggcgcgt ccattcgcca ttcaggctgc gcaactgttg ggaagggcga 60 teggtgeggg cetetteget attacgeeag etggegaaag ggggatgtge tgeaaggega 120 ttaagttggg taacgccagg gttttcccag tcacgacgtt gtaaaacgac ggccagtgaa 180 ttgtaatacg actcactata gggcgaattg ggcccgacgt cgcatgcacc aggcttctca 240 ggcgctgact tagaaaacct cttgaatgaa gctgcgcttg tagcggctcg tcaaaacaag 300 aaaaaaatcg atgcgcgtga tattgacgaa gcgacggacc gtgtaattgc cggacccgct 360 aagaagagcc gcgttatctc caagaaagaa cgcaatatcg tggcttatca cgaaggcgga 420 cacaccgtta tcggtctcgt tttagatgag gcagatatgg ttcataaagt aacgattgtt 480 cctcggggcc aggctggcgg ttatgctgtt atgctgccaa gagaagaccg ttatttccaa 540 acaaagccgg agctgcttga taaaattgtc ggcctcttgg gcggacgtgt tgctgaagag 600 attatetteg gtgaagteag cacaggggeg cacaatgaet tecagegtge gaegaatatt 660 gcaagacgaa tggttacaga attcggtatg tcagaaaaac tgggaccgtt gcaatttgga 720 cagtctcagg gcggtcaggt attcttaggc cgtgatttca acaacgaaca gaactacagt 780 gatcaaatcg cttacgaaat tgatcaggaa attcagcgca tcatcaaaga atgttatgag 840 cgtgcgaaac aaatcctgac tgaaaatcgt gacaagcttg aattgattgc ccaaacgctt 900 ctgaaagttg aaacgcttga cgctgaacaa atcaaacacc ttatcgatca tggaacatta 960 cctgagcgta atttctcaga tgatgaaaag aacgatgatg tgaaagtaaa cattctgaca 1020 aaaacagaag aaaagaaaga cgatacgaaa gagtaattcg ctttctttct aaaaaaactg 1080 ccggctgacg ctggcagttt ttttatgtaa atgattggct cagctgcggc ttttacaatc 1140 atccaattct ggtatcgatt tgtttacaaa tgagccgctg atcgtgtatg gtattgtaga 1200 atgtttgtaa aaagtaaagt agagaaacta ttcaaaagtg gtgatagagg ttgttactgg 1260 ttatcgatgt ggggaacacc ctgcagctcg agtgaaatac cgcacagatg cgtaaggaga 1320 aaataccgca tcaggcgata aacccagcga accatttgag gtgataggta agattatacc 1380 gaggtatgaa aacgagaatt ggacctttac agaattactc tatgaagcgc catatttaaa 1440 aagctaccaa gacgaagagg atgaagagga tgaggaggca gattgccttg aatatattga 1500

caatactgat aagataatat atcttttata tagaagatat cgccgtatgt aaggatttca 1560 gggggcaagg cataggcagc gcgcttatca atatatctat agaatgggca aaqcataaaa 1620 acttqcatqq actaatqctt qaaacccaqq acaataacct tataqcttqt aaattctatc 1680 ataattgtgg tttcaaaatc ggctccgtcg atactatgtt atacgccaac tttcaaaaca 1740 actttqaaaa agctgttttc tqqtatttaa qqttttagaa tqcaaqqaac aqtqaattqq 1800 agttcgtctt gttataatta gcttcttggg gtatctttaa atactgtaga aaagaggaag 1860 gaaataataa atggctaaaa tgagaatatc accggaattg aaaaaactga tcgaaaaata 1920 ccgctgcgta aaagatacgg aaggaatgtc tcctgctaag gtatataagc tggtgggaga 1980 aaatgaaaac ctatatttaa aaatgacgga cagccggtat aaagggacca cctatgatgt 2040 ggaacgggaa aaggacatga tgctatggct ggaaggaaag ctgcctgttc caaaggtcct 2100 gcactttgaa cggcatgatg gctggagcaa tctgctcatg agtgaggccg atggcgtcct 2160 ttgctcggaa gagtatgaag atgaacaaag ccctgaaaag attatcgagc tgtatgcgga 2220 gtgcatcagg ctctttcact ccatcgacat atcggattgt ccctatacga atagcttaga 2280 cagccgctta gccgaattgg attacttact gaataacgat ctggccgatg tggattgcga 2340 aaactgggaa gaagacactc catttaaaga tccgcgcgag ctgtatgatt ttttaaagac 2400 ggaaaagccc gaagaggaac ttgtcttttc ccacggcgac ctgggagaca gcaacatctt 2460 tgtgaaagat ggcaaagtaa gtggctttat tgatcttggg agaagcggca gggcggacaa 2520 gtggtatgac attgccttct gcgtccggtc gatcagggag gatatcgggg aagaacagta 2580 tgtcgagcta ttttttgact tactggggat caagcctgat tgggagaaaa taaaatatta 2640 tattttactg gatgaattgt tttagtacct agatttagat gtctaaaaag ctttaactac 2700 aagettttta gacatetaat ettttetgaa gtacateege aactgtecat actetgatgt 2760 tttatatett ttetaaaagt tegetagata ggggteeega gegeetaega ggaatttgta 2820 tegecatteg ceatteagge tgegeaactg ttgggaaggg egateggtge ggtegaetgg 2880 caggeaaaac aggaeccaag gteattgega caggaggeet ggegeegete attgegaaeg 2940 aatcagattg tatagacatc gttgatccat tcttaaccct aaaagggctg gaattgattt 3000 atgaaagaaa ccgcgtagga agtgtatagg aggtttagta atggattatt tagtaaaagc 3060 acttgcgtat gacggaaaag ttcgggctta tgcagcgaga acgactgata tggtaaatga 3120 ggggcagaga cgccatggta cgtggccgac agcatccgct gcactaggcc gtacaatgac 3180 agetteactt atgeteggeg etatgetgaa gggegatgat aagetgaeeg tgaaaatega 3240 gggcggaggt ccgatcggag ctattgtagc tgatgccaat gccaaaggag aagtcagagc 3300 ctatgtctct aacccgcaag ttcattttga tttaaatgaa caaggtaagc ttgatgtcag 3360 acgtgcggtt ggaacaaacg gaacgttaag tgtcgtaaaa gatttaggtt tgcgcgagtt 3420 cttcacagga caagtagaaa tcgtttcagg agaattagga gatgatttta cttactatct 3480 tgtgtcatct gagcaggttc cttcatcagt gggcgtaggt gtgctcgtaa atcctgacaa 3540 taccattett geggeagggg getttattat teagetgatg eegggaacag atgatgaaac 3600 aatcacaaaa attgaacagc gtctatctca agtagagccg atttctaagc tcatccaaaa 3660 agggctgaca ccagaagaaa ttttagaaga agtcctaggc gagaaacctg agattttgga 3720 aacgatgcct gtcagattcc attgcccttg ttcaaaagaa cggttcgaaa cagccatttt 3780 aggactaggc aaaaaagaaa ttcaagatat gatagaagaa gatggacaag ccgaagcagt 3840 atgccatttt tgtaatgaaa agtacttatt tacaaaagaa gagctggaag ggcttcgtga 3900 ccaaactacc cgctaagctc tttagcgggt ttttaatttg agaaaagggg ctgaaagcag 3960 gtttgaaatc aagaacaatc tggacgcgtt ggatgcatag cttgagtatt ctatagtgtc 4020 acctaaatag cttggcgtaa tcatggtcat agctgtttcc tgtgtgaaat tgttatccgc 4080 tcacaattcc acacaacata cgagccggaa gcataaagtg taaagcctgg ggtgcctaat 4140 gagtgageta acteacatta attgegttge geteactgee egettteeag tegggaaace 4200 tgtcgtgcca gctgcattaa tgaatcggcc aacgcgcggg gagaggcggt ttgcgtattg 4260 ggegetette egetteeteg eteactgaet egetgegete ggtegttegg etgeggegag 4320 cggtatcagc tcactcaaag gcggtaatac ggttatccac agaatcaggg gataacgcag 4380 gaaagaacat gtgagcaaaa ggccagcaaa aggccaggaa ccgtaaaaag gccgcgttgc 4440 tggcgttttt cgataggctc cgccccctg acgagcatca caaaaatcga cgctcaagtc 4500 agaggtggcg aaacccgaca ggactataaa gataccaggc gtttccccct ggaagctccc 4560 tegtgegete teetgtteeg accetgeege ttaceggata cetgteegee ttteteeett 4620 cgggaagcgt ggcgctttct catagctcac gctgtaggta tctcagttcg gtgtaggtcg 4680 ttcgctccaa gctgggctgt gtgcacgaac cccccgttca gcccgaccgc tgcgccttat 4740 ccggtaacta tcgtcttgag tccaacccgg taagacacga cttatcgcca ctggcagcag 4800 ccactggtaa caggattagc agagcgaggt atgtaggcgg tgctacagag ttcttgaagt 4860 ggtggcctaa ctacggctac actagaagga cagtatttgg tatctgcgct ctgctgaagc 4920 cagttacctt cggaaaaaga gttggtagct cttgatccgg caaacaaacc accgctggta 4980 gcggtggttt ttttgtttgc aagcagcaga ttacgcgcag aaaaaaagga tctcaagaag 5040 atcetttgat ettttetaeg gggtetgaeg eteagtggaa egaaaaetea egttaaggga 5100 ttttggtcat gagattatca aaaaggatct tcacctagat ccttttaaat taaaaatgaa 5160 gttttaaatc aatctaaagt atatatgagt aaacttggtc tgacagttac caatgcttaa 5220 tcagtgaggc acctatctca gcgatctgtc tatttcgttc atccatagtt gcctgactcc 5280 ccgtcgtgta gataactacg atacgggagg gcttaccatc tggccccagt gctgcaatga 5340 taccgcgaga cccacgctca ccggctccag atttatcagc aataaaccag ccagccggaa 5400 gggccgagcg cagaagtggt cctgcaactt tatccgcctc catccagtct attaattgtt 5460 geegggaage tagagtaagt agttegeeag ttaatagttt gegeaaegtt gttggeattg 5520 ctacaggcat cgtggtgtca cgctcgtcgt ttggtatggc ttcattcagc tccggttccc 5580 aacgatcaag gcgagttaca tgatccccca tgttgtgcaa aaaagcggtt agctccttcg 5640 gtcctccgat cgttgtcaga agtaagttgg ccgcagtgtt atcactcatg gttatggcag 5700 cactgcataa ttctcttact gtcatgccat ccgtaagatg cttttctgtg actggtgagt 5760 actcaaccaa gtcattctga gaataccgcg cccggcgacc gagttgctct tgcccggcgt 5820 caatacggga taatagtgta tgacatagca gaactttaaa agtgctcatc attggaaaac 5880 gttcttcggg gcgaaaactc tcaaggatct taccgctgtt gagatccagt tcgatgtaac 5940 ccactcgtgc acccaactga tcttcagcat cttttacttt caccagcgtt tctgggtgag 6000 caaaaacagg aaggcaaaat gccgcaaaaa agggaataag ggcgacacgg aaatgttgaa 6060 tactcatact cttccttttt caatattatt gaagcattta tcagggttat tgtctcatga 6120 gcggatacat atttgaatgt atttagaaaa ataaacaaat aggggttccg cgcacatttc 6180 cccgaaaagt gccacctgta tgcggtgtga aataccgcac agatgcgtaa ggagaaaata 6240 ccgcatcagg cgaaattgta aacgttaata ttttgttaaa attcgcgtta aatatttgtt 6300 aaatcagctc attttttaac caataggccg aaatcggcaa aatcccttat aaatcaaaag 6360 aatagaccga gatagggttg agtgttgttc cagtttggaa caagagtcca ctattaaaga 6420 acgtggactc caacgtcaaa gggcgaaaaa ccgtctatca gggcgatggc ccactacgtg 6480 aaccatcacc caaatcaagt tttttgcggt cgaggtgccg taaagctcta aatcggaacc 6540 ctaaagggag cccccgattt agagcttgac ggggaaagcc ggcgaacgtg gcgagaaagg 6600 aagggaagaa agcgaaagga gcgggcgcta gggcgctggc aagtgtagcg gtcacgctgc 6660 6688 gcgtaaccac cacacccgcc gcgcttaa

<210> 93

<211> 8503

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant

pAN004 plasmid

<400> 93 gaattttgcg gccgcttcga aagctgtaat ataaaaacct tcttcaacta acggggcagg 60 ttagtgacat tagaaaaccg actgtaaaaa gtacagtcgg cattatctca tattataaaa 120 gccagtcatt aggcctatct gacaattcct gaatagagtt cataaacaat cctgcatgat 180 aaccatcaca aacagaatga tgtacctgta aagatagcgg taaatatatt gaattacctt 240 tattaatgaa ttttcctgct gtaataatgg gtagaaggta attactatta ttattgatat 300 ttaagttaaa cccagtaaat gaagtccatg gaataataga aagagaaaaa gcattttcag 360 gtataggtgt tttgggaaac aatttccccg aaccattata tttctctaca tcagaaaggt 420 ataaatcata aaactctttg aagtcattct ttacaggagt ccaaatacca gagaatgttt 480 tagatacacc atcaaaaatt gtataaagtg gctctaactt atcccaataa cctaactctc 540 cgtcgctatt gtaaccagtt ctaaaagctg tatttgagtt tatcaccctt gtcactaaga 600 aaataaatgc agggtaaaat ttatatcctt cttgttttat gtttcggtat aaaacactaa 660 tatcaatttc tgtggttata ctaaaagtcg tttgttggtt caaataatga ttaaatatct 720 cttttctctt ccaattgtct aaatcaattt tattaaagtt catttgatat gcctcctaaa 780 tttttatcta aagtgaattt aggaggetta ettgtetget ttetteatta gaateaatee 840 ttttttaaaa gtcaatatta ctgtaacata aatatatatt ttaaaaatat cccactttat 900 ccaattttcg tttgttgaac taatgggtgc tttagttgaa gaataaagac cacattaaaa 960 aatgtggtet titgtgtttt titaaaggat tigagegtag egaaaaatee tittettet 1020 tatettgata ataagggtaa etattgaatt eggtaceaag agtttgtaga aaegeaaaaa 1080 ggccatccgt caggatggcc ttctgcttaa tttgatgcct ggcagtttat ggcgggcgtc 1140 ctgcccgcca ccctccgggc cgttgcttcg caacgttcaa atccgctccc ggcggatttg 1200 tectaeteag gagagegtte accgacaaac aacagataaa acgaaaggee cagtettteg 1260 actgagcctt tcgttttatt tgatgcctgg cagttcccta ctctcgcatg gggagacccc 1320 acactaccat eggegetacg gegttteact tetgagtteg geatggggte aggtgggace 1380 accgcgctac tgccgccagg caaattctgt tttatcagac cgcttctgcg ttctgattta 1440 atctgtatca ggctgaaaat cttctctcat ccgccaaaac aggatcctac ggaaatggag 1500 cggcaaaacc gttttactct caaaatctta aaagaaaacc cccgataaag ggggcttttc 1560 ttctacaaaa ttgtacgggc tggttcgttc cccagcattt gttcaatttt gttttgatca 1620 ttcagaacag ccactttcgg ctcatggctt gccgcttctt gatcagacat cattttgtag 1680 gaaataataa tgaccttatc teetteetge acaaggegtg eggetgeace gtttaageat 1740 atgacgccgc ttccccgttt accaggaata atatacgttt caagacgtgc tccattatta 1800 ttattcacaa tttgtacttt ttcattagga agcattccca cagcatcaat gagatcttca 1860 tcaattgtaa tgcttcccac atagttcagg tttgcttccg taacagttgc cctgtgaagt 1920 ttgccgctca tcattgttcg atacatatta tattctctcc atttctcgaa tatcaataat 1980 gatattatct attaaacgcg cttttgaaaa agcaactgca acagcgagaa tcatctttcc 2040 agcaatttca ttcacaggct cgagttccgg ataggaataa agctctacat agtctatggt 2100 tccgctagtc gtttcaatga tatcttttgc agcttttatc accgcttcag gatctctttc 2160 accggcttgg acaagttccg cacttgtttg aagggcccga tacagcttag gcgcttcttt 2220 tettteetea getgttaagt atacattgeg agagettttg getaageegt etteetetet 2280 gaccgtatcg acaggaacca attcaatatc catgaagaag tcgctgatta acccatcaac 2340 aacagctacc tgctgcgcat cttttaaacc gaaataggca cgagtcggct tgactagatt 2400 gaaaagette gteagtaega tegegaeeee gteaaaatgt eettetettg agegeeegea 2460 taacacgtct gtgcgtcttt ctacatgaat cgtgacattc ttttcaccgg gatacatatc 2520 atgagcatct ggcgtaaaaa gaatatcgac tccggcgttt tctgcaagag ctgcatcccg 2580 ctcaatatcg cgcggatatg cttcaaaatc ttcattaggg ccgaattgtg caggattcac 2640 aaaaatactc ataataacgg cgtcgttttc ttgtcttgct ttgtctgcta aggttaaatg 2700 cccctcatgc agaaacccca tcgtcggaac aaatccgatt gacttgccct ctgaatggta 2760 ttgttttatg gcttctttca gctgtgaaat atcagtaatc tgtctcatct tattttcccc 2820 cgtacaagcc gtcaagcact gtctggttca tttgaaagga atgcttttgt tcagggaaag 2880 cacgatgtct tacatcctga acatatccgc tgattgctgt ttcgatggtt tcatcaatgc 2940 gcgtatattg ctttacaaat ttaggtgttc tctcaacacc gtggccgata atatcatgat 3000 aaacgagaac ttgtccgtcc gctttcacac cagccccgat tccaatgacc ggtatgctta 3060 gegtetegge aattttgget gtgagttetg eeggeacaca tteeageaca ageateatag 3120 ctcctgcttc ttcgcatttt atactgtctt ctattaattt tttggcgctt tgttcgtctt 3180 tgccctgtac tttatagccg cccagtacgc cgactgactg cggtgtcaaa cctaagtgac 3240 tgactactgg aatgcctcca agcgtcaatg cgcgaatgga ttcaaacacg ccttctccgc 3300 cctcaagctt cagtgcgtca gctccgcttt cctgaacgat agccgctgca tttttcagcg 3360 tatcttcctt agacaggtga taagacataa acggcatatc tgtcacaata aaggtattcg 3420 gcgcacccct tttaacggct tttgtatgat ggatcatgtc cgcaactgtc acaccgacag 3480 ttgaatcaag gccgaggacg accattccaa gtgaatcacc gactaaaatc atgtcaactc 3540 ccgcttgttc agcaagttta gctgccggat aatcataagc ggtcagcatg acaatcggtt 3600 gagcgtcctg ctgttgttaa gattattata ccacaccttg tagataaagt caacaacttt 3720 ttgcaaaatt tttcaggaat tttagcagag gttgttctgg atgtagaaca aaacatcttt 3780 ccgctcttgt gctgttagga tatctttctt ggaagctagg taggcctcga gttatggcag 3840 ttggttaaaa ggaaacaaaa agaccgtttt cacacaaaac ggtctttttc gatttctttt 3900 tacagtcaca gccacttttg caaaaaccgg acagettcat geettataac tgctgtttcg 3960 gtcgacaagc ttcgcgaagc ggccgcaaaa ttcactggcc gtcgttttac aacgtcgtga 4020 ctgggaaaac cctggcgtta cccaacttaa tcgccttgca gcacatcccc ctttcgccag 4080 ctggcgtaat agcgaagagg cccgcaccga tcgcccttcc caacagttgc gcagcctgaa 4140 tggcgaatgg cgcctgatgc ggtattttct ccttacgcat ctgtgcggta tttcacaccg 4200 catatggtgc actctcagta caatctgctc tgatgccgca tagttaagcc agccccgaca 4260 cccgccaaca cccgctgact atgcttgtaa accgttttgt gaaaaaattt ttaaaataaa 4320 aaaggggacc tctagggtcc ccaattaatt agtaatataa tctattaaag gtcattcaaa 4380 aggtcatcca ccggatcagc ttagtaaagc cctcgctaga ttttaatgcg gatgttgcga 4440 ttacttcgcc aactattgcg ataacaagaa aaagccagcc tttcatgata tatctcccaa 4500 tttgtgtagg gcttattatg cacgcttaaa aataataaaa gcagacttga cctgatagtt 4560 tggctgtgag caattatgtg cttagtgcat ctaacgcttg agttaagccg cgccgcgaag 4620 cggcgtcggc ttgaacgaat tgttagacat tatttgccga ctaccttggt gatctcgcct 4680 ttcacgtagt ggacaaattc ttccaactga tctgcgcgcg aggccaagcg atcttcttct 4740 tgtccaagat aagcctgtct agcttcaagt atgacgggct gatactgggc cggcaggcgc 4800 tecattgeee agteggeage gaeateette ggegegattt tgeeggttae tgegetgtae 4860 caaatgcggg acaacgtaag cactacattt cgctcatcgc cagcccagtc gggcggcgag 4920 ttccatagcg ttaaggtttc atttagcgcc tcaaatagat cctgttcagg aaccggatca 4980 aagagtteet eegeegetgg acetaceaag geaacgetat gttetettge ttttgteage 5040 aagatagcca gatcaatgtc gatcgtggct ggctcgaaga tacctgcaag aatgtcattg 5100 cgctgccatt ctccaaattg cagttcgcgc ttagctggat aacgccacgg aatgatgtcg 5160 tegtgeacaa caatggtgae ttetacageg eggagaatet egetetetee aggggaagee 5220 gaagttteca aaaggtegtt gateaaaget egeegegttg ttteateaag eettaeggte 5280 accgtaacca gcaaatcaat atcactgtgt ggcttcaggc cgccatccac tgcggagccg 5340 tacaaatgta eggeeageaa egteggtteg agatggeget egatgaegee aactaeetet 5400 gatagttgag tcgatacttc ggcgatcacc gcttccctca tgatgtttaa ctttgtttta 5460 gggcgactgc cctgctgcgt aacatcgttg ctgctccata acatcaaaca tcgacccacg 5520 gcgtaacgcg cttgctgctt ggatgcccga ggcatagact gtaccccaaa aaaacagtca 5580 taacaagcca tgaaaaccgc cactgcgccg ttaccaccgc tgcgttcggt caaggttctg 5640 qaccagttgc gtgagcgcat acgctacttg cattacagct tacgaaccga acaggcttat 5700 gtccactggg ttcgtgcctt catccgtttc cacggtgtgc gtcacccggc aaccttgggc 5760 agcageqaaq tegaggeatt tetgteetgg etggegaaeg agegeaaggt tteggtetee 5820 acqcatcqtc aqqcattqqc qqccttqctg ttcttctacg gcaaggtgct gtgcacggat 5880 ctgccctggc ttcaggagat cggaagacct cggccgtcgc ggcgcttgcc ggtggtgctg 5940 accordgatg aagtggttcg catcotcggt tttctggaag gcgagcatcg tttgttcgcc 6000 cagettetgt atggaacggg catgeggate agtgagggtt tgcaactgeg ggtcaaggat 6060 ctqqatttcq atcacggcac gatcatcgtg cgggagggca agggctccaa ggatcgggcc 6120 ttgatgttac ccgagagctt ggcacccagc ctgcgcgagc aggggaattg atccggtgga 6180 tgaccttttg aatgaccttt aatagattat attactaatt aattggggac cctagaggtc 6240 ccctttttta ttttaaaaat tttttcacaa aacggtttac aagcataacg ggttttgctg 6300 cccqcaaacq ggctgttctg gtgttgctag tttgttatca gaatcgcaga tccggcttca 6360 ggtttgccgg ctgaaagcgc tatttcttcc agaattgcca tgattttttc cccacgggag 6420 gegteactgg etecegtgtt gteggeaget ttgattegat aageageate geetgtttea 6480 ggctgtctat gtgtgactgt tgagctgtaa caagttgtct caggtgttca atttcatgtt 6540 ctagttgctt tgttttactg gtttcacctg ttctattagg tgttacatgc tgttcatctg 6600 ttacattqtc gatctqttca tggtgaacag ctttaaatgc accaaaaact cgtaaaagct 6660 ctgatgtate tatettttt acaccgtttt catetgtgca tatggacagt tttccctttg 6720 atatctaacg gtgaacagtt gttctacttt tgtttgttag tcttgatgct tcactgatag 6780 atacaagagc cataagaacc tcagatcctt ccgtatttag ccagtatgtt ctctagtgtg 6840 gttcgttgtt tttgcgtgag ccatgagaac gaaccattga gatcatgctt actttgcatg 6900 tcactcaaaa attttgcctc aaaactggtg agctgaattt ttgcagttaa agcatcgtgt 6960 agtqtttttc ttagtccgtt acgtaggtag gaatctgatg taatggttgt tggtattttg 7020 tcaccattca tttttatctg gttgttctca agttcggtta cgagatccat ttgtctatct 7080 agttcaactt ggaaaatcaa cgtatcagtc gggcggcctc gcttatcaac caccaatttc 7140 atattgctgt aagtgtttaa atctttactt attggtttca aaacccattg gttaagcctt 7200 ttaaactcat ggtagttatt ttcaagcatt aacatgaact taaattcatc aaggctaatc 7260 tctatatttg ccttgtgagt tttcttttgt gttagttctt ttaataacca ctcataaatc 7320 ctcatagagt atttgttttc aaaagactta acatgttcca gattatattt tatgaatttt 7380 tttaactgga aaagataagg caatatetet teactaaaaa etaattetaa tttttegett 7440 gagaacttgg catagtttgt ccactggaaa atctcaaagc ctttaaccaa aggattcctg 7500 atttccacag ttctcgtcat cagctctctg gttgctttag ctaatacacc ataagcattt 7560 tecetactga tgtteateat etgagegtat tggttataag tgaaegatae egteegttet 7620 ttccttgtag ggttttcaat cgtggggttg agtagtgcca cacagcataa aattagcttg 7680 qtttcatgct ccgttaagtc atagcgacta atcgctagtt catttgcttt gaaaacaact 7740 aattcagaca tacatctcaa ttggtctagg tgattttaat cactatacca attgagatgg 7800 gctagtcaat gataattact agtccttttc ctttgagttg tgggtatctg taaattctgc 7860 tagacetttg etggaaaact tgtaaattet getagaceet etgtaaatte egetagacet 7920 ttgtgtgttt tttttgttta tattcaagtg gttataattt atagaataaa gaaagaataa 7980 aaaaagataa aaagaataga tcccagccct gtgtataact cactacttta gtcagttccg 8040 cagtattaca aaaggatgtc gcaaacgctg tttgctcctc tacaaaacag accttaaaac 8100 cctaaagget taagtageae eetegeaage tegggeaaat egetgaatat teettttgte 8160 teegaeeate aggeaeetga gtegetgtet tittegtgae atteagtieg etgegeteae 8220 ggctctggca gtgaatgggg gtaaatggca ctacaggcgc cttttatgga ttcatgcaag 8280 gaaactaccc ataatacaag aaaagcccgt cacgggcttc tcagggcgtt ttatggcggg 8340 tetgetatgt ggtgetatet gaetttttge tgtteageag tteetgeeet etgattttee 8400 agtetgacea etteggatta tecegtgaca ggteatteag aetggetaat geaceeagta 8460 8503 aggcagcggt atcatcaaca ggcttacccg tcttactgtc aac

<210> 94

<211> 7381

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:recombinant
pAN006 plasmid

<400> 94

ttgcggccgc ttcgaaagct gtaatataaa aaccttcttc aactaacggg gcaggttagt 60 gacattagaa aaccgactgt aaaaagtaca gtcggcatta tctcatatta taaaagccag 120 tcattaggcc tatctgacaa ttcctgaata gagttcataa acaatcctgc atgataacca 180 tcacaaacag aatgatgtac ctgtaaagat agcggtaaat atattgaatt acctttatta 240

BGI-141CP - 138 -

atgaattttc ctgctgtaat aatgggtaga aggtaattac tattattatt gatatttaag 300 ttaaacccag taaatgaagt ccatggaata atagaaagag aaaaagcatt ttcaggtata 360 ggtgttttgg gaaacaattt ccccgaacca ttatatttct ctacatcaga aaggtataaa 420 tcataaaact ctttgaagtc attctttaca ggagtccaaa taccagagaa tgttttagat 480 acaccatcaa aaattgtata aagtggctct aacttatccc aataacctaa ctctccgtcg 540 ctattgtaac cagttctaaa agctgtattt gagtttatca cccttgtcac taagaaaata 600 aatgcagggt aaaatttata teettettgt tttatgttte ggtataaaae actaatatea 660 atttctgtgg ttatactaaa agtcgtttgt tggttcaaat aatgattaaa tatctctttt 720 ctcttccaat tgtctaaatc aattttatta aagttcattt gatatgcctc ctaaattttt 780 atctaaagtg aatttaggag gettaettgt etgetttett eattagaate aateettttt 840 taaaagtcaa tattactgta acataaatat atattttaaa aatatcccac tttatccaat 900 tttcgtttgt tgaactaatg ggtgctttag ttgaagaata aagaccacat taaaaaatgt 960 ggtcttttgt gtttttttaa aggatttgag cgtagcgaaa aatccttttc tttcttatct 1020 tgataataag ggtaactatt gaattcggta ccaagagttt gtagaaacgc aaaaaggcca 1080 teegteagga tggeettetg ettaatttga tgeetggeag tttatggegg gegteetgee 1140 egecaecete egggeegttg ettegeaacg tteaaateeg eteeeggegg atttgteeta 1200 ctcaggagag cgttcaccga caaacaacag ataaaacgaa aggcccagtc tttcgactga 1260 geetttegtt ttatttgatg eetggeagtt eeetactete geatggggag acceeacaet 1320 accateggeg ctaeggegtt teaettetga gtteggeatg gggteaggtg ggaecaeege 1380 gctactgccg ccaggcaaat tctgttttat cagaccgctt ctgcgttctg atttaatctg 1440 tatcaggctg aaaatcttct ctcatccgcc aaaacaggat cctacggaaa tggagcggca 1500 aaaccgtttt actctcaaaa tcttaaaaga aaacccccga taaagggggc ttttcttcta 1560 caaaattgta cgggctggtt cgttccccag catttgttca attttgtttt gatcattcag 1620 aacagccact ttcggctcat ggcttgccgc ttcttgatca gacatcattt tgtaggaaat 1680 aataatgacc ttatctcctt cctgcacaag gcgtgcggct gcaccgttta agcatatgac 1740 gccgcttccc cgtttaccag gaataatata cgtttcaaga cgtgctccat tattattatt 1800 cacaatttgt actttttcat taggaagcat tcccacagca tcaatgagat cttcatcaat 1860 tgtaatgctt cccacatagt tcaggtttgc ttccgtaaca gttgccctgt gaagtttgcc 1920 gctcatcatt gttcgataca tattatattc tctccatttc tcgaatatca ataatgatat 1980 tatctattaa acgcgctttt gaaaaagcaa ctgcaacagc gagaatcatc tttccagcaa 2040

tttcattcac aggetegagt teeggatagg aataaagete tacatagtet atggtteege 2100 tagtcgtttc aatgatatct tttgcagctt ttatcaccgc ttcaggatct ctttcaccgg 2160 cctcagctgt taagtataca ttgcgagagc ttttggctaa gccgtcttcc tctctgaccg 2280 tatcgacagg aaccaattca atatccatga agaagtcgct gattaaccca tcaacaacag 2340 ctacctgctg cgcatctttt aaaccgaaat aggcacgagt cggcttgact agattgaaaa 2400 gcttcgtcag tacgatcgcg accccgtcaa aatgtccttc tcttgagcgc ccgcataaca 2460 cgtctgtgcg tctttctaca tgaatcgtga cattcttttc accgggatac atatcatgag 2520 catctggcgt aaaaagaata tcgactccgg cgttttctgc aagagctgca tcccgctcaa 2580 tatcgcgcgg atatgcttca aaatcttcat tagggccgaa ttgtgcagga ttcacaaaaa 2640 tactcataat aacggcgtcg ttttcttgtc ttgctttgtc tgctaaggtt aaatgcccct 2700 catgcagaaa ccccatcgtc ggaacaaatc cgattgactt gccctctgaa tggtattgtt 2760 ttatggette tttcagetgt gaaatateag taatetgtet eatettattt teeceegtae 2820 aagcegteaa geactgtetg gtteatttga aaggaatget tttgtteagg gaaageaega 2880 tgtcttacat cctgaacata tccgctgatt gctgtttcga tggtttcatc aatgcgcgta 2940 tattgcttta caaatttagg tgttctctca acaccgtggc cgataatatc atgataaacg 3000 agaacttgtc cgtccgcttt cacaccagcc ccgattccaa tgaccggtat gcttagcgtc 3060 teggeaattt tggetgtgag ttetgeegge acaeatteea geacaageat catageteet 3120 gettettege attitataet giettetatt aattitiigg egettigite giettigeee 3180 tgtactttat agccgcccag tacgccgact gactgcggtg tcaaacctaa gtgactgact 3240 actggaatgc ctccaagcgt caatgcgcga atggattcaa acacgccttc tccgccctca 3300 agetteagtg egteagetee gettteetga aegatageeg etgeattttt eagegtatet 3360 tecttagaca ggtgataaga cataaacgge atatetgtea caataaaggt atteggegea 3420 ccccttttaa cggcttttgt atgatggatc atgtccgcaa ctgtcacacc gacagttgaa 3480 tcaaggccga ggacgaccat tccaagtgaa tcaccgacta aaatcatgtc aactcccgct 3540 tgttcagcaa gtttagctgc cggataatca taagcggtca gcatgacaat cggttcttca 3600 gacteettea tittitagaaa ateeagtiit giitteatgi titeteetee tetagagegi 3660 cctgctgttg ttaagattat tataccacac cttgtagata aagtcaacaa ctttttgcaa 3720 aatttttcag gaattttagc agaggttgtt ctggatgtag aacaaaacat ctttccqctc 3780 ttgtgctgtt aggatatctt tcttggaagc taggtaggcc tcgagttatg gcagttggtt 3840 aaaaggaaac aaaaagaccg ttttcacaca aaacggtctt tttcgatttc tttttacagt 3900 cacagocact tttgcaaaaa coggacagot tcatgootta taactgotgt ttcqqtcqac 3960 ctgcaggcat gcaagcttcg cgaagcggcc gccgacgcga ggctggatgg ccttccccat 4020 tatgattett etegetteeg geggeategg gatgeeegeg ttgeaggeea tgetgteeag 4080 geaggtagat gacgaceate agggaeaget teaaggateg etegeggete ttaccageet 4140 aacttegate actggacege tgategteae ggegatttat geegeetegg egageaeatg 4200 gaacgggttg gcatggattg taggcgccgc cctatacctt gtctgcctcc ccgcgttgcg 4260 tegeggtgea tggageeggg ceacetegae etgaatggaa geeggeggea eetegetaae 4320 ggattcacca ctccaagaat tggagccaat caattcttgc ggagaactgt gaatgcgcaa 4380 accaaccett ggcagaacat atccategeg teegecatet eeageageeg caegeggege 4440 atctegggea gegttgggte etggeeaegg gtgegeatga tegtgeteet gtegttgagg 4500 acceggetag getggegggg ttgeettaet ggttageaga atgaateace gataegegag 4560 cgaacgtgaa gcgactgctg ctgcaaaacg tctgcgacct gagcaacaac atgaatggtc 4620 tteggtttee gtgtttegta aagtetggaa aegeggaagt eagegeeetg eaceattatg 4680 ttccggatct gcatcgcagg atgctgctgg ctaccctgtg gaacacctac atctgtatta 4740 acgaageget ggeattgace etgagtgatt tttetetggt eeegeegeat ceatacegee 4800 agttgtttac cctcacaacg ttccagtaac cgggcatgtt catcatcagt aacccgtatc 4860 gtgagcatcc tetetegttt categgtate attacececa tgaacagaaa tteeceetta 4920 cacggaggca tcaagtgacc aaacaggaaa aaaccgccct taacatggcc cgctttatca 4980 gaagccagac attaacgctt ctggagaaac tcaacgagct ggacgcggat gaacaggcag 5040 acatetytya ategetteae gaceaegety atgagettta eegeagetye etegegegtt 5100 teggtgatga eggtgaaaac etetgaeaca tgeageteee ggagaeggte acagettgte 5160 tgtaagcgga tgccgggagc agacaagccc gtcagggcgc gtcagcgggt gttggcgggt 5220 gtcggggcgc agccatgacc cagtcacgta gcgatagcgg agtgtatact ggcttaacta 5280 tgcggcatca gagcagattg tactgagagt gcaccatatg cggtgtgaaa taccgcacag 5340 atgegtaagg agaaaataee geateaggeg etetteeget teetegetea etgaeteget 5400 gegeteggte gtteggetge ggegageggt ateageteae teaaaggegg taataeggtt 5460 atccacagaa tcaggggata acgcaggaaa gaacatgtga gcaaaaaggcc agcaaaaggc 5520 caggaaccgt aaaaaggccg cgttgctggc gtttttccat aggctccgcc cccctgacga 5580 gcatcacaaa aatcgacgct caagtcagag gtggcgaaac ccgacaggac tataaagata 5640 ccaggogttt ccccctggaa getecctegt gegeteteet gtteegacee tgeegettae 5700

eggatacetg teegeettte teeetteggg aagegtggeg ettteteata geteaegetg 5760 taggtatete agtteggtgt aggtegtteg etecaagetg ggetgtgtge aegaaceee 5820 cgttcagccc gaccgctgcg ccttatccgg taactatcgt cttgagtcca acccgqtaag 5880 acacgactta tegecactgg cageageeac tggtaacagg attageagag egaggtatgt 5940 aggeggtget acagagttet tgaagtggtg geetaactae ggetacacta gaaggacagt 6000 atttggtatc tgcgctctgc tgaagccagt taccttcgga aaaagagttg gtagctcttg 6060 atccggcaaa caaaccaccg ctggtagcgg tggttttttt gtttgcaagc agcagattac 6120 gcgcagaaaa aaaggatctc aagaagatcc tttgatcttt tctacggggt ctgacgctca 6180 gtggaacgaa aactcacgtt aagggatttt ggtcatgaga ttatcaaaaa ggatcttcac 6240 ctagatcctt ttaaattaaa aatgaagttt taaatcaatc taaagtatat atgagtaaac 6300 ttggtctgac agttaccaat gcttaatcag tgaggcacct atctcagcga tctgtctatt 6360 tegtteatee atagttgeet gaeteeegt egtgtagata actaegatae gggagggett 6420 accatctggc cccagtgctg caatgatacc gcgagaccca cgctcaccgg ctccagattt 6480 atcagcaata aaccagccag ccggaagggc cgagcgcaga agtggtcctg caactttatc 6540 cgcctccatc cagtctatta attgttgccg ggaagctaga gtaagtagtt cgccagttaa 6600 tagtttgcgc aacgttgttg ccattgctgc aggcatcgtg gtgtcacgct cgtcgtttgg 6660 tatggcttca ttcagctccg gttcccaacg atcaaggcga gttacatgat cccccatgtt 6720 gtgcaaaaaa geggttaget eetteggtee teegategtt gteagaagta agttggeege 6780 agtgttatca ctcatggtta tggcagcact gcataattct cttactgtca tgccatccgt 6840 aagatgettt tetgtgactg gtgagtaete aaccaagtea ttetgagaat agtgtatgeg 6900 gcgaccgagt tgctcttgcc cggcgtcaat acgggataat accgcgccac atagcagaac 6960 tttaaaagtg ctcatcattg gaaaacgttc ttcggggcga aaactctcaa ggatcttacc 7020 getgttgaga tecagttega tgtaacceae tegtgeaece aactgatett cageatettt 7080 tactttcacc agcgtttctg ggtgagcaaa aacaggaagg caaaatgccg caaaaaaggg 7140 aataagggcg acacggaaat gttgaatact catactcttc ctttttcaat attattgaag 7200 catttatcag ggttattgtc tcatgagcgg atacatattt gaatgtattt agaaaaataa 7260 acaaataggg gttccgcgca catttccccg aaaagtgcca cctgacgtct aagaaaccat 7320 tattatcatg acattaacct ataaaaatag gcgtatcacg aggccctttc gtcttcaaga 7380 7381 a

United States Patent & Trademark Office

Office of Initial Patent Examination -- Scanning Division



Application deficiencies were found during scanning:

Page(s)	of Transmi	Ha /	were not present
for scanning.	(Document title)		
	6		
\square Page(s)	of		were not present
for scanning.	(D	ocument title)	

☐ Scanned copy is best available.